

# Saranya Khunkhet

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2379401/publications.pdf>

Version: 2024-02-01

13  
papers

277  
citations

1307594

7  
h-index

1199594

12  
g-index

13  
all docs

13  
docs citations

13  
times ranked

284  
citing authors

| #  | ARTICLE   | IF  | CITATIONS |
|----|---|-----|-----------|
| 1  | Triple combination therapy of narrowband ultraviolet B, fractional carbon dioxide laser and topical bimatoprost 0.01% for non-segmental vitiligo on non-facial areas: A randomized half-body, double-blind, placebo-controlled, comparative study. <i>Dermatologic Therapy</i> , 2022, 35, e15198.                    | 1.7 | 2         |
| 2  | Minimum Contact Time of 1.25%, 2.5%, 5%, and 10% Benzoyl Peroxide for a Bactericidal Effect Against <i>Cutibacterium acnes</i> . <i>Clinical, Cosmetic and Investigational Dermatology</i> , 2022, Volume 15, 403-409.  | 1.8 | 4         |
| 3  | Histopathological Evidence of Occipital Involvement in Male Androgenetic Alopecia. <i>Frontiers in Medicine</i> , 2021, 8, 790597.  | 2.6 | 8         |
| 4  | Frontal fibrosing alopecia in Asians: a retrospective clinical study. <i>International Journal of Dermatology</i> , 2020, 59, 184-190.  | 1.0 | 11        |
| 5  | Alopecia Induced by Autologous Fat Injection into the Temporal Area: Case Report and Review of the Literature. <i>Case Reports in Dermatology</i> , 2019, 11, 150-156.  | 0.8 | 12        |
| 6  | A randomized, investigator-blinded, controlled, split-scalp study of the efficacy and safety of a 1550-nm fractional erbium-glass laser, used in combination with topical 5% minoxidil versus 5% minoxidil alone, for the treatment of androgenetic alopecia. <i>Lasers in Medical Science</i> , 2019, 34, 1857-1864. | 2.1 | 36        |
| 7  | Low-level laser therapy for the treatment of androgenetic alopecia in Thai men and women: a 24-week, randomized, double-blind, sham device-controlled trial. <i>Lasers in Medical Science</i> , 2019, 34, 1107-1114.  | 2.1 | 63        |
| 8  | A randomized, double-blind controlled study of the efficacy and safety of topical solution of 0.25% finasteride admixed with 3% minoxidil vs. 3% minoxidil solution in the treatment of male androgenetic alopecia. <i>Journal of the European Academy of Dermatology and Venereology</i> , 2018, 32, 2257-2263.      | 2.4 | 64        |
| 9  | Frontal fibrosing alopecia in Asians: A retrospective clinical study. <i>Journal of the American Academy of Dermatology</i> , 2018, 79, AB144.  | 1.2 | 0         |
| 10 | Trichoscopic clues for diagnosis of alopecia areata and trichotillomania in Asians. <i>International Journal of Dermatology</i> , 2017, 56, 161-165.  | 1.0 | 36        |
| 11 | A retrospective comparative study of the efficacy and safety of two regimens of diphenylcyclopropanone in the treatment of recalcitrant alopecia areata. <i>Dermatology Reports</i> , 2017, 9, 7399.  | 0.8 | 33        |
| 12 | Pythiosis presenting with digital gangrene and subcutaneous nodules mimicking medium vessel vasculitis. <i>JAAD Case Reports</i> , 2015, 1, 399-402.  | 0.8 | 6         |
| 13 | The Possible Role of Contact Sensitization to Fragrances and Preservatives in Poikiloderma of Civatte. <i>Case Reports in Dermatology</i> , 2014, 6, 258-263.   | 0.8 | 2         |