GülÅÄh Saç

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/2388716/publications.pdf

Version: 2024-02-01

20 173
papers citations

1684188 1199594 12
h-index g-index

20 20 docs citations

20 times ranked 234 citing authors

#	Article	IF	CITATIONS
1	Are introduced gibel carp <i>Carassius gibelio</i> in Turkey more invasive in artificial than in natural waters?. Fisheries Management and Ecology, 2012, 19, 178-187.	2.0	50
2	Circumstantial evidence of gibel carp, <i>Carassius gibelio, </i> reproductive competition exerted on native fish species in a mesotrophic reservoir. Fisheries Management and Ecology, 2012, 19, 167-177.	2.0	45
3	Neotype designation of Aphanius iconii, first reviser action to stabilise the usage of A. fontinalis and A. meridionalis and comments on the family group names of fishes placed in Cyprinodontidae (Teleostei: Cyprinodontiformes). Zootaxa, 2017, 4294, .	0.5	23
4	Length-weight relationships of six freshwater fishes from small streams flowing into Lake Sapanca, NW Turkey. Journal of Applied Ichthyology, 2009, 25, 230-231.	0.7	10
5	Title is missing!. Turkish Journal of Fisheries and Aquatic Sciences, 2019, 19, .	0.9	7
6	İstanbul İli (TÃ⅓rkiye) Tatlısu Balık Faunası. Turkish Journal of Bioscience and Collections, 2019, 3, 19-36	50.4	7
7	Length–weight relationship of seven freshwater fish species from Turkish Thrace. Journal of Applied Ichthyology, 2019, 35, 808-811.	0.7	5
8	Fish fauna of Yıldız Mountains (Northeastern Thrace, Turkey). Su ÜrÃ⅓nleri Dergisi, 2019, 36, 65-73.	0.3	5
9	New records of <i>Pseudophoxinus firati </i> from Turkey (Teleostei: Leuciscidae). Journal of Applied Ichthyology, 2019, 35, 769-774.	0.7	4
10	THE OCCURRENCE OF Ligula intestinalis IN ITS FISH HOST Rutilus rutilus (L.) AND THE EFFECTS OF PARASITE ON THE FISH GROWTH (BÜYÜKÇEKMECE RESERVOIR, TURKEY). Journal of Aquaculture Engineering and Fisheries Research, 0, , 142-150.	0.6	4
11	Büyükçekmece Rezervuarı (İstanbul, Türkiye)'ndaki Bazı Balık Türlerinin Boy-Ağırlık ve Boy Kondisyon Faktörleri. Journal of Limnology and Freshwater Fisheries Research, 2016, 2, 43-43.	r-Boy İliŒ	Ÿķileri ile
12	Pseudophoxinus cilicicus, a new spring minnow from southern Anatolia (Teleostei: Leuciscidae). Zootaxa, 2019, 4671, zootaxa.4671.1.8.	0.5	3
13	Relationships between Body Size, Weight and Fecundity of the Endangered Fish Alburnus carinatus Battalgil, 1941 in the Manyas Lake (Turkey). Aquatic Sciences and Engineering, 2019, 35, 27-30.	0.8	2
14	Length-weight relationship of 16 fish species belonging to genus <i>Pseudophoxinus</i> (Teleostei:) Tj ETQq0 0 0 r	gBI /Over	lock 10 Tf 50
15	Bio-Ecological Traits of Western Tubenose Goby Proterorhinus semilunaris (Heckel, 1837): A Key to Understand Its Invasion Success. Water (Switzerland), 2019, 11, 1247.	2.7	1
16	Feeding of Barbus cyclolepis Heckel, 1837 (Teleostei: Cyprinidae) and its relationship with benthic macroinvertebrate fauna in the Istranca Stream (İstanbul, Turkey). Su Ürünleri Dergisi, 2021, 38, 345-353.	0.3	1
17	A NEW LOCALITY FOR THE ENDEMIC Gobio sakaryaensis TURAN, EKMEKÇİ, LUSKOVA & MENDEL, 2012 IN THE MARMARA REGION (TURKEY) AND THE LENGTH-WEIGHT RELATIONSHIP OF THE SPECIES. Aquatic Research, 0, 36-40.		1
18	Pressure of Urbanisation on the Fish Community Structure in Küçük Menderes River Basin (Turkey). Turkish Journal of Water Science and Management, 2021, 5, 43-62.	0.2	0

#	Article	IF	CITATIONS
19	The ichthyofaunal diversity of freshwater ecosystems in Gökçeada Island (NW Turkey) under the pressure of non-native species. Turkish Journal of Zoology, 0, , .	0.9	O
20	New Localities and Length-Weight Relationship for Alburnus caeruleus in the Euphrates and Tigris River Basins (Turkey). Aquatic Sciences and Engineering, 0, , 105-109.	0.8	0