
Catherine F Kimber

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/2391897/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Development and testing of relative risk-based health messages for electronic cigarette products. Harm Reduction Journal, 2021, 18, 96.	1.3	2
2	The Time Course of Compensatory Puffing With an Electronic Cigarette: Secondary Analysis of Real-World Puffing Data With High and Low Nicotine Concentration Under Fixed and Adjustable Power Settings. Nicotine and Tobacco Research, 2021, 23, 1153-1159.	1.4	9
3	Communicating the relative health risks of E-cigarettes: An online experimental study exploring the effects of a comparative health message versus the EU nicotine addiction warnings on smokers' and non-smokers' risk perceptions and behavioural intentions. Addictive Behaviors, 2020, 101, 106177.	1.7	21
4	Daily exposure to formaldehyde and acetaldehyde and potential health risk associated with use of high and low nicotine e-liquid concentrations. Scientific Reports, 2020, 10, 6546.	1.6	11
5	Nicotine absorption from e-cigarettes over 12â€ [−] months. Addictive Behaviors, 2019, 91, 102-105.	1.7	16
6	Compensatory Puffing With Lower Nicotine Concentration E-liquids Increases Carbonyl Exposure in E-cigarette Aerosols. Nicotine and Tobacco Research, 2018, 20, 998-1003.	1.4	51
7	The effects of the European e-cigarette health warnings and comparative health messages on non-smokers' and smokers' risk perceptions and behavioural intentions. BMC Public Health, 2018, 18, 1259.	1.2	7
8	â€~Realâ€world' compensatory behaviour with low nicotine concentration eâ€liquid: subjective effects and nicotine, acrolein and formaldehyde exposure. Addiction, 2018, 113, 1874-1882.	1.7	77
9	Self-titration by experienced e-cigarette users: blood nicotine delivery and subjective effects. Psychopharmacology, 2016, 233, 2933-2941.	1.5	132
10	E-cigarette puffing patterns associated with high and low nicotine e-liquid strength: effects on toxicant and carcinogen exposure. BMC Public Health, 2016, 16, 999.	1.2	20
11	First―versus secondâ€generation electronic cigarettes: predictors of choice and effects on urge to smoke and withdrawal symptoms. Addiction, 2015, 110, 669-677.	1.7	74
12	Text messages to support e-cigarette use for smoking cessation: a tool for researchers. Qeios, 0, , .	0.0	1
13	Tailored interventions to assist smokers to stop smoking using e-cigarettes (TASSE): Study protocol. Qeios, 0, , .	0.0	2
14	Tailored interventions to assist smokers to stop smoking using e-cigarettes (TASSE): Study protocol. Qeios, 0, , .	0.0	0
15	Tailored interventions to assist smokers to stop smoking using e-cigarettes (TASSE): Study protocol. Qeios, 0, , .	0.0	2