

Ladislav Senolt

List of Publications by Citations

Source: <https://exaly.com/author-pdf/239363/ladislav-senolt-publications-by-citations.pdf>

Version: 2024-04-19

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

126
papers

5,903
citations

32
h-index

76
g-index

189
ext. papers

8,623
ext. citations

3.9
avg, IF

6.58
L-index

#	Paper	IF	Citations
126	Plasma Hsp90 levels in patients with systemic sclerosis and relation to lung and skin involvement: a cross-sectional and longitudinal study. <i>Scientific Reports</i> , 2021 , 11, 1	4.9	2785
125	The role of resistin as a regulator of inflammation: Implications for various human pathologies. <i>Clinical Immunology</i> , 2009 , 133, 157-70	9	262
124	Resistin in rheumatoid arthritis synovial tissue, synovial fluid and serum. <i>Annals of the Rheumatic Diseases</i> , 2007 , 66, 458-63	2.4	184
123	Activation of STAT3 integrates common profibrotic pathways to promote fibroblast activation and tissue fibrosis. <i>Nature Communications</i> , 2017 , 8, 1130	17.4	155
122	Association of circulating miR-223 and miR-16 with disease activity in patients with early rheumatoid arthritis. <i>Annals of the Rheumatic Diseases</i> , 2014 , 73, 1898-904	2.4	140
121	Increased adiponectin is negatively linked to the local inflammatory process in patients with rheumatoid arthritis. <i>Cytokine</i> , 2006 , 35, 247-52	4	127
120	Increased level of cytokines and matrix metalloproteinases in osteoarthritic subchondral bone. <i>Cytokine</i> , 2007 , 38, 151-6	4	125
119	Prospective new biological therapies for rheumatoid arthritis. <i>Autoimmunity Reviews</i> , 2009 , 9, 102-7	13.6	107
118	Increased serum adiponectin levels in female patients with erosive compared with non-erosive osteoarthritis. <i>Annals of the Rheumatic Diseases</i> , 2009 , 68, 295-6	2.4	96
117	Hyaluronic acid levels may have predictive value for the progression of knee osteoarthritis. <i>Osteoarthritis and Cartilage</i> , 2004 , 12, 277-83	6.2	79
116	Vaspin and omentin: new adipokines differentially regulated at the site of inflammation in rheumatoid arthritis. <i>Annals of the Rheumatic Diseases</i> , 2010 , 69, 1410-1	2.4	78
115	Pentosidine in serum and synovial fluid in patients with knee osteoarthritis and its potential role of prediction of osteoarthritis progression. <i>Arthritis Research</i> , 2003 , 5, 167		78
114	Up-regulation of metastasis-promoting S100A4 (Mts-1) in rheumatoid arthritis: putative involvement in the pathogenesis of rheumatoid arthritis. <i>Arthritis and Rheumatism</i> , 2007 , 56, 779-89		74
113	Heat shock protein 90 (Hsp90) inhibition targets canonical TGF- β signalling to prevent fibrosis. <i>Annals of the Rheumatic Diseases</i> , 2014 , 73, 1215-22	2.4	60
112	S100A4 is expressed at site of invasion in rheumatoid arthritis synovium and modulates production of matrix metalloproteinases. <i>Annals of the Rheumatic Diseases</i> , 2006 , 65, 1645-8	2.4	60
111	Increased pentosidine, an advanced glycation end product, in serum and synovial fluid from patients with knee osteoarthritis and its relation with cartilage oligomeric matrix protein. <i>Annals of the Rheumatic Diseases</i> , 2005 , 64, 886-90	2.4	59
110	Decreases in serum levels of S100A8/9 (calprotectin) correlate with improvements in total swollen joint count in patients with recent-onset rheumatoid arthritis. <i>Arthritis Research and Therapy</i> , 2011 , 13, R122	5.7	52

109	An update on biomarkers in axial spondyloarthritis. <i>Autoimmunity Reviews</i> , 2016 , 15, 501-9	13.6	49
108	Demonstration of extracellular peptidylarginine deiminase (PAD) activity in synovial fluid of patients with rheumatoid arthritis using a novel assay for citrullination of fibrinogen. <i>Arthritis Research and Therapy</i> , 2014 , 16, 498	5.7	49
107	Increasing the infliximab dose in rheumatoid arthritis patients: a randomised, double blind study failed to confirm its efficacy. <i>Annals of the Rheumatic Diseases</i> , 2009 , 68, 1285-9	2.4	49
106	The metastasis associated protein S100A4: a potential novel link to inflammation and consequent aggressive behaviour of rheumatoid arthritis synovial fibroblasts. <i>Annals of the Rheumatic Diseases</i> , 2008 , 67, 1499-504	2.4	47
105	The metastasis-associated protein S100A4 promotes the inflammatory response of mononuclear cells via the TLR4 signalling pathway in rheumatoid arthritis. <i>Rheumatology</i> , 2014 , 53, 1520-6	3.9	45
104	Low circulating Dickkopf-1 and its link with severity of spinal involvement in diffuse idiopathic skeletal hyperostosis. <i>Annals of the Rheumatic Diseases</i> , 2012 , 71, 71-4	2.4	45
103	Serum calprotectin (S100A8/9): an independent predictor of ultrasound synovitis in patients with rheumatoid arthritis. <i>Arthritis Research and Therapy</i> , 2015 , 17, 252	5.7	42
102	Pro-inflammatory effects of interleukin-35 in rheumatoid arthritis. <i>Cytokine</i> , 2015 , 73, 36-43	4	39
101	The IL-20 Cytokine Family in Rheumatoid Arthritis and Spondyloarthritis. <i>Frontiers in Immunology</i> , 2018 , 9, 2226	8.4	36
100	MicroRNA-125b: association with disease activity and the treatment response of patients with early rheumatoid arthritis. <i>Arthritis Research and Therapy</i> , 2016 , 18, 124	5.7	35
99	Serum hyaluronic acid as a potential marker with a predictive value for further radiographic progression of hand osteoarthritis. <i>Osteoarthritis and Cartilage</i> , 2009 , 17, 1615-9	6.2	35
98	S100A4 amplifies TGF- β -induced fibroblast activation in systemic sclerosis. <i>Annals of the Rheumatic Diseases</i> , 2015 , 74, 1748-55	2.4	34
97	Monoclonal antibodies to human cartilage oligomeric matrix protein: epitope mapping and characterization of sandwich ELISA. <i>Clinica Chimica Acta</i> , 2003 , 328, 59-69	6.2	34
96	The metastasis promoting protein S100A4 is increased in idiopathic inflammatory myopathies. <i>Rheumatology</i> , 2011 , 50, 1766-72	3.9	33
95	Metastasis-inducing S100A4 protein is associated with the disease activity of rheumatoid arthritis. <i>Rheumatology</i> , 2009 , 48, 1590-4	3.9	32
94	Clinical improvement and reduction in serum calprotectin levels after an intensive exercise programme for patients with ankylosing spondylitis and non-radiographic axial spondyloarthritis. <i>Arthritis Research and Therapy</i> , 2016 , 18, 275	5.7	32
93	Efficacy and Safety of Anti-Interleukin-20 Monoclonal Antibody in Patients With Rheumatoid Arthritis: A Randomized Phase IIa Trial. <i>Arthritis and Rheumatology</i> , 2015 , 67, 1438-48	9.5	31
92	The transcription factor GLI2 as a downstream mediator of transforming growth factor- β -induced fibroblast activation in SSc. <i>Annals of the Rheumatic Diseases</i> , 2017 , 76, 756-764	2.4	28

91	Increased serum concentration of immune cell derived microparticles in polymyositis/dermatomyositis. <i>Immunology Letters</i> , 2010 , 128, 124-30	4.1	28
90	Increased levels of peptidylarginine deiminase 2 in synovial fluid from anti-CCP-positive rheumatoid arthritis patients: Association with disease activity and inflammatory markers. <i>Rheumatology</i> , 2016 , 55, 918-27	3.9	28
89	Association between circulating miRNAs and spinal involvement in patients with axial spondyloarthritis. <i>PLoS ONE</i> , 2017 , 12, e0185323	3.7	25
88	Adiponectin relation to skin changes and dyslipidemia in systemic sclerosis. <i>Cytokine</i> , 2012 , 58, 165-8	4	25
87	Calgizzarin (S100A11): a novel inflammatory mediator associated with disease activity of rheumatoid arthritis. <i>Arthritis Research and Therapy</i> , 2017 , 19, 79	5.7	24
86	Tribbles homologue 3 stimulates canonical TGF- β signalling to regulate fibroblast activation and tissue fibrosis. <i>Annals of the Rheumatic Diseases</i> , 2016 , 75, 609-16	2.4	24
85	The level of serum visfatin (PBEF) is associated with total number of B cells in patients with rheumatoid arthritis and decreases following B cell depletion therapy. <i>Cytokine</i> , 2011 , 55, 116-21	4	24
84	S100A4 (Mts1): is there any relation to the pathogenesis of rheumatoid arthritis?. <i>Autoimmunity Reviews</i> , 2006 , 5, 129-31	13.6	24
83	Interleukin 35 Synovial Fluid Levels Are Associated with Disease Activity of Rheumatoid Arthritis. <i>PLoS ONE</i> , 2015 , 10, e0132674	3.7	24
82	Progranulin Is Associated with Disease Activity in Patients with Rheumatoid Arthritis. <i>Mediators of Inflammation</i> , 2015 , 2015, 740357	4.3	23
81	Emerging therapies in rheumatoid arthritis: focus on monoclonal antibodies. <i>F1000Research</i> , 2019 , 8,	3.6	23
80	Relationship between serum calprotectin (S100A8/9) and clinical, laboratory and ultrasound parameters of disease activity in rheumatoid arthritis: A large cohort study. <i>PLoS ONE</i> , 2017 , 12, e0183427	3.7	19
79	Relative efficiencies of peptidylarginine deiminase 2 and 4 in generating target sites for anti-citrullinated protein antibodies in fibrinogen, alpha-enolase and histone H3. <i>PLoS ONE</i> , 2018 , 13, e0203214	3.7	19
78	Decreased circulating visfatin is associated with improved disease activity in early rheumatoid arthritis: data from the PERAC cohort. <i>PLoS ONE</i> , 2014 , 9, e103495	3.7	17
77	Resistin in idiopathic inflammatory myopathies. <i>Arthritis Research and Therapy</i> , 2012 , 14, R111	5.7	17
76	MicroRNA-331 and microRNA-151-3p as biomarkers in patients with ST-segment elevation myocardial infarction. <i>Scientific Reports</i> , 2020 , 10, 5845	4.9	15
75	No effect of physiotherapy on the serum levels of adipocytokines in patients with ankylosing spondylitis. <i>Clinical Rheumatology</i> , 2012 , 31, 67-71	3.9	15
74	Circulating S100 proteins effectively discriminate SLE patients from healthy controls: a cross-sectional study. <i>Rheumatology International</i> , 2019 , 39, 469-478	3.6	15

73	Interleukin-35 is upregulated in systemic sclerosis and its serum levels are associated with early disease. <i>Rheumatology</i> , 2015 , 54, 2273-82	3.9	14
72	The role of resistin in inflammatory myopathies. <i>Current Rheumatology Reports</i> , 2013 , 15, 336	4.9	14
71	The pre-clinical phase of rheumatoid arthritis: From risk factors to prevention of arthritis. <i>Autoimmunity Reviews</i> , 2021 , 20, 102797	13.6	14
70	The level of fatty acid-binding protein 4, a novel adipokine, is increased in rheumatoid arthritis and correlates with serum cholesterol levels. <i>Cytokine</i> , 2013 , 64, 441-7	4	13
69	The metastasis promoting protein S100A4 levels associate with disease activity rather than cancer development in patients with idiopathic inflammatory myopathies. <i>Arthritis Research and Therapy</i> , 2014 , 16, 468	5.7	13
68	Serum calprotectin may reflect inflammatory activity in patients with active rheumatoid arthritis despite normal to low C-reactive protein. <i>Clinical Rheumatology</i> , 2018 , 37, 2055-2062	3.9	12
67	Chronic repetitive nonprogressive epilepsy partialis continua due to rheumatoid meningitis. <i>Seizure: the Journal of the British Epilepsy Association</i> , 2013 , 22, 80-2	3.2	12
66	Serum tenascin-C discriminates patients with active SLE from inactive patients and healthy controls and predicts the need to escalate immunosuppressive therapy: a cohort study. <i>Arthritis Research and Therapy</i> , 2015 , 17, 341	5.7	12
65	Lower serum clusterin levels in patients with erosive hand osteoarthritis are associated with more pain. <i>BMC Musculoskeletal Disorders</i> , 2018 , 19, 264	2.8	12
64	Pro-inflammatory S100A11 is elevated in inflammatory myopathies and reflects disease activity and extramuscular manifestations in myositis. <i>Cytokine</i> , 2019 , 116, 13-20	4	11
63	Identification of Novel Native Autoantigens in Rheumatoid Arthritis. <i>Biomedicines</i> , 2020 , 8,	4.8	11
62	Expanding the citrullinome of synovial fibrinogen from rheumatoid arthritis patients. <i>Journal of Proteomics</i> , 2019 , 208, 103484	3.9	11
61	Biomarkers of hand osteoarthritis. <i>Rheumatology International</i> , 2018 , 38, 725-735	3.6	11
60	Interleukin-20 is triggered by TLR ligands and associates with disease activity in patients with rheumatoid arthritis. <i>Cytokine</i> , 2017 , 97, 187-192	4	10
59	High levels of metastasis-inducing S100A4 protein and treatment outcome in early rheumatoid arthritis: data from the PERAC cohort. <i>Biomarkers</i> , 2015 , 20, 47-51	2.6	9
58	Muscle and fat metabolism in obesity after kidney transplantation: no effect of peritoneal dialysis or hemodialysis. <i>Journal of Renal Nutrition</i> , 2012 , 22, 166-70	3	9
57	Adipokine profile is modulated in subcutaneous adipose tissue by TNF α inhibitors in patients with rheumatoid arthritis. <i>Annals of the Rheumatic Diseases</i> , 2011 , 70, 2054-6	2.4	8
56	Serum Calprotectin Discriminates Subclinical Disease Activity from Ultrasound-Defined Remission in Patients with Rheumatoid Arthritis in Clinical Remission. <i>PLoS ONE</i> , 2016 , 11, e0165498	3.7	8

55	Cross-sectional study of patients with axial spondyloarthritis fulfilling imaging arm of ASAS classification criteria: baseline clinical characteristics and subset differences in a single-centre cohort. <i>BMJ Open</i> , 2019 , 9, e024713	3	7
54	Real world use of secukinumab for treatment of axial spondyloarthritis and psoriatic arthritis: nationwide results from the ATTRA registry. <i>Clinical and Experimental Rheumatology</i> , 2019 , 37, 342-343	2.2	7
53	Pentosidine, an advanced glycation end-product, may reflect clinical and morphological features of hand osteoarthritis. <i>Open Rheumatology Journal</i> , 2012 , 6, 64-9	0.2	5
52	Early changes in serum visfatin after abdominal surgery: a new pro-inflammatory marker in diagnosis?. <i>Biomedical Papers of the Medical Faculty of the University Palacky&#x0301;, Olomouc, Czechoslovakia</i> , 2015 , 159, 489-96	1.7	5
51	Serum visfatin levels in patients with axial spondyloarthritis and their relationship to disease activity and spinal radiographic damage: a cross-sectional study. <i>Rheumatology International</i> , 2019 , 39, 1037-1043	3.6	4
50	Cytokines and inflammatory mediators as promising markers of polymyositis/dermatomyositis. <i>Current Opinion in Rheumatology</i> , 2020 , 32, 534-541	5.3	4
49	S100A11 (calgizzarin) is released via NETosis in rheumatoid arthritis (RA) and stimulates IL-6 and TNF secretion by neutrophils. <i>Scientific Reports</i> , 2021 , 11, 6063	4.9	3
48	Clusterin serum levels are elevated in patients with early rheumatoid arthritis and predict disease activity and treatment response. <i>Scientific Reports</i> , 2021 , 11, 11525	4.9	3
47	Applicability of Small-Molecule Inhibitors in the Study of Peptidyl Arginine Deiminase 2 (PAD2) and PAD4. <i>Frontiers in Immunology</i> , 2021 , 12, 716250	8.4	2
46	Chemokine and Cytokine Profiles in Patients with Hand Osteoarthritis. <i>Biomolecules</i> , 2020 , 11,	5.9	2
45	P187 Secukinumab significantly decreased joint synovitis measured by Power Doppler ultrasonography in biologic-naive patients with active psoriatic arthritis: primary (12week) results from a randomised, placebo-controlled Phase 3 study. <i>Rheumatology</i> , 2021 , 60,	3.9	2
44	The effect of a 24-week training focused on activities of daily living, muscle strengthening, and stability in idiopathic inflammatory myopathies: a monocentric controlled study with follow-up. <i>Arthritis Research and Therapy</i> , 2021 , 23, 173	5.7	2
43	Plasma heat shock protein 90 levels in patients with spondyloarthritis and their relation to structural changes: a cross-sectional study. <i>Biomarkers in Medicine</i> , 2021 , 15, 5-13	2.3	2
42	A6.26 Analysis of circulating mirnas in patients with axial sponyloarthritis with different spinal involvement. <i>Annals of the Rheumatic Diseases</i> , 2015 , 74, A66.1-A66	2.4	1
41	A3.10 Plasma levels of heat shock protein 90 correlate with disease activity, lung involvement and skin fibrosis in systemic sclerosis. <i>Annals of the Rheumatic Diseases</i> , 2015 , 74, A35.1-A35	2.4	1
40	THU0528 Decrease in Circulating Visfatin Levels is Associated with Disease Activity Improvement in Early Rheumatoid Arthritis. <i>Annals of the Rheumatic Diseases</i> , 2014 , 73, 366.1-366	2.4	1
39	A1.48 Enhanced expression of CD11c on non-classical CD16+ peripheral blood monocytes in early rheumatoid arthritis. <i>Annals of the Rheumatic Diseases</i> , 2014 , 73, A20.2-A21	2.4	1
38	A3.16 Serum S100A4 correlates with skin fibrosis, lung involvement and disease activity in systemic sclerosis. <i>Annals of the Rheumatic Diseases</i> , 2014 , 73, A48.1-A48	2.4	1

37	AB0203 S100A4 Serum Levels Correlate with Disease Activity, Skin Fibrosis and Lung Involvement in Systemic Sclerosis. <i>Annals of the Rheumatic Diseases</i> , 2014 , 73, 871.1-871	2.4	1
36	MiRs in RA: possible biomarkers and therapeutic targets. <i>Arthritis Research and Therapy</i> , 2012 , 14,	5.7	1
35	Calprotectin (a S100 protein) as a sensitive biomarker for rheumatoid arthritis: new perspectives for an old finding. <i>International Journal of Clinical Rheumatology</i> , 2012 , 7, 127-129	1.5	1
34	Inhibition of Hsp90 Counteracts the Established Experimental Dermal Fibrosis Induced by Bleomycin. <i>Biomedicines</i> , 2021 , 9,	4.8	1
33	Interleukin-35 in idiopathic inflammatory myopathies. <i>Cytokine</i> , 2021 , 137, 155350	4	1
32	High miR-451 expression in peripheral blood mononuclear cells from subjects at risk of developing rheumatoid arthritis. <i>Scientific Reports</i> , 2021 , 11, 4719	4.9	1
31	Identification of potential autoantigens in anti-CCP-positive and anti-CCP-negative rheumatoid arthritis using citrulline-specific protein arrays. <i>Scientific Reports</i> , 2021 , 11, 17300	4.9	1
30	Response to Secukinumab on Synovitis using Power Doppler Ultrasound in Psoriatic Arthritis: 12-week Results from a Phase III Study, ULTIMATE. <i>Rheumatology</i> , 2021 ,	3.9	1
29	Increased visfatin levels are associated with higher disease activity in anti-Jo-1-positive myositis patients. <i>Clinical and Experimental Rheumatology</i> , 2016 , 34, 222-9	2.2	1
28	S100A11 (calgizzarin) is released by circulating mononuclear cells and its elevated plasma levels distinguish systemic lupus erythematosus patients from healthy individuals. <i>Clinical and Experimental Rheumatology</i> , 2019 , 37, 338-339	2.2	1
27	Clusterin is upregulated in serum and muscle tissue in idiopathic inflammatory myopathies and associates with clinical disease activity and cytokine profile. <i>Clinical and Experimental Rheumatology</i> , 2021 , 39, 1021-1032	2.2	1
26	S100A4 is elevated in axial spondyloarthritis: a potential link to disease severity. <i>BMC Rheumatology</i> , 2020 , 4, 13	2.9	0
25	Hsp90 Levels in Idiopathic Inflammatory Myopathies and Their Association With Muscle Involvement and Disease Activity: A Cross-Sectional and Longitudinal Study.. <i>Frontiers in Immunology</i> , 2022 , 13, 811045	8.4	0
24	IL-40: A New B Cell-Associated Cytokine Up-Regulated in Rheumatoid Arthritis Decreases Following the Rituximab Therapy and Correlates With Disease Activity, Autoantibodies, and NETosis. <i>Frontiers in Immunology</i> , 2021 , 12, 745523	8.4	0
23	Sexual function in patients with idiopathic inflammatory myopathies: a cross-sectional study. <i>Rheumatology</i> , 2021 , 60, 5060-5072	3.9	0
22	A3.10 Serum calprotectinis elevated in patients with early rheumatoid arthritis but not in patients at risk of developing rheumatoid arthritis. <i>Annals of the Rheumatic Diseases</i> , 2016 , 75, A36.1-A36	2.4	
21	A10.17 Interleukin-20 serum levels correlate with disease activity in rheumatoid arthritis patients and decrease in response to rituximab treatment. <i>Annals of the Rheumatic Diseases</i> , 2016 , 75, A79.2-A79 ^{2.4}		
20	THU0353 Serum Tweak Levels in Patients with Neuropsychiatric Systemic Lupus Erythematosus. <i>Annals of the Rheumatic Diseases</i> , 2016 , 75, 315.1-315	2.4	

- 19 SAT0068 Circulating MIR-126 Is A Marker of Disease Activity in Patients with Early RA but Does Not Differentiate Patients at Risk of Developing RA. *Annals of the Rheumatic Diseases*, **2016**, 75, 689.2-689 2.4
- 18 AB0696 Clinical Characteristics of Spondyloarthritis in The Czech Republic: A Single Centre Cohort. *Annals of the Rheumatic Diseases*, **2016**, 75, 1142.2-1143 2.4
- 17 A8.06 MicroRNA-125b expression in PBMCS is inversely associated with disease activity in patients with early rheumatoid arthritis. *Annals of the Rheumatic Diseases*, **2016**, 75, A67.1-A67 2.4
- 16 AB0599 Interleukin-35 in Idiopathic Inflammatory Myopathies. *Annals of the Rheumatic Diseases*, **2016**, 75, 1109.2-1109 2.4
- 15 AB0163 Comprehensive Analysis of Circulating Mirnas in Patients with Axial Spondyloarthritis. *Annals of the Rheumatic Diseases*, **2014**, 73, 856.4-857 2.4
- 14 SAT0565 The Expression and Regulation of the Argonaute Protein Family Member PIWIL4 in RA. *Annals of the Rheumatic Diseases*, **2014**, 73, 795.1-795 2.4
- 13 SAT0188 Serum Calprotectin (S100A8/9) Correlates with Clinical and Ultrasound Outcomes in Patients with Early Rheumatoid Arthritis. *Annals of the Rheumatic Diseases*, **2014**, 73, 658.1-658 2.4
- 12 THU0168 Monotherapy with Biological Disease-Modifying Anti-Rheumatic Drugs-Data from the Czech Registry Attra. *Annals of the Rheumatic Diseases*, **2014**, 73, 238.2-238 2.4
- 11 OP0240 Signal Transducer and Activator of Transcription 3 Regulates Fibroblast Activation as A Downstream Mediator of Transforming Growth Factor-Beta. *Annals of the Rheumatic Diseases*, **2014**, 73, 153.1-153 2.4
- 10 AB0114 Comparative Analysis of the Expression of Protein Tyrosine Phosphatase Non-Receptor Type 2 (PTPN2) in Autoimmune Disease. *Annals of the Rheumatic Diseases*, **2014**, 73, 841.3-842 2.4
- 9 SAT0205 Potential of US 7 Score in Evaluating of Disease Activity of Patients with Rheumatoid Arthritis in the State of Remission. *Annals of the Rheumatic Diseases*, **2014**, 73, 664.1-664 2.4
- 8 A8.3 Deficit of S100A4 Prevents Joint Destruction and Systemic Bone Loss in hTNFtg Mouse Model. *Annals of the Rheumatic Diseases*, **2013**, 72, A58.1-A58 2.4
- 7 A7.3 Association of Circulating miR-223 and miR-16 with Disease Activity in Patients with Early Rheumatoid Arthritis. *Annals of the Rheumatic Diseases*, **2013**, 72, A48.3-A49 2.4
- 6 Metastasis-Promoting protein S100A4 (MTS1) upregulates production of proinflammatory cytokines via TLR-4. *Annals of the Rheumatic Diseases*, **2010**, 69, A38-A39 2.4
- 5 Modulation of subcutaneous adipose tissue adipokines by TNF- α blockade therapy in patients with inflammatory arthritides. *Annals of the Rheumatic Diseases*, **2011**, 70, A85-A85 2.4
- 4 Potential role of S100A4 in an immune response of rheumatoid arthritis. *Annals of the Rheumatic Diseases*, **2011**, 70, A85-A85 2.4
- 3 SAT0466 Progression of Pain and Ultrasound Detected Synovitis in Patients with Erosive Hand Osteoarthritis over One Year. *Annals of the Rheumatic Diseases*, **2016**, 75, 839.3-840 2.4
- 2 AB0952 Serum Visfatin and Resistin, but Not Adiponectin or Leptin, Are Associated with Ultrasound Synovitis in Patients with Rheumatoid Arthritis. *Annals of the Rheumatic Diseases*, **2016**, 75, 1227.1-1227 2.4

- 1 A9.02 Heat shock protein 90 plasma levels correlate with disease activity, skeletal muscle, lung and heart involvement in idiopathic inflammatory myopathies. *Annals of the Rheumatic Diseases*, **2016**, 75, A70.2-A71 2.4