Katsushi Tokunaga

List of Publications by Citations

Source: https://exaly.com/author-pdf/239523/katsushi-tokunaga-publications-by-citations.pdf

Version: 2024-04-19

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

173
papers

4,031
citations

h-index

55
g-index

183
ext. papers

4,742
ext. citations

5
avg, IF

L-index

#	Paper	IF	Citations
173	An ancient retrotransposal insertion causes Fukuyama-type congenital muscular dystrophy. <i>Nature</i> , 1998 , 394, 388-92	50.4	665
172	Genome-wide association study identifies TNFSF15 and POU2AF1 as susceptibility loci for primary biliary cirrhosis in the Japanese population. <i>American Journal of Human Genetics</i> , 2012 , 91, 721-8	11	201
171	FTO polymorphisms in oceanic populations. <i>Journal of Human Genetics</i> , 2007 , 52, 1031-1035	4.3	118
170	Genetic origins of the Ainu inferred from combined DNA analyses of maternal and paternal lineages. <i>Journal of Human Genetics</i> , 2004 , 49, 187-193	4.3	99
169	Strong association between hla-a*02:06 and acetaminophen-related stevens-johnson syndrome with severe mucosal involvements in the Japanese. <i>Clinical and Translational Allergy</i> , 2014 , 4, P11	5.2	78
168	HLA association with antipyretic analgesics-induced Stevens-Johnson Syndrome/toxic epidermal necrolysis with severe ocular surface complications in japanese patients. <i>Clinical and Translational Allergy</i> , 2014 , 4, P89	5.2	78
167	Genome-Wide Association Study Identifies TLL1 Variant Associated With Development of Hepatocellular Carcinoma After Eradication of Hepatitis C Virus Infection. <i>Gastroenterology</i> , 2017 , 152, 1383-1394	13.3	76
166	Association of TCF7L2 polymorphisms with susceptibility to type 2 diabetes in 4,087 Japanese subjects. <i>Journal of Human Genetics</i> , 2008 , 53, 174-180	4.3	76
165	Independent strong association of HLA-A*02:06 and HLA-B*44:03 with cold medicine-related Stevens-Johnson syndrome with severe mucosal involvement. <i>Scientific Reports</i> , 2014 , 4, 4862	4.9	67
164	Recombination and gene conversion-like events may contribute to ABO gene diversity causing various phenotypes. <i>Immunogenetics</i> , 2001 , 53, 190-9	3.2	64
163	A genome-wide association study for diabetic retinopathy in a Japanese population: potential association with a long intergenic non-coding RNA. <i>PLoS ONE</i> , 2014 , 9, e111715	3.7	62
162	Genome-wide Association Study of Autism Spectrum Disorder in the East Asian Populations. <i>Autism Research</i> , 2016 , 9, 340-9	5.1	59
161	Trans-ethnic study confirmed independent associations of HLA-A*02:06 and HLA-B*44:03 with cold medicine-related Stevens-Johnson syndrome with severe ocular surface complications. <i>Scientific Reports</i> , 2014 , 4, 5981	4.9	55
160	IKZF1, a new susceptibility gene for cold medicine-related Stevens-Johnson syndrome/toxic epidermal necrolysis with severe mucosal involvement. <i>Journal of Allergy and Clinical Immunology</i> , 2015 , 135, 1538-45.e17	11.5	50
159	Identification of the gene variations in human CD22. <i>Immunogenetics</i> , 1999 , 49, 280-6	3.2	50
158	Different Alleles Cause an Imbalance in A2 and A2B Phenotypes of the ABO Blood Group. <i>Vox Sanguinis</i> , 1998 , 74, 242-247	3.1	49
157	Genetic variations on the Y chromosome in the Japanese population and implications for modern human Y chromosome lineage. <i>Journal of Human Genetics</i> , 1999 , 44, 240-5	4.3	47

156	Contribution of HLA genes to genetic predisposition in diffuse panbronchiolitis. <i>American Journal of Respiratory and Critical Care Medicine</i> , 1998 , 158, 846-50	10.2	45	
155	Associations of human leukocyte antigens with autoimmune diseases: challenges in identifying the mechanism. <i>Journal of Human Genetics</i> , 2015 , 60, 697-702	4.3	42	
154	Epigenome-wide association study of DNA methylation in panic disorder. <i>Clinical Epigenetics</i> , 2017 , 9, 6	7.7	39	
153	HLA class II alleles in Ainu living in Hidaka District, Hokkaido, northern Japan. <i>American Journal of Physical Anthropology</i> , 1996 , 101, 1-9	2.5	38	
152	DBTSS/DBKERO for integrated analysis of transcriptional regulation. <i>Nucleic Acids Research</i> , 2018 , 46, D229-D238	20.1	37	
151	Strong Association of the Locus with Childhood Steroid-Sensitive Nephrotic Syndrome in the Japanese Population. <i>Journal of the American Society of Nephrology: JASN</i> , 2018 , 29, 2189-2199	12.7	37	
150	Association of Cw11 in Japanese patients with psoriasis vulgaris. <i>Tissue Antigens</i> , 1990 , 36, 241-2		37	
149	Genome-wide association studies identify PRKCB as a novel genetic susceptibility locus for primary biliary cholangitis in the Japanese population. <i>Human Molecular Genetics</i> , 2017 , 26, 650-659	5.6	36	
148	A 12-kb structural variation in progressive myoclonic epilepsy was newly identified by long-read whole-genome sequencing. <i>Journal of Human Genetics</i> , 2019 , 64, 359-368	4.3	36	
147	Founder-haplotype analysis in Fukuyama-type congenital muscular dystrophy (FCMD). <i>Human Genetics</i> , 1998 , 103, 323-7	6.3	35	
146	Appropriate data cleaning methods for genome-wide association study. <i>Journal of Human Genetics</i> , 2008 , 53, 886-893	4.3	34	
145	Possible association of human leucocyte antigen DR1 with delayed sleep phase syndrome. <i>Psychiatry and Clinical Neurosciences</i> , 1999 , 53, 527-9	6.2	34	
144	HLA and Human Mate Choice. Tests on Japanese Couples Anthropological Science, 2000, 108, 199-214	1.3	32	
143	Pathogen lineage-based genome-wide association study identified CD53 as susceptible locus in tuberculosis. <i>Journal of Human Genetics</i> , 2017 , 62, 1015-1022	4.3	31	
142	A polymorphism in CCR1/CCR3 is associated with narcolepsy. <i>Brain, Behavior, and Immunity</i> , 2015 , 49, 148-55	16.6	30	
141	NAT2 variants are associated with drug-induced liver injury caused by anti-tuberculosis drugs in Indonesian patients with tuberculosis. <i>Journal of Human Genetics</i> , 2016 , 61, 533-7	4.3	30	
140	Evidence for Host-Bacterial Co-evolution via Genome Sequence Analysis of 480 Thai Mycobacterium tuberculosis Lineage 1 Isolates. <i>Scientific Reports</i> , 2018 , 8, 11597	4.9	30	
139	Discerning the Origins of the Negritos, First Sundaland People: Deep Divergence and Archaic Admixture. <i>Genome Biology and Evolution</i> , 2017 , 9, 2013-2022	3.9	30	

138	Analysis of human leukocyte antigen-G polymorphism including intron 4 in Japanese couples with habitual abortion. <i>American Journal of Reproductive Immunology</i> , 1999 , 41, 159-63	3.8	30
137	Understanding of HLA-conferred susceptibility to chronic hepatitis B infection requires HLA genotyping-based association analysis. <i>Scientific Reports</i> , 2016 , 6, 24767	4.9	29
136	New susceptibility and resistance HLA-DP alleles to HBV-related diseases identified by a trans-ethnic association study in Asia. <i>PLoS ONE</i> , 2014 , 9, e86449	3.7	28
135	Polymorphisms of promoter and coding regions of the arylamine N-acetyltransferase 2 (NAT2) gene in the Indonesian population: proposal for a new nomenclature. <i>Journal of Human Genetics</i> , 2008 , 53, 201-209	4.3	28
134	Power of genome-wide linkage disequilibrium testing by using microsatellite markers. <i>Journal of Human Genetics</i> , 2003 , 48, 487-491	4.3	28
133	Sequence Variation of Human Platelet Membrane Glycoprotein IIIa Associated with the Yuka/Yukb Alloantigen System <i>Proceedings of the Japan Academy Series B: Physical and Biological Sciences</i> , 1991 , 67, 102-106	4	28
132	Association and interaction analyses of NRG1 and ERBB4 genes with schizophrenia in a Japanese population. <i>Journal of Human Genetics</i> , 2008 , 53, 929-935	4.3	27
131	High-density Association Mapping and Interaction Analysis of PLA2R1 and HLA Regions with Idiopathic Membranous Nephropathy in Japanese. <i>Scientific Reports</i> , 2016 , 6, 38189	4.9	27
130	Genetics of narcolepsy. Human Genome Variation, 2019, 6, 4	1.8	26
129	Key HLA-DRB1-DQB1 haplotypes and role of the BTNL2 gene for response to a hepatitis B vaccine. <i>Hepatology</i> , 2018 , 68, 848-858	11.2	26
128	Profiling of genes expressed in human monocytes and monocyte-derived dendritic cells using cDNA expression array. <i>British Journal of Haematology</i> , 2001 , 114, 191-7	4.5	25
127	Determination of Granulocyte-Specific Antigens on Neutrophil FclReceptor IIIb by PCR-Preferential Homoduplex Formation Assay, and Gene Frequencies in the Japanese Population. <i>Vox Sanguinis</i> , 1999 , 77, 218-222	3.1	25
126	Human leukocyte antigen variants and risk of hepatocellular carcinoma modified by hepatitis C virus genotypes: A genome-wide association study. <i>Hepatology</i> , 2018 , 67, 651-661	11.2	24
125	Association of diffuse panbronchiolitis with microsatellite polymorphism of the human interleukin 8 (IL-8) gene. <i>Journal of Human Genetics</i> , 1999 , 44, 169-72	4.3	24
124	Human Leukocyte Antigen Class I Genes Associated With Stevens-Johnson Syndrome and Severe Ocular Complications Following Use of Cold Medicine in a Brazilian Population. <i>JAMA Ophthalmology</i> , 2017 , 135, 355-360	3.9	23
123	Genome-wide association study identified new susceptible genetic variants in HLA class I region for hepatitis B virus-related hepatocellular carcinoma. <i>Scientific Reports</i> , 2018 , 8, 7958	4.9	23
122	Support for an association between HLA-DR1 and schizophrenia in the Japanese population. <i>American Journal of Medical Genetics Part A</i> , 2000 , 96, 725-7		22
121	The First Pilot Genome-Wide Gene-Environment Study of Depression in the Japanese Population. <i>PLoS ONE</i> , 2016 , 11, e0160823	3.7	22

120	Human primary biliary cirrhosis-susceptible allele of rs4979462 enhances TNFSF15 expression by binding NF-1. <i>Human Genetics</i> , 2015 , 134, 737-47	6.3	21
119	A functional SNP in the NKX2.5-binding site of ITPR3 promoter is associated with susceptibility to systemic lupus erythematosus in Japanese population. <i>Journal of Human Genetics</i> , 2008 , 53, 151-162	4.3	21
118	Phenome-wide association study maps new diseases to the human major histocompatibility complex region. <i>Journal of Medical Genetics</i> , 2016 , 53, 681-9	5.8	21
117	New susceptibility variants to narcolepsy identified in HLA class II region. <i>Human Molecular Genetics</i> , 2015 , 24, 891-8	5.6	20
116	GA4GH: International policies and standards for data sharing across genomic research and healthcare <i>Cell Genomics</i> , 2021 , 1, 100029-100029		20
115	NAT2 ultra-slow acetylator and risk of anti-tuberculosis drug-induced liver injury: a genotype-based meta-analysis. <i>Pharmacogenetics and Genomics</i> , 2018 , 28, 167-176	1.9	20
114	Unique characteristics of the Ainu population in Northern Japan. <i>Journal of Human Genetics</i> , 2015 , 60, 565-71	4.3	19
113	Genetic diversity and dynamic distribution of Mycobacterium tuberculosis isolates causing pulmonary and extrapulmonary tuberculosis in Thailand. <i>Journal of Clinical Microbiology</i> , 2014 , 52, 4267	-9:7	19
112	Development of a prediction system for anti-tuberculosis drug-induced liver injury in Japanese patients. <i>Human Genome Variation</i> , 2016 , 3, 16014	1.8	19
111	Genome-wide association study of self-reported food reactions in Japanese identifies shrimp and peach specific loci in the HLA-DR/DQ gene region. <i>Scientific Reports</i> , 2018 , 8, 1069	4.9	18
110	Principal contribution of HLA-DQ alleles, DQB1*06:04 and DQB1*03:01, to disease resistance against primary biliary cholangitis in a Japanese population. <i>Scientific Reports</i> , 2017 , 7, 11093	4.9	18
109	HLA-A*02:06 and PTGER3 polymorphism exert additive effects in cold medicine-related Stevens-Johnson syndrome with severe ocular complications. <i>Human Genome Variation</i> , 2015 , 2, 15023	1.8	18
108	Reply to: "Lack of HIF-2[In limb bud mesenchyme causes a modest and transient delay of endochondral bone development" and "Replication studies in various ethnic populations do not support the association of the HIF-2[Is]NP rs17039192 with knee osteoarthritis". <i>Nature Medicine</i> ,	50.5	18
107	2011, 17, 27-29 Genome-Wide Association Study Identifies Risk Variants for Lichen Planus in Patients With Hepatitis C Virus Infection. <i>Clinical Gastroenterology and Hepatology</i> , 2017, 15, 937-944.e5	6.9	17
106	Common risk variants in NPHS1 and TNFSF15 are associated with childhood steroid-sensitive nephrotic syndrome. <i>Kidney International</i> , 2020 , 98, 1308-1322	9.9	17
105	Evaluation of polygenic risks for narcolepsy and essential hypersomnia. <i>Journal of Human Genetics</i> , 2016 , 61, 873-878	4.3	17
104	Mutations in rrs, rpsL and gidB in streptomycin-resistant Mycobacterium tuberculosis isolates from Thailand. <i>Journal of Global Antimicrobial Resistance</i> , 2016 , 4, 5-10	3.4	17
103	HLA-A26 subtype A pockets accommodate acidic N-termini of ligands. <i>Immunogenetics</i> , 1998 , 48, 350-3	3.2	17

102	Genome-Wide Association Study Confirming a Strong Effect of HLA and Identifying Variants in on Chromosome 12q13.13 Associated With Susceptibility to Fulminant Type 1 Diabetes. <i>Diabetes</i> , 2019 , 68, 665-675	0.9	17
101	Genome-wide association study using the ethnicity-specific Japonica array: identification of new susceptibility loci for cold medicine-related Stevens-Johnson syndrome with severe ocular complications. <i>Journal of Human Genetics</i> , 2017 , 62, 485-489	4.3	16
100	Disease susceptibility genes shared by primary biliary cirrhosis and Crohn@ disease in the Japanese population. <i>Journal of Human Genetics</i> , 2015 , 60, 525-31	4.3	16
99	An association analysis of HLA-DQB1 with narcolepsy without cataplexy and idiopathic hypersomnia with/without long sleep time in a Japanese population. <i>Human Genome Variation</i> , 2015 , 2, 15031	1.8	16
98	Analysis of whole Y-chromosome sequences reveals the Japanese population history in the Jomon period. <i>Scientific Reports</i> , 2019 , 9, 8556	4.9	15
97	Identification of the functional variant driving ORMDL3 and GSDMB expression in human chromosome 17q12-21 in primary biliary cholangitis. <i>Scientific Reports</i> , 2017 , 7, 2904	4.9	15
96	Role of B cell inhibitory receptor polymorphisms in systemic lupus erythematosus: a negative times a negative makes a positive. <i>Journal of Human Genetics</i> , 2006 , 51, 741-750	4.3	15
95	Polymorphisms in the ABO blood group gene in three populations in the New Georgia group of the Solomon Islands. <i>Journal of Human Genetics</i> , 2006 , 51, 407-411	4.3	15
94	A common Japanese haplotype HLA-A26-Cw3-B61-DR9-DQ3 carries HLA-B*4002. <i>Tissue Antigens</i> , 1992 , 40, 257-60		15
93	POGLUT1, the putative effector gene driven by rs2293370 in primary biliary cholangitis susceptibility locus chromosome 3q13.33. <i>Scientific Reports</i> , 2019 , 9, 102	4.9	14
92	De novo polyalanine expansion of PHOX2B in congenital central hypoventilation syndrome: unequal sister chromatid exchange during paternal gametogenesis. <i>Journal of Human Genetics</i> , 2007 , 52, 921-925	4.3	14
91	Comparative FISH mapping of the ancestral fusion point of human chromosome 2. <i>Chromosome Research</i> , 2000 , 8, 727-35	4.4	14
90	Digenic mutations in and impair formaldehyde clearance and cause a multisystem disorder, AMeD syndrome. <i>Science Advances</i> , 2020 , 6,	14.3	14
89	The first genome-wide association study identifying new susceptibility loci for obstetric antiphospholipid syndrome. <i>Journal of Human Genetics</i> , 2017 , 62, 831-838	4.3	13
88	Genomewide Association Study Confirming the Association of with Susceptibility to Antituberculosis Drug-Induced Liver Injury in Thai Patients. <i>Antimicrobial Agents and Chemotherapy</i> , 2019 , 63,	5.9	13
87	Glypican-5 Increases Susceptibility to Nephrotic Damage in Diabetic Kidney. <i>American Journal of Pathology</i> , 2015 , 185, 1889-98	5.8	13
86	Association of HLA class I and II gene polymorphisms with acetaminophen-related Stevens-Johnson syndrome with severe ocular complications in Japanese individuals. <i>Human Genome Variation</i> , 2019 , 6, 50	1.8	13
85	Association of Human Leukocyte Antigen Class 1 genes with Stevens Johnson Syndrome with severe ocular complications in an Indian population. <i>Scientific Reports</i> , 2017 , 7, 15960	4.9	13

(2021-2015)

84	Detection of Hereditary 1,25-Hydroxyvitamin D-Resistant Rickets Caused by Uniparental Disomy of Chromosome 12 Using Genome-Wide Single Nucleotide Polymorphism Array. <i>PLoS ONE</i> , 2015 , 10, e013	3131757	13	
83	MHC (major histocompatibility complex)-DRB genes and polymorphisms in common marmoset. <i>Journal of Molecular Evolution</i> , 2000 , 51, 214-22	3.1	13	
82	Epigenome-wide association study of DNA methylation in narcolepsy: an integrated genetic and epigenetic approach. <i>Sleep</i> , 2018 , 41,	1.1	12	
81	Amino acid position 37 of HLA-DRI affects susceptibility to Crohn@ disease in Asians. <i>Human Molecular Genetics</i> , 2018 , 27, 3901-3910	5.6	12	
8o	Haplotype analyses with the human leucocyte antigen and tumour necrosis factor-alpha genes in narcolepsy families. <i>Psychiatry and Clinical Neurosciences</i> , 2001 , 55, 37-9	6.2	12	
79	Role of HLA-DP and HLA-DQ on the clearance of hepatitis B virus and the risk of chronic infection in a multiethnic population. <i>Liver International</i> , 2017 , 37, 1476-1487	7.9	11	
78	A missense variant in PER2 is associated with delayed sleep-wake phase disorder in a Japanese population. <i>Journal of Human Genetics</i> , 2019 , 64, 1219-1225	4.3	11	
77	Whole-exome sequencing and gene-based rare variant association tests suggest that PLA2G4E might be a risk gene for panic disorder. <i>Translational Psychiatry</i> , 2018 , 8, 41	8.6	11	
76	Immune-related pathways including HLA-DRB1(*)13:02 are associated with panic disorder. <i>Brain, Behavior, and Immunity</i> , 2015 , 46, 96-103	16.6	11	
75	Negative association of the HLA-DRB1*1502-DQB1*0601 haplotype with human narcolepsy. <i>Immunogenetics</i> , 2001 , 52, 299-301	3.2	11	
74	New variations of human SHP-1. <i>Immunogenetics</i> , 1999 , 49, 577-9	3.2	11	
73	A Genome-wide Association Study Identifying RAP1A as a Novel Susceptibility Gene for Crohn® Disease in Japanese Individuals. <i>Journal of Crohn</i> and Colitis, 2019 , 13, 648-658	1.5	11	
72	NFKB1 and MANBA Confer Disease Susceptibility to Primary Biliary Cholangitis via Independent Putative Primary Functional Variants. <i>Cellular and Molecular Gastroenterology and Hepatology</i> , 2019 , 7, 515-532	7.9	10	
71	Identification of a telomeric boundary of the HLA region with potential for predisposition to human narcolepsy. <i>Immunogenetics</i> , 2000 , 52, 12-8	3.2	10	
70	Fine-mapping analysis revealed complex pleiotropic effect and tissue-specific regulatory mechanism of TNFSF15 in primary biliary cholangitis, Crohn@ disease and leprosy. <i>Scientific Reports</i> , 2016 , 6, 31429	4.9	9	
69	Novel rare variations of the oxytocin receptor (OXTR) gene in autism spectrum disorder individuals. <i>Human Genome Variation</i> , 2015 , 2, 15024	1.8	9	
68	Establishment of a method of anonymization of DNA samples in genetic research. <i>Journal of Human Genetics</i> , 2003 , 48, 327-330	4.3	9	
67	X Chromosome Contribution to the Genetic Architecture of Primary Biliary Cholangitis. Gastroenterology, 2021 , 160, 2483-2495.e26	13.3	9	

66	Homoplastic single nucleotide polymorphisms contributed to phenotypic diversity in Mycobacterium tuberculosis. <i>Scientific Reports</i> , 2020 , 10, 8024	4.9	8
65	A high-resolution HLA imputation system for the Taiwanese population: a study of the Taiwan Biobank. <i>Pharmacogenomics Journal</i> , 2020 , 20, 695-704	3.5	8
64	Identification of ITPA on chromosome 20 as a susceptibility gene for young-onset tuberculosis. <i>Human Genome Variation</i> , 2016 , 3, 15067	1.8	8
63	Novel DNA chip based on a modified DigiTag2 assay for high-throughput species identification and genotyping of Mycobacterium tuberculosis complex isolates. <i>Journal of Clinical Microbiology</i> , 2014 , 52, 1962-8	9.7	8
62	Determination of granulocyte-specific antigens on neutrophil FcA receptor IIIb by PCR-preferential homoduplex formation assay, and gene frequencies in the Japanese population. <i>Vox Sanguinis</i> , 1999 , 77, 218-22	3.1	8
61	The Origin and Composition of Korean Ethnicity Analyzed by Ancient and Present-Day Genome Sequences. <i>Genome Biology and Evolution</i> , 2020 , 12, 553-565	3.9	7
60	An epigenome-wide methylation study of healthy individuals with or without depressive symptoms. Journal of Human Genetics, 2018 , 63, 319-326	4.3	7
59	Identification of HLA-A*02:06:01 as the primary disease susceptibility HLA allele in cold medicine-related Stevens-Johnson syndrome with severe ocular complications by high-resolution NGS-based HLA typing. <i>Scientific Reports</i> , 2019 , 9, 16240	4.9	7
58	Genome-Wide Association Study Identifies ZNF354C Variants Associated with Depression from Interferon-Based Therapy for Chronic Hepatitis C. <i>PLoS ONE</i> , 2016 , 11, e0164418	3.7	7
57	Polymorphisms in the TMEM132D region are associated with panic disorder in HLA-DRB1*13:02-negative individuals of a Japanese population. <i>Human Genome Variation</i> , 2016 , 3, 160	0 1 .8	7
56	Identification of a two-SNP PLA2R1 Haplotype and HLA-DRB1 Alleles as Primary Risk Associations in Idiopathic Membranous Nephropathy. <i>Scientific Reports</i> , 2018 , 8, 15576	4.9	7
55	NELFCD and CTSZ loci are associated with jaundice-stage progression in primary biliary cholangitis in the Japanese population. <i>Scientific Reports</i> , 2018 , 8, 8071	4.9	7
54	Integrative genome analysis identified the KANNO blood group antigen as prion protein. <i>Transfusion</i> , 2019 , 59, 2429-2435	2.9	6
53	Association of functional polymorphisms in interferon regulatory factor 2 (IRF2) with susceptibility to systemic lupus erythematosus: a case-control association study. <i>PLoS ONE</i> , 2014 , 9, e109764	3.7	6
52	Comparative study of the haplotype structure and linkage disequilibrium of chromosome 1p36.2 region in the Korean and Japanese populations. <i>Journal of Human Genetics</i> , 2004 , 49, 603-609	4.3	6
51	Haplotype-specific sequence encoding the protein kinase, interferon-inducible double-stranded RNA-dependent activator in the human leukocyte antigen class II region. <i>Immunogenetics</i> , 2001 , 52, 186	3-3-2	6
50	A human PSMB11 variant affects thymoproteasome processing and CD8+ T cell production. <i>JCI Insight</i> , 2017 , 2,	9.9	6
49	Protective association of HLA-DRB1*13:02, HLA-DRB1*04:06, and HLA-DQB1*06:04 alleles with cervical cancer in a Korean population. <i>Human Immunology</i> , 2019 , 80, 107-111	2.3	6

48	Haplotype-specific PCR for diplotyping. Human Genome Variation, 2020, 7, 13	1.8	5
47	Genome-wide analysis of CNV (copy number variation) and their associations with narcolepsy in a Japanese population. <i>Journal of Human Genetics</i> , 2014 , 59, 235-40	4.3	5
46	Linkage disequilibrium structure of the 5q31-33 region in a Thai population. <i>Journal of Human Genetics</i> , 2008 , 53, 850-856	4.3	5
45	HLA-DRB1 polymorphism on Ha@no island of the Kingdom of Tonga. <i>Anthropological Science</i> , 2006 , 114, 193-198	1.3	5
44	Sojourn times and substitution rate at overdominant and linked neutral loci. <i>Genetics</i> , 2000 , 155, 921-7	4	5
43	Metabolome analysis using cerebrospinal fluid from narcolepsy type 1 patients. <i>Sleep</i> , 2020 , 43,	1.1	4
42	Significance of functional disease-causal/susceptible variants identified by whole-genome analyses for the understanding of human diseases. <i>Proceedings of the Japan Academy Series B: Physical and Biological Sciences</i> , 2017 , 93, 657-676	4	4
41	Increased expression and altered localization of cathepsin Z are associated with progression to jaundice stage in primary biliary cholangitis. <i>Scientific Reports</i> , 2018 , 8, 11808	4.9	4
40	Human lymphocytes obtained from decidual tissue express killer activatory receptors as well as killer inhibitory receptors: analysis using a single strand conformation polymorphism method. <i>American Journal of Reproductive Immunology</i> , 1998 , 39, 271-8	3.8	4
39	Novel single nucleotide polymorphisms (SNPs) at positions 497 (T/G) and 829 (T/C) in the human c-FOS gene and haplotype association. <i>Human Mutation</i> , 2000 , 16, 279	4.7	4
38	Computer simulation analysis suggests weak balancing selection operative at the MICA locus. <i>Hereditas</i> , 2000 , 133, 25-8	2.4	4
37	DNA-based typing of human platelet antigen systems by polymerase chain reaction-single-strand conformation polymorphism method. <i>Vox Sanguinis</i> , 1995 , 69, 347-51	3.1	4
36	Narcolepsy susceptibility gene CCR3 modulates sleep-wake patterns in mice. <i>PLoS ONE</i> , 2017 , 12, e018	7 § 88	4
35	Genome-Wide Association Study Reveals Host Genetic Factors for Liver Diseases. <i>Journal of Clinical and Translational Hepatology</i> , 2013 , 1, 45-50	5.2	4
34	HLA Class I Analysis Provides Insight Into the Genetic and Epigenetic Background of Immune Evasion in Colorectal Cancer With High Microsatellite Instability. <i>Gastroenterology</i> , 2021 ,	13.3	4
33	Epigenome-wide association study of narcolepsy-affected lateral hypothalamic brains, and overlapping DNA methylation profiles between narcolepsy and multiple sclerosis. <i>Sleep</i> , 2020 , 43,	1.1	4
32	The association of integration patterns of human papilloma virus and single nucleotide polymorphisms on immune- or DNA repair-related genes in cervical cancer patients. <i>Scientific Reports</i> , 2019 , 9, 13132	4.9	3
31	Identification of the novel HLA-DQB1 allele, HLA-DQB1*05:03:01:04, in a Japanese individual. <i>Hla</i> , 2019 , 94, 466-468	1.9	3

30	Lessons from Genome-Wide Search for Disease-Related Genes with Special Reference to HLA-Disease Associations. <i>Genes</i> , 2014 , 5, 84-96	4.2	3
29	Selecting a contingency table in a population-based association study: allele frequency or positivity?. <i>Journal of Human Genetics</i> , 1999 , 44, 246-8	4.3	3
28	An optimized genomic VCF workflow for precise identification of Mycobacterium tuberculosis cluster from cross-platform whole genome sequencing data. <i>Infection, Genetics and Evolution</i> , 2020 , 79, 104152	4.5	3
27	A variant at 9q34.11 is associated with HLA-DQB1*06:02 hegative essential hypersomnia. <i>Journal of Human Genetics</i> , 2018 , 63, 1259-1267	4.3	3
26	Genome-wide haplotype association analysis of primary biliary cholangitis risk in Japanese. <i>Scientific Reports</i> , 2018 , 8, 7806	4.9	3
25	Polo-like kinase 4 and Stromal antigen 3 are not associated with recurrent pregnancy loss caused by embryonic aneuploidy. <i>Human Genome Variation</i> , 2020 , 7, 18	1.8	2
24	Detection of the novel HLA-A allele, HLA-A*02:01:01:58, in a Japanese individual. <i>Hla</i> , 2019 , 94, 435-436	1.9	2
23	Discovery of the novel HLA-B allele, HLA-B*51:01:01:36 in a Japanese individual. <i>Hla</i> , 2019 , 94, 445-447	1.9	2
22	Power of association test for detecting minor histocompatibility gene causing graft-versus-host disease following bone marrow transplantation [correction]. <i>Journal of Human Genetics</i> , 2003 , 48, 502-5	d7 ³	2
21	Strong linkage disequilibrium of a HbE variant with the (AT)9(T)5 repeat in the BP1 binding site upstream of the beta-globin gene in the Thai population. <i>Journal of Human Genetics</i> , 2005 , 50, 7-11	4.3	2
20	Sequence analysis of a serological subtype, LA-A9HH, observed in Japanese and the confirmatory sequence of A*2408 <i>Major Histocompatibility Complex</i> , 1996 , 3, 9-14	0.1	2
19	HLA-DRB1 Polymorphism of Balopa Islanders in Papua New Guinea. <i>Anthropological Science</i> , 2003 , 111, 157-164	1.3	2
18	MEL1 at 1p36.3 Is Fused to Several Partner Genes Including HOXA9 Located at Different Chromosomal Bands from 3q21.3 in t(1;3)(p36.3;q21.3)-Leukemia <i>Blood</i> , 2007 , 110, 3499-3499	2.2	2
17	Risk factors associated with large clusters of tuberculosis patients determined by whole-genome sequencing in a high-tuberculosis-burden country. <i>Tuberculosis</i> , 2020 , 125, 101991	2.6	2
16	MGeND: an integrated database for Japanese clinical and genomic information. <i>Human Genome Variation</i> , 2019 , 6, 53	1.8	2
15	CD14 and IL18 gene polymorphisms associated with colorectal cancer subsite risks among atomic bomb survivors. <i>Human Genome Variation</i> , 2015 , 2, 15035	1.8	1
14	Combining effects of polymorphism of tumor necrosis factor IbGflanking region and HLA-DRB1 on radiological progression in patients with rheumatoid arthritis. <i>Modern Rheumatology</i> , 2009 , 19, 134-139	3.3	1
13	DNA typing of HLAA2, A26 and B61 alleles using the PCRMPH method <i>Major Histocompatibility Complex</i> , 1997 , 3, 205-212	0.1	1

LIST OF PUBLICATIONS

12	The genotypes of GYPA and GYPB carrying the MNSs antigens are not associated with cerebral malaria. <i>Journal of Human Genetics</i> , 2007 , 52, 476-479	4.3	1
11	HLA-DRB1 alleles encoding the shared epitope associated with rheumatoid arthritis confer additional susceptibility to seronegative spondyloarthropathies in HLA-B27-positive Japanese individuals. <i>Japanese Journal of Rheumatology</i> , 1999 , 9, 55-64		1
10	Genome-wide association study of idiopathic hypersomnia in a Japanese population. <i>Sleep and Biological Rhythms</i> ,1	1.3	1
9	Mapping of susceptible variants for cold medicine-related Stevens-Johnson syndrome by whole-genome resequencing. <i>Npj Genomic Medicine</i> , 2021 , 6, 9	6.2	1
8	Characterization of intermediate-sized insertions using whole-genome sequencing data and analysis of their functional impact on gene expression. <i>Human Genetics</i> , 2021 , 140, 1201-1216	6.3	О
7	Genome-wide copy number variation analysis of hepatitis B infection in a Japanese population. <i>Human Genome Variation</i> , 2021 , 8, 22	1.8	O
6	A rare genetic variant in the cleavage site of prepro-orexin is associated with idiopathic hypersomnia <i>Npj Genomic Medicine</i> , 2022 , 7, 29	6.2	О
5	A new sister journal to the Journal of Human Genetics-for the interest and benefit of the global community of human genome researchers. <i>Human Genome Variation</i> , 2014 , 1, 14003	1.8	
4	A Subgroup Analysis of Genome-wide Association Study for Panic Disorder. <i>Major Histocompatibility Complex</i> , 2017 , 24, 54-64	0.1	
3	HLA-DRB1 alleles encoding the shared epitope associated with rheumatoid arthritis confer additional susceptibility to seronegative spondyloarthropathies in HLA-B27-positive Japanese individuals. <i>Japanese Journal of Rheumatology</i> , 1999 , 9, 55-64		
2	Genetic polymorphisms in the fourth component of complement (C4) and in properdin factor B (BF) in Japanese patients with palmoplantar pustulosis. <i>Journal of Dermatology</i> , 1992 , 19, 353-5	1.6	
1	Applications of HLA gene polymorphisms. <i>Major Histocompatibility Complex</i> , 2014 , 21, 87-95	0.1	