

Makoto Asashima

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/2400152/makoto-asashima-publications-by-year.pdf>

Version: 2024-04-26

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

106
papers

3,458
citations

29
h-index

55
g-index

106
ext. papers

4,031
ext. citations

5.1
avg, IF

4.55
L-index

#	Paper	IF	Citations
106	Toward global standardization of conducting fair investigations of allegations of research misconduct. <i>Accountability in Research</i> , 2020 , 27, 327-346	1.9	3
105	A Novel Therapeutic Strategy for Pancreatic Cancer: Targeting Cell Surface Glycan Using rBC2LC-N Lectin-Drug Conjugate (LDC). <i>Molecular Cancer Therapeutics</i> , 2018 , 17, 183-195	6.1	30
104	Development of a practical sandwich assay to detect human pluripotent stem cells using cell culture media. <i>Regenerative Therapy</i> , 2017 , 6, 1-8	3.7	5
103	Genome evolution in the allotetraploid frog <i>Xenopus laevis</i> . <i>Nature</i> , 2016 , 538, 336-343	50.4	510
102	Complete mitochondrial genome of "Xenopus tropicalis" Asashima line (Anura: Pipidae), a possible undescribed species. <i>Mitochondrial DNA Part A: DNA Mapping, Sequencing, and Analysis</i> , 2016 , 27, 3341-3 ¹⁻³		2
101	Physicochemical and biological characterizations of Pxt peptides from amphibian (<i>Xenopus tropicalis</i>) skin. <i>Journal of Biochemistry</i> , 2016 , 159, 619-29	3.1	3
100	Functional Overload Enhances Satellite Cell Properties in Skeletal Muscle. <i>Stem Cells International</i> , 2016 , 2016, 7619418	5	13
99	Generation of stomach tissue from mouse embryonic stem cells. <i>Nature Cell Biology</i> , 2015 , 17, 984-93	23.4	71
98	Insulin-like factor regulates neural induction through an IGF1 receptor-independent mechanism. <i>Scientific Reports</i> , 2015 , 5, 11603	4.9	3
97	Establishment and culture optimization of a new type of pituitary immortalized cell line. <i>Biochemical and Biophysical Research Communications</i> , 2015 , 463, 1218-24	3.4	6
96	Identification of novel peptides from amphibian (<i>Xenopus tropicalis</i>) skin by direct tissue MALDI-MS analysis. <i>FEBS Journal</i> , 2015 , 282, 102-13	5.7	6
95	Dullard/Ctdnep1 regulates endochondral ossification via suppression of TGF- β signaling. <i>Journal of Bone and Mineral Research</i> , 2015 , 30, 318-29	6.3	12
94	Diabetes and stem cell function. <i>BioMed Research International</i> , 2015 , 2015, 592915	3	21
93	Elimination of tumorigenic human pluripotent stem cells by a recombinant lectin-toxin fusion protein. <i>Stem Cell Reports</i> , 2015 , 4, 811-20	8	80
92	The requirement of histone modification by PRDM12 and Kdm4a for the development of pre-placodal ectoderm and neural crest in <i>Xenopus</i> . <i>Developmental Biology</i> , 2015 , 399, 164-176	3.1	29
91	Improved efficiency of definitive endoderm induction from human induced pluripotent stem cells in feeder and serum-free culture system. <i>In Vitro Cellular and Developmental Biology - Animal</i> , 2015 , 51, 1-8	2.6	18
90	A stable chimeric fibroblast growth factor (FGF) can successfully replace basic FGF in human pluripotent stem cell culture. <i>PLoS ONE</i> , 2015 , 10, e0118931	3.7	10

89	Lipase member H is a novel secreted protein selectively upregulated in human lung adenocarcinomas and bronchioloalveolar carcinomas. <i>Biochemical and Biophysical Research Communications</i> , 2014 , 443, 1141-7	3.4	13
88	Biosynthesis of ribosomal RNA in nucleoli regulates pluripotency and differentiation ability of pluripotent stem cells. <i>Stem Cells</i> , 2014 , 32, 3099-111	5.8	53
87	MicroRNAs and epigenetics in adult neurogenesis. <i>Advances in Genetics</i> , 2014 , 86, 27-44	3.3	39
86	Occupancy of tissue-specific cis-regulatory modules by Otx2 and TLE/Groucho for embryonic head specification. <i>Nature Communications</i> , 2014 , 5, 4322	17.4	38
85	Wnt protein-mediated satellite cell conversion in adult and aged mice following voluntary wheel running. <i>Journal of Biological Chemistry</i> , 2014 , 289, 7399-412	5.4	49
84	Enhanced bone-forming activity of side population cells in the periodontal ligament. <i>Cell Transplantation</i> , 2014 , 23, 691-701	4	14
83	A medium hyperglycosylated podocalyxin enables noninvasive and quantitative detection of tumorigenic human pluripotent stem cells. <i>Scientific Reports</i> , 2014 , 4, 4069	4.9	22
82	Enzyme-free passage of human pluripotent stem cells by controlling divalent cations. <i>Scientific Reports</i> , 2014 , 4, 4646	4.9	23
81	Prohibitin 2 regulates the proliferation and lineage-specific differentiation of mouse embryonic stem cells in mitochondria. <i>PLoS ONE</i> , 2014 , 9, e81552	3.7	27
80	Microfluidic perfusion culture of human induced pluripotent stem cells under fully defined culture conditions. <i>Biotechnology and Bioengineering</i> , 2014 , 111, 937-47	4.9	33
79	Monitoring neurodegeneration in diabetes using adult neural stem cells derived from the olfactory bulb. <i>Stem Cell Research and Therapy</i> , 2013 , 4, 51	8.3	6
78	N-Cadherin is a prospective cell surface marker of human mesenchymal stem cells that have high ability for cardiomyocyte differentiation. <i>Biochemical and Biophysical Research Communications</i> , 2013 , 438, 753-9	3.4	26
77	The phosphatase Dullard negatively regulates BMP signalling and is essential for nephron maintenance after birth. <i>Nature Communications</i> , 2013 , 4, 1398	17.4	15
76	rBC2LCN, a new probe for live cell imaging of human pluripotent stem cells. <i>Biochemical and Biophysical Research Communications</i> , 2013 , 431, 524-9	3.4	53
75	BMP signaling regulates the differentiation of mouse embryonic stem cells into lung epithelial cell lineages. <i>In Vitro Cellular and Developmental Biology - Animal</i> , 2013 , 49, 230-7	2.6	8
74	Intrinsic ability of adult stem cell in skeletal muscle: an effective and replenishable resource to the establishment of pluripotent stem cells. <i>Stem Cells International</i> , 2013 , 2013, 420164	5	16
73	Podocalyxin is a glycoprotein ligand of the human pluripotent stem cell-specific probe rBC2LCN. <i>Stem Cells Translational Medicine</i> , 2013 , 2, 265-73	6.9	57
72	Activin A regulates growth of gastro-intestinal epithelial cells by mediating epithelial-mesenchymal interaction. <i>Development Growth and Differentiation</i> , 2013 , 55, 786-91	3	7

71	Induction of intermediate mesoderm by retinoic acid receptor signaling from differentiating mouse embryonic stem cells. <i>International Journal of Developmental Biology</i> , 2013 , 57, 383-9	1.9	21
70	Serum protein isoform profiles indicate the progression of hepatitis C virus-induced liver diseases. <i>International Journal of Molecular Medicine</i> , 2013 , 31, 943-50	4.4	2
69	Hippo signaling components, Mst1 and Mst2, act as a switch between self-renewal and differentiation in <i>Xenopus</i> hematopoietic and endothelial progenitors. <i>International Journal of Developmental Biology</i> , 2013 , 57, 407-14	1.9	12
68	A lectin-based glycomic approach to identify characteristic features of <i>Xenopus</i> embryogenesis. <i>PLoS ONE</i> , 2013 , 8, e56581	3.7	6
67	Dullard/Ctdnep1 modulates WNT signalling activity for the formation of primordial germ cells in the mouse embryo. <i>PLoS ONE</i> , 2013 , 8, e57428	3.7	19
66	Xnr3 affects brain patterning via cell migration in the neural-epidermal tissue boundary during early <i>Xenopus</i> embryogenesis. <i>International Journal of Developmental Biology</i> , 2013 , 57, 779-86	1.9	2
65	Characterization of CXC-type chemokine molecules in early <i>Xenopus laevis</i> development. <i>International Journal of Developmental Biology</i> , 2013 , 57, 41-7	1.9	3
64	Novel cell surface genes expressed in the stomach primordium during gastrointestinal morphogenesis of mouse embryos. <i>Gene Expression Patterns</i> , 2012 , 12, 154-63	1.5	2
63	Ubc9 negatively regulates BMP-mediated osteoblastic differentiation in cultured cells. <i>Bone</i> , 2012 , 50, 1092-9	4.7	11
62	mNanog possesses dorsal mesoderm-inducing ability by modulating both BMP and Activin/nodal signaling in <i>Xenopus</i> ectodermal cells. <i>PLoS ONE</i> , 2012 , 7, e46630	3.7	5
61	Induction of differentiation of undifferentiated cells into pancreatic beta cells in vertebrates. <i>International Journal of Developmental Biology</i> , 2012 , 56, 313-23	1.9	25
60	KDEL tagging: a method for generating dominant-negative inhibitors of the secretion of TGF-beta superfamily proteins. <i>International Journal of Developmental Biology</i> , 2012 , 56, 351-6	1.9	7
59	Structural and quantitative evidence for dynamic glycome shift on production of induced pluripotent stem cells. <i>Molecular and Cellular Proteomics</i> , 2012 , 11, 1913-23	7.6	68
58	xCOUP-TF-B regulates xCyp26 transcription and modulates retinoic acid signaling for anterior neural patterning in <i>Xenopus</i> . <i>International Journal of Developmental Biology</i> , 2012 , 56, 239-44	1.9	3
57	Mechanobiology During Vertebrate Organ Development 2011 , 39-47		
56	Rapamycin treatment causes developmental delay, pigmentation defects, and gastrointestinal malformation on <i>Xenopus</i> embryogenesis. <i>Biochemical and Biophysical Research Communications</i> , 2011 , 404, 974-8	3.4	10
55	Development of defective and persistent Sendai virus vector: a unique gene delivery/expression system ideal for cell reprogramming. <i>Journal of Biological Chemistry</i> , 2011 , 286, 4760-71	5.4	251
54	The transcriptional coactivators Yap and TAZ are expressed during early <i>Xenopus</i> development. <i>International Journal of Developmental Biology</i> , 2011 , 55, 121-6	1.9	12

53	Development of Ca ²⁺ (+) signaling mechanisms and cell motility in presumptive ectodermal cells during amphibian gastrulation. <i>Development Growth and Differentiation</i> , 2011 , 53, 37-47	3	5
52	Chemokine ligand Xenopus CXCLC (XCXCLC) regulates cell movements during early morphogenesis. <i>Development Growth and Differentiation</i> , 2011 , 53, 971-81	3	4
51	Possible linkages between the inner and outer cellular states of human induced pluripotent stem cells. <i>BMC Systems Biology</i> , 2011 , 5 Suppl 1, S17	3.5	19
50	Translocon-associated protein subunit Trap-7/Ssr3 is required for vascular network formation in the mouse placenta. <i>Developmental Dynamics</i> , 2011 , 240, 394-403	2.9	15
49	The conserved Rieske oxygenase DAF-36/Neverland is a novel cholesterol-metabolizing enzyme. <i>Journal of Biological Chemistry</i> , 2011 , 286, 25756-62	5.4	118
48	Glycome diagnosis of human induced pluripotent stem cells using lectin microarray. <i>Journal of Biological Chemistry</i> , 2011 , 286, 20345-53	5.4	151
47	In vitro organogenesis using multipotent cells. <i>Human Cell</i> , 2010 , 23, 1-14	4.5	5
46	Claudin5 genes encoding tight junction proteins are required for Xenopus heart formation. <i>Development Growth and Differentiation</i> , 2010 , 52, 665-75	3	7
45	Induction of neural crest cells from mouse embryonic stem cells in a serum-free monolayer culture. <i>International Journal of Developmental Biology</i> , 2010 , 54, 1287-94	1.9	24
44	Reduction of N-glycolylneuraminic acid in human induced pluripotent stem cells generated or cultured under feeder- and serum-free defined conditions. <i>PLoS ONE</i> , 2010 , 5, e14099	3.7	41
43	XRASGRP2 is essential for blood vessel formation during Xenopus development. <i>International Journal of Developmental Biology</i> , 2010 , 54, 609-15	1.9	10
42	Zygotic VegT is required for Xenopus paraxial mesoderm formation and is regulated by Nodal signaling and Eomesodermin. <i>International Journal of Developmental Biology</i> , 2010 , 54, 81-92	1.9	19
41	Xenopus furry contributes to release of microRNA gene silencing. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2010 , 107, 19344-9	11.5	10
40	TIF1beta regulates the pluripotency of embryonic stem cells in a phosphorylation-dependent manner. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2010 , 107, 10926-31	11.5	60
39	Activin plays a key role in the maintenance of long-term memory and late-LTP. <i>Learning and Memory</i> , 2010 , 17, 176-85	2.8	43
38	Pdx1-transfected adipose tissue-derived stem cells differentiate into insulin-producing cells in vivo and reduce hyperglycemia in diabetic mice. <i>International Journal of Developmental Biology</i> , 2010 , 54, 699-705	1.9	61
37	An in vitro reconstitution system for the assessment of chromatin protein fluidity during Xenopus development. <i>Biochemical and Biophysical Research Communications</i> , 2010 , 400, 200-6	3.4	
36	Inhibitory Smad proteins promote the differentiation of mouse embryonic stem cells into ependymal-like ciliated cells. <i>Biochemical and Biophysical Research Communications</i> , 2010 , 401, 1-6	3.4	11

35	BMP4 induction of trophoblast from mouse embryonic stem cells in defined culture conditions on laminin. <i>In Vitro Cellular and Developmental Biology - Animal</i> , 2010 , 46, 416-30	2.6	57
34	Intensely fluorescent azobenzenes: synthesis, crystal structures, effects of substituents, and application to fluorescent vital stain. <i>Chemistry - A European Journal</i> , 2010 , 16, 5026-35	4.8	85
33	Cloning of noggin gene from hydra and analysis of its functional conservation using <i>Xenopus laevis</i> embryos. <i>Evolution & Development</i> , 2010 , 12, 267-74	2.6	11
32	Creating frog heart as an organ: in vitro-induced heart functions as a circulatory organ in vivo. <i>International Journal of Developmental Biology</i> , 2010 , 54, 851-6	1.9	6
31	Directed induction of anterior and posterior primitive streak by Wnt from embryonic stem cells cultured in a chemically defined serum-free medium. <i>FASEB Journal</i> , 2009 , 23, 114-22	0.9	68
30	The RNA-binding protein Mex3b has a fine-tuning system for mRNA regulation in early <i>Xenopus</i> development. <i>Development (Cambridge)</i> , 2009 , 136, 2413-22	6.6	14
29	Molecular analyses of <i>Xenopus laevis</i> Mesp-related genes. <i>Integrative Zoology</i> , 2009 , 4, 387-94	1.9	7
28	Directional migration of neuronal PC12 cells in a ratchet wheel shaped microchamber. <i>Journal of Bioscience and Bioengineering</i> , 2009 , 108, 76-83	3.3	6
27	Separation with zwitterionic hydrophilic interaction liquid chromatography improves protein identification by matrix-assisted laser desorption/ionization-based proteomic analysis. <i>Biomedical Chromatography</i> , 2009 , 23, 607-14	1.7	21
26	Proteomic analysis of membrane proteins expressed specifically in pluripotent murine embryonic stem cells. <i>Proteomics</i> , 2009 , 9, 126-37	4.8	22
25	Decreased expression of CXXC4 promotes a malignant phenotype in renal cell carcinoma by activating Wnt signaling. <i>Oncogene</i> , 2009 , 28, 297-305	9.2	67
24	Bestrophin genes are expressed in <i>Xenopus</i> development. <i>Biochemical and Biophysical Research Communications</i> , 2009 , 384, 290-5	3.4	2
23	The <i>Xenopus</i> Bowline/Ripply family proteins negatively regulate the transcriptional activity of T-box transcription factors. <i>International Journal of Developmental Biology</i> , 2009 , 53, 631-9	1.9	21
22	IGFBP-4 is an inhibitor of canonical Wnt signalling required for cardiogenesis. <i>Nature</i> , 2008 , 454, 345-9	50.4	171
21	A crucial role of a high mobility group protein HMGA2 in cardiogenesis. <i>Nature Cell Biology</i> , 2008 , 10, 567-74	23.4	68
20	Elucidation of the role of activin in organogenesis using a multiple organ induction system with amphibian and mouse undifferentiated cells in vitro. <i>Development Growth and Differentiation</i> , 2008 , 50 Suppl 1, S35-45	3	16
19	Nucleoredoxin regulates the Wnt/planar cell polarity pathway in <i>Xenopus</i> . <i>Genes To Cells</i> , 2008 , 13, 965-75	2.5	24
18	Tbx6, Thylacine1, and E47 synergistically activate bowline expression in <i>Xenopus</i> somitogenesis. <i>Developmental Biology</i> , 2008 , 313, 816-28	3.1	19

17	Physical interaction between Tbx6 and mespb is indispensable for the activation of bowline expression during <i>Xenopus</i> somitogenesis. <i>Biochemical and Biophysical Research Communications</i> , 2008 , 372, 607-12	3.4	6
16	TRIQK, a novel family of small proteins localized to the endoplasmic reticulum membrane, is conserved across vertebrates. <i>Zoological Science</i> , 2008 , 25, 706-13	0.8	4
15	Global expression of simulated microgravity-responsive genes in <i>Xenopus</i> liver cells. <i>Zoological Science</i> , 2008 , 25, 828-37	0.8	4
14	Molecular links among the causative genes for ocular malformation: Otx2 and Sox2 coregulate Rax expression. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2008 , 105, 5408-13	11.5	82
13	Mitf contributes to melanosome distribution and melanophore dendricity. <i>Pigment Cell and Melanoma Research</i> , 2008 , 21, 56-62	4.5	23
12	Retinoic acid metabolizing factor xCyp26c is specifically expressed in neuroectoderm and regulates anterior neural patterning in <i>Xenopus laevis</i> . <i>International Journal of Developmental Biology</i> , 2008 , 52, 893-901	1.9	17
11	XSUMO-1 is required for normal mesoderm induction and axis elongation during early <i>Xenopus</i> development. <i>Developmental Dynamics</i> , 2007 , 236, 2757-66	2.9	13
10	Gene expression profile of <i>Xenopus</i> A6 cells cultured under random positioning machine shows downregulation of ion transporter genes and inhibition of dome formation. <i>Advances in Space Research</i> , 2007 , 40, 1694-1702	2.4	4
9	TSC-box is essential for the nuclear localization and antiproliferative effect of XTSC-22. <i>Development Growth and Differentiation</i> , 2007 , 49, 197-204	3	5
8	<i>Xenopus</i> galectin-VIa shows highly specific expression in cement glands and is regulated by canonical Wnt signaling. <i>Gene Expression Patterns</i> , 2007 , 7, 852-7	1.5	4
7	A novel gene, BENI is required for the convergent extension during <i>Xenopus laevis</i> gastrulation. <i>Developmental Biology</i> , 2007 , 303, 270-80	3.1	2
6	Bowline mediates association of the transcriptional corepressor XGrg-4 with Tbx6 during somitogenesis in <i>Xenopus</i> . <i>Biochemical and Biophysical Research Communications</i> , 2007 , 359, 959-64	3.4	23
5	Ripply2 is essential for precise somite formation during mouse early development. <i>FEBS Letters</i> , 2007 , 581, 2691-6	3.8	30
4	Recombinant Tol2 transposase with activity in <i>Xenopus</i> embryos. <i>FEBS Letters</i> , 2007 , 581, 4333-6	3.8	17
3	N-cadherin is a useful marker for the progenitor of cardiomyocytes differentiated from mouse ES cells in serum-free condition. <i>Biochemical and Biophysical Research Communications</i> , 2006 , 351, 877-82	3.4	22
2	Dullard promotes degradation and dephosphorylation of BMP receptors and is required for neural induction. <i>Developmental Cell</i> , 2006 , 11, 763-74	10.2	65
1	Xapelin and Xmsr are required for cardiovascular development in <i>Xenopus laevis</i> . <i>Developmental Biology</i> , 2006 , 298, 188-200	3.1	69