

Satoshi Oeda

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2401417/publications.pdf>

Version: 2024-02-01

23
papers

553
citations

777949

13
h-index

721071

23
g-index

25
all docs

25
docs citations

25
times ranked

1054
citing authors

#	ARTICLE	IF	CITATIONS
1	Clinical Outcomes in Biopsy-Proven Nonalcoholic Fatty Liver Disease Patients: A Multicenter Registry-based Cohort Study. <i>Clinical Gastroenterology and Hepatology</i> , 2023, 21, 370-379.	2.4	30
2	Ipragliflozin Improves the Hepatic Outcomes of Patients With Diabetes with NAFLD. <i>Hepatology Communications</i> , 2022, 6, 120-132.	2.0	60
3	Trends in hepatocellular carcinoma incident cases in Japan between 1996 and 2019. <i>Scientific Reports</i> , 2022, 12, 1517.	1.6	10
4	Prediction of Nonalcoholic Fatty Liver Disease Using Noninvasive and Non-Imaging Procedures in Japanese Health Checkup Examinees. <i>Diagnostics</i> , 2021, 11, 132.	1.3	18
5	Effect of skin-capsular distance on controlled attenuation parameter for diagnosing liver steatosis in patients with nonalcoholic fatty liver disease. <i>Scientific Reports</i> , 2021, 11, 15641.	1.6	4
6	Accuracy of liver stiffness measurement and controlled attenuation parameter using FibroScan® M/XL probes to diagnose liver fibrosis and steatosis in patients with nonalcoholic fatty liver disease: a multicenter prospective study. <i>Journal of Gastroenterology</i> , 2020, 55, 428-440.	2.3	50
7	Diagnostic Accuracy of FibroScan and Factors Affecting Measurements. <i>Diagnostics</i> , 2020, 10, 940.	1.3	44
8	Metabolomic/lipidomic-based analysis of plasma to diagnose hepatocellular ballooning in patients with nonalcoholic fatty liver disease: A multicenter study. <i>Hepatology Research</i> , 2020, 50, 955-965.	1.8	12
9	Accuracy of the Enhanced Liver Fibrosis test, and combination of the Enhanced Liver Fibrosis and noninvasive tests for the diagnosis of advanced liver fibrosis in patients with nonalcoholic fatty liver disease. <i>Hepatology Research</i> , 2020, 50, 682-692.	1.8	32
10	Discordant pathological diagnosis of nonalcoholic fatty liver disease: A prospective multicenter study. <i>JGH Open</i> , 2020, 4, 497-502.	0.7	22
11	Diagnostic accuracy of FibroScan-AST score to identify nonalcoholic steatohepatitis with significant activity and fibrosis in Japanese patients with nonalcoholic fatty liver disease: Comparison between M and XL probes. <i>Hepatology Research</i> , 2020, 50, 831-839.	1.8	32
12	Generation Gap for Screening and Treatment of Hepatitis C Virus in Saga Prefecture, Japan: An Administrative Database Study of 35,625 Subjects. <i>Internal Medicine</i> , 2020, 59, 169-174.	0.3	3
13	Recommendations from primary care physicians, family, friends and work colleagues influence patients' decisions related to hepatitis screening, medical examinations and antiviral treatment. <i>Experimental and Therapeutic Medicine</i> , 2020, 19, 2973-2982.	0.8	6
14	Predictors of hepatocellular carcinoma recurrence associated with the use of direct-acting antiviral agent therapy for hepatitis C virus after curative treatment: A prospective multicenter cohort study. <i>Cancer Medicine</i> , 2019, 8, 2646-2653.	1.3	27
15	Categorization and Characterization of Activities Designed to Help Health-care Professionals Involved in Hepatitis Care Increase Their Awareness of the Disease: The Classification of Hepatitis Medical Care Coordinators. <i>Internal Medicine</i> , 2019, 58, 1825-1834.	0.3	4
16	Direct-acting antiviral agents do not increase the incidence of hepatocellular carcinoma development: a prospective, multicenter study. <i>Hepatology International</i> , 2019, 13, 293-301.	1.9	38
17	Prevalence of pruritus in patients with chronic liver disease: A multicenter study. <i>Hepatology Research</i> , 2018, 48, E252-E262.	1.8	37
18	Genome-wide association study identified new susceptible genetic variants in HLA class I region for hepatitis B virus-related hepatocellular carcinoma. <i>Scientific Reports</i> , 2018, 8, 7958.	1.6	38

#	ARTICLE	IF	CITATIONS
19	“Weariness” and “unpleasantness” reduce adherence to branched-chain amino acid granules among Japanese patients with liver cirrhosis: results of a single-center cross-sectional survey. <i>Hepatology Research</i> , 2017, 47, E169-E177.	1.8	4
20	Reducing the mortality rate of hepatitis and liver cancer in Japan. <i>Journal of General and Family Medicine</i> , 2017, 18, 205-211.	0.3	6
21	Efficacy of glutathione for the treatment of nonalcoholic fatty liver disease: an open-label, single-arm, multicenter, pilot study. <i>BMC Gastroenterology</i> , 2017, 17, 96.	0.8	65
22	Optimal Follow-up of Patients with Viral Hepatitis Improves the Detection of Early-stage Hepatocellular Carcinoma and the Prognosis of Survival. <i>Internal Medicine</i> , 2016, 55, 2749-2758.	0.3	10
23	The aging of subjects with hepatitis C virus and the impact of dual oral therapy for elder hepatitis C patients. <i>Acta Hepatologica Japonica</i> , 2015, 56, 273-279.	0.0	1