

Rmi Coudroy

List of Publications by Citations

Source: <https://exaly.com/author-pdf/2411745/remi-coudroy-publications-by-citations.pdf>

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

69

papers

1,515

citations

18

h-index

38

g-index

70

ext. papers

1,985

ext. citations

9.3

avg, IF

4.29

L-index

#	Paper	IF	Citations
69	Early use of polymyxin B hemoperfusion in patients with septic shock due to peritonitis: a multicenter randomized control trial. <i>Intensive Care Medicine</i> , 2015 , 41, 975-84	14.5	220
68	Enteral versus parenteral early nutrition in ventilated adults with shock: a randomised, controlled, multicentre, open-label, parallel-group study (NUTRIREA-2). <i>Lancet, The</i> , 2018 , 391, 133-143	40	218
67	Effect of non-invasive oxygenation strategies in immunocompromised patients with severe acute respiratory failure: a post-hoc analysis of a randomised trial. <i>Lancet Respiratory Medicine, the</i> , 2016 , 4, 646-652	35.1	137
66	Sequential application of oxygen therapy via high-flow nasal cannula and noninvasive ventilation in acute respiratory failure: an observational pilot study. <i>Respiratory Care</i> , 2015 , 60, 170-8	2.1	105
65	Predictors of Intubation in Patients With Acute Hypoxemic Respiratory Failure Treated With a Noninvasive Oxygenation Strategy. <i>Critical Care Medicine</i> , 2018 , 46, 208-215	1.4	90
64	High-flow nasal cannula oxygen therapy versus noninvasive ventilation in immunocompromised patients with acute respiratory failure: an observational cohort study. <i>Annals of Intensive Care</i> , 2016 , 6, 45	8.9	70
63	High-flow nasal oxygen therapy and noninvasive ventilation in the management of acute hypoxemic respiratory failure. <i>Annals of Translational Medicine</i> , 2017 , 5, 297	3.2	59
62	Non-invasive ventilation versus high-flow nasal cannula oxygen therapy with apnoeic oxygenation for preoxygenation before intubation of patients with acute hypoxaemic respiratory failure: a randomised, multicentre, open-label trial. <i>Lancet Respiratory Medicine, the</i> , 2019 , 7, 303-312	35.1	58
61	Incidence and impact of skin mottling over the knee and its duration on outcome in critically ill patients. <i>Intensive Care Medicine</i> , 2015 , 41, 452-9	14.5	54
60	Influence of ICU-bed availability on ICU admission decisions. <i>Annals of Intensive Care</i> , 2015 , 5, 55	8.9	36
59	Impact of sleep alterations on weaning duration in mechanically ventilated patients: a prospective study. <i>European Respiratory Journal</i> , 2018 , 51,	13.6	32
58	Clinical features and outcome of patients with acute respiratory failure revealing anti-synthetase or anti-MDA-5 dermatopulmonary syndrome: a French multicenter retrospective study. <i>Annals of Intensive Care</i> , 2018 , 8, 87	8.9	32
57	Functional outcome of prolonged refractory status epilepticus. <i>Critical Care</i> , 2015 , 19, 199	10.8	31
56	Early Identification of Acute Respiratory Distress Syndrome in the Absence of Positive Pressure Ventilation: Implications for Revision of the Berlin Criteria for Acute Respiratory Distress Syndrome. <i>Critical Care Medicine</i> , 2018 , 46, 540-546	1.4	26
55	ADAMTS13 deficiency in severe postpartum HELLP syndrome. <i>British Journal of Haematology</i> , 2013 , 163, 409-10	4.5	25
54	Septic shock with no diagnosis at 24 hours: a pragmatic multicenter prospective cohort study. <i>Critical Care</i> , 2016 , 20, 360	10.8	23
53	Mechanism of airway closure in acute respiratory distress syndrome: a possible role of surfactant depletion. <i>Intensive Care Medicine</i> , 2019 , 45, 290-291	14.5	20

52	Multicentric Standardized Flow Cytometry Routine Assessment of Patients With Sepsis to Predict Clinical Worsening. <i>Chest</i> , 2018 , 154, 617-627	5.3	19
51	Clinical spectrum and short-term outcome of adult patients with purpura fulminans: a French multicenter retrospective cohort study. <i>Intensive Care Medicine</i> , 2018 , 44, 1502-1511	14.5	18
50	Modulation by Polymyxin-B Hemoperfusion of Inflammatory Response Related to Severe Peritonitis. <i>Shock</i> , 2017 , 47, 93-99	3.4	17
49	Reliability of methods to estimate the fraction of inspired oxygen in patients with acute respiratory failure breathing through non-rebreather reservoir bag oxygen mask. <i>Thorax</i> , 2020 , 75, 805-807	7.3	15
48	Role of ICU-acquired weakness on extubation outcome among patients at high risk of reintubation. <i>Critical Care</i> , 2020 , 24, 86	10.8	14
47	Prevalence of Complete Airway Closure According to Body Mass Index in Acute Respiratory Distress Syndrome. <i>Anesthesiology</i> , 2020 , 133, 867-878	4.3	14
46	Preoxygenation with non-invasive ventilation versus high-flow nasal cannula oxygen therapy for intubation of patients with acute hypoxaemic respiratory failure in ICU: the prospective randomised controlled FLORALI-2 study protocol. <i>BMJ Open</i> , 2017 , 7, e018611	3	13
45	Long-term Quality of Life in Adult Patients Surviving Purpura Fulminans: An Exposed-Unexposed Multicenter Cohort Study. <i>Clinical Infectious Diseases</i> , 2019 , 69, 332-340	11.6	13
44	Non-invasive ventilation or high-flow oxygen therapy: When to choose one over the other?. <i>Respirology</i> , 2019 , 24, 724-731	3.6	12
43	High-flow nasal cannula oxygen therapy alone or with non-invasive ventilation during the weaning period after extubation in ICU: the prospective randomised controlled HIGH-WEAN protocol. <i>BMJ Open</i> , 2018 , 8, e023772	3	11
42	In-Hospital Mortality-Associated Factors in Patients With Thrombotic Antiphospholipid Syndrome Requiring ICU Admission. <i>Chest</i> , 2020 , 157, 1158-1166	5.3	9
41	Automated detection and quantification of reverse triggering effort under mechanical ventilation. <i>Critical Care</i> , 2021 , 25, 60	10.8	9
40	Predictors of successful separation from high-flow nasal oxygen therapy in patients with acute respiratory failure: a retrospective monocenter study. <i>Annals of Intensive Care</i> , 2019 , 9, 101	8.9	8
39	Pneumococcal purpura fulminans in asplenic or hyposplenic patients: a French multicenter exposed-unexposed retrospective cohort study. <i>Critical Care</i> , 2020 , 24, 68	10.8	7
38	Influence of Hypoxic Preservation Temperature on Endothelial Cells and Kidney Integrity. <i>BioMed Research International</i> , 2019 , 2019, 8572138	3	7
37	High-flow nasal oxygen therapy alone or with non-invasive ventilation in immunocompromised patients admitted to ICU for acute hypoxemic respiratory failure: the randomised multicentre controlled FLORALI-IM protocol. <i>BMJ Open</i> , 2019 , 9, e029798	3	7
36	Survival in Immunocompromised Patients Ultimately Requiring Invasive Mechanical Ventilation: A Pooled Individual Patient Data Analysis. <i>American Journal of Respiratory and Critical Care Medicine</i> , 2021 , 204, 187-196	10.2	7
35	Duration of diaphragmatic inactivity after endotracheal intubation of critically ill patients. <i>Critical Care</i> , 2021 , 25, 26	10.8	7

34	Effect of non-invasive remote ischemic preconditioning on intra-renal perfusion in volunteers. <i>Journal of Nephrology</i> , 2017 , 30, 393-395	4.8	6
33	Is immunosuppression status a risk factor for noninvasive ventilation failure in patients with acute hypoxemic respiratory failure? A post hoc matched analysis. <i>Annals of Intensive Care</i> , 2019 , 9, 90	8.9	6
32	Benefits and risks of noninvasive oxygenation strategy in COVID-19: a multicenter, prospective cohort study (COVID-ICU) in 137 hospitals. <i>Critical Care</i> , 2021 , 25, 421	10.8	6
31	Acute Respiratory Distress Syndrome: Respiratory Monitoring and Pulmonary Physiology. <i>Seminars in Respiratory and Critical Care Medicine</i> , 2019 , 40, 66-80	3.9	5
30	Impact of sleep alterations on weaning duration of mechanically ventilated patients: how much is bad?. <i>European Respiratory Journal</i> , 2018 , 52,	13.6	5
29	Influence of Noninvasive Ventilation Protocol on Intubation Rates in Subjects With De Novo Respiratory Failure. <i>Respiratory Care</i> , 2020 , 65, 525-534	2.1	4
28	Tidal volume and non-invasive ventilation failure - AuthorsReply. <i>Lancet Respiratory Medicine</i> , 2016 , 4, e52	35.1	4
27	CAPS criteria fail to identify most severely-ill thrombotic antiphospholipid syndrome patients requiring intensive care unit admission. <i>Journal of Autoimmunity</i> , 2019 , 103, 102292	15.5	4
26	Use of Type III procollagen measurement as predictor of lung fibroproliferation in ARDS: early measurement for earlier antifibroproliferative therapy?. <i>Intensive Care Medicine</i> , 2015 , 41, 1159-60	14.5	3
25	Beneficial Effects of Noninvasive Ventilation in Acute Hypoxemic Respiratory Failure: Caution With Findings From Meta-Analyses. <i>Critical Care Medicine</i> , 2017 , 45, e1100-e1101	1.4	3
24	Role of sleep on respiratory failure after extubation in the ICU. <i>Annals of Intensive Care</i> , 2021 , 11, 71	8.9	3
23	Noninvasive Ventilation and Outcomes Among Immunocompromised Patients. <i>JAMA - Journal of the American Medical Association</i> , 2016 , 315, 1901-2	27.4	3
22	High-flow nasal oxygen alone or alternating with non-invasive ventilation in critically ill immunocompromised patients with acute respiratory failure: a randomised controlled trial.. <i>Lancet Respiratory Medicine</i> , 2022 ,	35.1	3
21	Could Noninvasive Ventilation Failure Rates Be Underestimated in the LUNG SAFE Study?. <i>American Journal of Respiratory and Critical Care Medicine</i> , 2017 , 196, 395-396	10.2	2
20	Endotoxin Mass Concentration in Plasma Is Associated With Mortality in a Multicentric Cohort of Peritonitis-Induced Shock. <i>Frontiers in Medicine</i> , 2021 , 8, 749405	4.9	2
19	T-piece versus pressure-support ventilation for spontaneous breathing trials before extubation in patients at high risk of reintubation: protocol for a multicentre, randomised controlled trial (TIP-EX). <i>BMJ Open</i> , 2020 , 10, e042619	3	2
18	Non-invasive ventilation versus high-flow nasal oxygen for postextubation respiratory failure in ICU: a post-hoc analysis of a randomized clinical trial. <i>Critical Care</i> , 2021 , 25, 221	10.8	2
17	Early Identification and Diagnostic Approach in Acute Respiratory Distress Syndrome (ARDS).. <i>Diagnostics</i> , 2021 , 11,	3.8	2

16	High-Flow Oxygen Therapy for Respiratory Failure in Immunocompromised Patients. <i>JAMA - Journal of the American Medical Association</i> , 2019 , 321, 1826-1827	27.4	1
15	Is skin mottling a predictor of high mortality in non-selected patients with cirrhosis admitted to intensive care unit?. <i>Journal of Hepatology</i> , 2015 , 63, 770-1	13.4	1
14	Accuracy of fluoropropyl-2beta-carbomethoxy-3beta-(4-iodophenyl) nortropane datscan for the diagnosis of lewy body dementia. <i>Journal of the American Geriatrics Society</i> , 2010 , 58, 209-10	5.6	1
13	Clinical phenotype and outcomes of pneumococcal versus meningococcal purpura fulminans: a multicenter retrospective cohort study. <i>Critical Care</i> , 2021 , 25, 386	10.8	1
12	One-Year Readmission Following Undifferentiated Acute Hypercapnic Respiratory Failure. <i>COPD: Journal of Chronic Obstructive Pulmonary Disease</i> , 2021 , 1-10	2	1
11	A Singular Spectrum Analysis-Based Data-Driven Technique for the Removal of Cardiogenic Oscillations in Esophageal Pressure Signals. <i>IEEE Journal of Translational Engineering in Health and Medicine</i> , 2020 , 8, 3300211	3	1
10	Noninvasive ventilation vs. high-flow nasal cannula oxygen for preoxygenation before intubation in patients with obesity: a post hoc analysis of a randomized controlled trial. <i>Annals of Intensive Care</i> , 2021 , 11, 114	8.9	1
9	Noninvasive Oxygenation in Patients with Acute Respiratory Failure: Current Perspectives.. <i>International Journal of General Medicine</i> , 2022 , 15, 3121-3132	2.3	0
8	Acute Hypercapnic Respiratory Failure Due to Exacerbation of Chronic Obstructive Pulmonary Disease: Who Are the Patients Most at Risk for Intubation?. <i>Critical Care Medicine</i> , 2015 , 43, e261	1.4	
7	Images in gerontology: a case report. <i>Gerontology</i> , 2010 , 56, 39-40	5.5	
6	Presence of Cryofibrinogen in a Cannabis user with Digital Ischaemia. <i>EJVES Extra</i> , 2010 , 19, e41-e43		
5	High-flow nasal cannula oxygen therapy171-185		
4	Noninvasive ventilation and high-flow nasal oxygen for acute respiratory failure: is less more?. <i>Current Opinion in Critical Care</i> , 2021 , 27, 60-65	3.5	
3	Identifying and Applying Best PEEP in Ventilated Critically Ill Patients 2021 , 685-698		
2	The authors reply. <i>Critical Care Medicine</i> , 2018 , 46, e1019-e1020	1.4	
1	The authors reply. <i>Critical Care Medicine</i> , 2018 , 46, e1012	1.4	