

Yasuhiko Kawakami

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2415381/publications.pdf>

Version: 2024-02-01

88
papers

6,227
citations

81839

39
h-index

69214

77
g-index

93
all docs

93
docs citations

93
times ranked

8568
citing authors

#	ARTICLE	IF	CITATIONS
1	Normal embryonic development and neonatal digit regeneration in mice overexpressing a stem cell factor, Sall4. PLoS ONE, 2022, 17, e0267273.	1.1	2
2	Aberrant Nuclear Translocation of E2F1 and Its Association in Cushing's Disease. Endocrinology, 2022, 163, .	1.4	3
3	A nontoxic fungal natural product modulates fin regeneration in zebrafish larvae upstream of FGF-WNT developmental signaling. Developmental Dynamics, 2021, 250, 160-174.	0.8	6
4	Multimodal Non-Surgical Treatments of Aggressive Pituitary Tumors. Frontiers in Endocrinology, 2021, 12, 624686.	1.5	13
5	The FGF-AKT pathway is necessary for cardiomyocyte survival for heart regeneration in zebrafish. Developmental Biology, 2021, 472, 30-37.	0.9	15
6	Acute elevation of interleukin 6 and matrix metalloproteinase 9 during the onset of pituitary apoplexy in Cushing's disease. Pituitary, 2021, 24, 859-866.	1.6	1
7	Two Distinctive POMC Promoters Modify Gene Expression in Cushing Disease. Journal of Clinical Endocrinology and Metabolism, 2021, 106, e3346-e3363.	1.8	15
8	A Rare Case of Recurrent Pituitary Collision Tumors. Journal of the Endocrine Society, 2020, 4, bvaa089.	0.1	0
9	IRX3/5 regulate mitotic chromatid segregation and limb bud shape. Development (Cambridge), 2020, 147, .	1.2	4
10	Development of the Proximal-Anterior Skeletal Elements in the Mouse Hindlimb Is Regulated by a Transcriptional and Signaling Network Controlled by Sall4. Genetics, 2020, 215, 129-141.	1.2	8
11	Tuba8 Drives Differentiation of Cortical Radial Glia into Apical Intermediate Progenitors by Tuning Modifications of Tubulin C Termini. Developmental Cell, 2020, 52, 477-491.e8.	3.1	7
12	Sall4 regulates neuromesodermal progenitors and their descendants during body elongation in mouse embryos. Development (Cambridge), 2019, 146, .	1.2	22
13	HMGB proteins and arthritis. Human Cell, 2018, 31, 1-9.	1.2	75
14	Gata6 restricts Isl1 to the posterior of nascent hindlimb buds through Isl1 cis-regulatory modules. Developmental Biology, 2018, 434, 74-83.	0.9	6
15	Temporal changes of Sall4 lineage contribution in developing embryos and the contribution of Sall4-lineages to postnatal germ cells in mice. Scientific Reports, 2018, 8, 16410.	1.6	11
16	Corepressor SMRT is required to maintain Hox transcriptional memory during somitogenesis. Proceedings of the National Academy of Sciences of the United States of America, 2018, 115, 10381-10386.	3.3	10
17	Characterization of cis-regulatory elements for Fgf10 expression in the chick embryo. Developmental Dynamics, 2018, 247, 1253-1263.	0.8	1
18	Teratogenic effects of in utero exposure to di-(2-ethylhexyl)-phthalate (DEHP) in B6:129S4 mice. Toxicological Sciences, 2017, 157, kfx019.	1.4	12

#	ARTICLE	IF	CITATIONS
19	The two domain hypothesis of limb prepatterning and its relevance to congenital limb anomalies. Wiley Interdisciplinary Reviews: Developmental Biology, 2017, 6, e270.	5.9	9
20	Cover Image, Volume 6, Issue 4. Wiley Interdisciplinary Reviews: Developmental Biology, 2017, 6, e285.	5.9	0
21	Etv2-miR-130a-Jarid2 cascade regulates vascular patterning during embryogenesis. PLoS ONE, 2017, 12, e0189010.	1.1	22
22	Expression of <i>Noggin</i> and <i>Gremlin1</i> and its implications in fine-tuning BMP activities in mouse cartilage tissues. Journal of Orthopaedic Research, 2017, 35, 1671-1682.	1.2	11
23	Analysis of transcription factors expressed at the anterior mouse limb bud. PLoS ONE, 2017, 12, e0175673.	1.1	13
24	Abstract 448: Etv2-Mir130a-Jarid2 Cascade Regulates Vascular Patterning During Embryogenesis. Arteriosclerosis, Thrombosis, and Vascular Biology, 2017, 37, .	1.1	0
25	Cell migration during heart regeneration in zebrafish. Developmental Dynamics, 2016, 245, 774-787.	0.8	30
26	Endoglin integrates BMP and Wnt signalling to induce haematopoiesis through JDP2. Nature Communications, 2016, 7, 13101.	5.8	18
27	Gata6-Dependent GLI3 Repressor Function is Essential in Anterior Limb Progenitor Cells for Proper Limb Development. PLoS Genetics, 2016, 12, e1006138.	1.5	16
28	Identification and functional characterization of novel transcriptional enhancers involved in regulating human <i>GLI3</i> expression during early development. Development Growth and Differentiation, 2015, 57, 570-580.	0.6	9
29	<i>Sall4-Gli3</i> system in early limb progenitors is essential for the development of limb skeletal elements. Proceedings of the National Academy of Sciences of the United States of America, 2015, 112, 5075-5080.	3.3	52
30	An optical labeling-based proliferation assay system reveals the paracrine effect of interleukin-6 in breast cancer. Biochimica Et Biophysica Acta - Molecular Cell Research, 2015, 1853, 27-40.	1.9	8
31	Regenerative responses after mild heart injuries for cardiomyocyte proliferation in zebrafish. Developmental Dynamics, 2014, 243, 1477-1486.	0.8	18
32	PVT1 dependence in cancer with MYC copy-number increase. Nature, 2014, 512, 82-86.	13.7	617
33	Distinct populations within <i>Isl1</i> lineages contribute to appendicular and facial skeletogenesis through the β -catenin pathway. Developmental Biology, 2014, 387, 37-48.	0.9	15
34	Redefining the Role of Retinoic Acid in Limb Development. Cell Reports, 2013, 3, 1337-1338.	2.9	5
35	<i>Isl1</i> Deletion Causes Kidney Agenesis and Hydronephrosis Resembling CAKUT. Journal of the American Society of Nephrology: JASN, 2013, 24, 1242-1249.	3.0	25
36	<i>Isl1</i> regulates establishment of the posterior hindlimb field upstream of the <i>Hand2</i> - <i>Shh</i> morphoregulatory gene network in mouse embryos. Development (Cambridge), 2012, 139, 1620-1629.	1.2	63

#	ARTICLE	IF	CITATIONS
37	Migration of cardiomyocytes is essential for heart regeneration in zebrafish. <i>Development (Cambridge)</i> , 2012, 139, 4133-4142.	1.2	125
38	Life-long preservation of the regenerative capacity in the fin and heart in zebrafish. <i>Biology Open</i> , 2012, 1, 739-746.	0.6	60
39	β -Catenin signaling specifies progenitor cell identity in parallel with Shh signaling in the developing mammalian thalamus. <i>Development (Cambridge)</i> , 2012, 139, 2692-2702.	1.2	49
40	HMGB factors are required for posterior digit development through integrating signaling pathway activities. <i>Developmental Dynamics</i> , 2011, 240, 1151-1162.	0.8	30
41	Islet1-mediated activation of the β -catenin pathway is necessary for hindlimb initiation in mice. <i>Development (Cambridge)</i> , 2011, 138, 4465-4473.	1.2	51
42	Expression Patterns and Function of Chromatin Protein HMGB2 during Mesenchymal Stem Cell Differentiation. <i>Journal of Biological Chemistry</i> , 2011, 286, 41489-41498.	1.6	47
43	A Src-Tks5 Pathway Is Required for Neural Crest Cell Migration during Embryonic Development. <i>PLoS ONE</i> , 2011, 6, e22499.	1.1	80
44	Designer TGF β Superfamily Ligands with Diversified Functionality. <i>PLoS ONE</i> , 2011, 6, e26402.	1.1	35
45	BMP-2/6 Heterodimer Is More Effective than BMP-2 or BMP-6 Homodimers as Inductor of Differentiation of Human Embryonic Stem Cells. <i>PLoS ONE</i> , 2010, 5, e11167.	1.1	84
46	Bone Morphogenetic Protein-2 and -6 Heterodimer Illustrates the Nature of Ligand-Receptor Assembly. <i>Molecular Endocrinology</i> , 2010, 24, 1469-1477.	3.7	58
47	Chromatin protein HMGB2 regulates articular cartilage surface maintenance via β -catenin pathway. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2009, 106, 16817-16822.	3.3	63
48	Sall genes regulate region-specific morphogenesis in the mouse limb by modulating Hox activities. <i>Development (Cambridge)</i> , 2009, 136, 585-594.	1.2	66
49	Differential activity of Wnt/ β -catenin signaling in the embryonic mouse thalamus. <i>Developmental Dynamics</i> , 2009, 238, 3297-3309.	0.8	39
50	Maintenance of Embryonic Stem Cell Pluripotency by Nanog-Mediated Dedifferentiation of Committed Mesoderm Progenitors. , 2009, , 37-53.		0
51	Molecular Cloning and Developmental Expression of a Hyaluronan and Proteoglycan Link Protein Gene, <i>crtl1/hapln1</i> , in Zebrafish. <i>Zoological Science</i> , 2008, 25, 912-918.	0.3	15
52	Sall genes regulates limb patterning through modulation of region-specific Hox activities in mice. <i>FASEB Journal</i> , 2008, 22, 230.6.	0.2	0
53	miles-apart-Mediated regulation of cell-fibronectin interaction and myocardial migration in zebrafish. <i>Nature Clinical Practice Cardiovascular Medicine</i> , 2007, 4, S77-S82.	3.3	45
54	Lysosomal cathepsins in embryonic programmed cell death. <i>Developmental Biology</i> , 2007, 301, 205-217.	0.9	49

#	ARTICLE	IF	CITATIONS
55	BMP-3 and BMP-6 Structures Illuminate the Nature of Binding Specificity with Receptors. <i>Biochemistry</i> , 2007, 46, 12238-12247.	1.2	96
56	Tbx2 and Tbx3 Regulate the Dynamics of Cell Proliferation during Heart Remodeling. <i>PLoS ONE</i> , 2007, 2, e398.	1.1	82
57	Sp8 exhibits reciprocal induction with Fgf8 but has an opposing effect on anterior-posterior cortical area patterning. <i>Neural Development</i> , 2007, 2, 10.	1.1	115
58	Wnt/beta-catenin signaling regulates vertebrate limb regeneration. <i>Genes and Development</i> , 2006, 20, 3232-3237.	2.7	267
59	Maintenance of embryonic stem cell pluripotency by Nanog-mediated reversal of mesoderm specification. <i>Nature Clinical Practice Cardiovascular Medicine</i> , 2006, 3, S114-S122.	3.3	58
60	Cell lineage transport: a mechanism for molecular gradient formation. <i>Molecular Systems Biology</i> , 2006, 2, 57.	3.2	20
61	Regulation of primary cilia formation and left-right patterning in zebrafish by a noncanonical Wnt signaling mediator, <i>duboraya</i> . <i>Nature Genetics</i> , 2006, 38, 1316-1322.	9.4	117
62	The role of TGF β 2s and Sox9 during limb chondrogenesis. <i>Current Opinion in Cell Biology</i> , 2006, 18, 723-729.	2.6	142
63	Nanog binds to Smad1 and blocks bone morphogenetic protein-induced differentiation of embryonic stem cells. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2006, 103, 10294-10299.	3.3	226
64	Retinoic acid signalling links left-right asymmetric patterning and bilaterally symmetric somitogenesis in the zebrafish embryo. <i>Nature</i> , 2005, 435, 165-171.	13.7	256
65	Expression of Fgf19 in the developing chick eye. <i>Developmental Brain Research</i> , 2005, 156, 104-109.	2.1	28
66	Noncanonical Wnt signaling regulates midline convergence of organ primordia during zebrafish development. <i>Genes and Development</i> , 2005, 19, 164-175.	2.7	146
67	Epicardial retinoid X receptor α is required for myocardial growth and coronary artery formation. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2005, 102, 18455-18460.	3.3	320
68	Transcriptional coactivator PGC-1 α regulates chondrogenesis via association with Sox9. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2005, 102, 2414-2419.	3.3	145
69	The Zebrafish as a Model of Heart Regeneration. <i>Cloning and Stem Cells</i> , 2004, 6, 345-351.	2.6	45
70	Sp8 and Sp9, two closely related buttonhead-like transcription factors, regulate Fgf8 expression and limb outgrowth in vertebrate embryos. <i>Development (Cambridge)</i> , 2004, 131, 4763-4774.	1.2	149
71	Notch activity acts as a sensor for extracellular calcium during vertebrate left-right determination. <i>Nature</i> , 2004, 427, 121-128.	13.7	255
72	Characterization of <i>dermacan</i> , a novel zebrafish lectican gene, expressed in dermal bones. <i>Mechanisms of Development</i> , 2004, 121, 301-312.	1.7	38

#	ARTICLE	IF	CITATIONS
73	MKP3 mediates the cellular response to FGF8 signalling in the vertebrate limb. <i>Nature Cell Biology</i> , 2003, 5, 513-519.	4.6	247
74	Notch activity induces Nodal expression and mediates the establishment of left-right asymmetry in vertebrate embryos. <i>Genes and Development</i> , 2003, 17, 1213-1218.	2.7	171
75	Activation of Notch signaling pathway precedes heart regeneration in zebrafish. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2003, 100, 11889-11895.	3.3	302
76	Expression of the chick vascular endothelial growth factor D gene during limb development. <i>Mechanisms of Development</i> , 2002, 116, 239-242.	1.7	8
77	The limb identity gene <i>Tbx5</i> promotes limb initiation by interacting with <i>Wnt2b</i> and <i>Fgf10</i> . <i>Development (Cambridge)</i> , 2002, 129, 5161-5170.	1.2	161
78	The limb identity gene <i>Tbx5</i> promotes limb initiation by interacting with <i>Wnt2b</i> and <i>Fgf10</i> . <i>Development (Cambridge)</i> , 2002, 129, 5161-70.	1.2	60
79	WNT Signals Control FGF-Dependent Limb Initiation and AER Induction in the Chick Embryo. <i>Cell</i> , 2001, 104, 891-900.	13.5	319
80	Involvement of Frizzled-10 in Wnt-7a signaling during chick limb development. <i>Development Growth and Differentiation</i> , 2000, 42, 561-569.	0.6	33
81	Identification of chick frizzled-10 expressed in the developing limb and the central nervous system. <i>Mechanisms of Development</i> , 2000, 91, 375-378.	1.7	23
82	Involvement of Wnt-5a in chondrogenic pattern formation in the chick limb bud. <i>Development Growth and Differentiation</i> , 1999, 41, 29-40.	0.6	126
83	Sonic hedgehog signaling during digit pattern duplication after application of recombinant protein and expressing cells. <i>Development Growth and Differentiation</i> , 1999, 41, 567-574.	0.6	17
84	Expression of the fibroblast growth factor receptor 1 genes in glomeruli in anti-Thy1.1 mesangial proliferative glomerulonephritis. <i>Virchows Archiv Fur Pathologische Anatomie Und Physiologie Und Fur Klinische Medizin</i> , 1999, 435, 501-508.	1.4	6
85	Cloning and expression pattern of <i>Xenopus prx-1</i> (<i>Xprx-1</i>) during embryonic development. <i>Development Growth and Differentiation</i> , 1998, 40, 97-104.	0.6	11
86	Bone Morphogenetic Protein Signaling Is Required for Maintenance of Differentiated Phenotype, Control of Proliferation, and Hypertrophy in Chondrocytes. <i>Journal of Cell Biology</i> , 1998, 140, 409-418.	2.3	166
87	Differential Expression of the Two Closely Related LIM-Class Homeobox Genes <i>LH-2A</i> and <i>LH-2B</i> during Limb Development. <i>Biochemical and Biophysical Research Communications</i> , 1997, 238, 506-511.	1.0	20
88	Cloning and characterization of <i>Wnt-4</i> and <i>Wnt-11</i> cDNAs from chick embryo. <i>DNA Sequence</i> , 1995, 5, 277-281.	0.7	14