
Ibrahim Rencuzogullari

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/2428305/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Prognostic value of C-reactive protein to albumin ratio for long-term outcomes of patients with peripheral arterial disease underwent endovascular treatment. Vascular, 2022, 30, 481-489.	0.9	9
2	Predictors of left ventricular ejection function decline in young patients with ST-segment elevation myocardial infarction. Revista Da Associação Médica Brasileira, 2022, 68, 802-807.	0.7	24
3	Endothelin-1 and C Reactive Protein as Potential Biomarkers for Restenosis in Patients with Arteriosclerosis Obliterans. Journal of Investigative Surgery, 2021, 34, 771-772.	1.3	3
4	The effect of low flow anesthesia on hemodynamic and peripheral oxygenation parameters in obesity surgery. Journal of King Abdulaziz University, Islamic Economics, 2021, 42, 264-269.	1.1	3
5	Relationship between RS time and the severity of chronic obstructive pulmonary disease. Cukurova Medical Journal, 2021, 46, 756-763.	0.2	0
6	The investigation of TIMI risk index for prediction of contrast-induced acute kidney injury in patients with ST elevation myocardial infarction. Acta Cardiologica, 2020, 75, 77-84.	0.9	12
7	Successful treatment of supravalvular pulmonary membranous stenosis with percutaneous balloon valvuloplasty. Acta Cardiologica, 2020, 75, 473-474.	0.9	0
8	P Wave Peak Time for Predicting an Increased Left Atrial Volume Index in Hemodialysis Patients. Medical Principles and Practice, 2020, 29, 262-269.	2.4	6
9	Reply to the Letter to the Editor Entitled "Serum Osmolarity and Contrast-Induced Nephropathy― Angiology, 2020, 71, 99-100.	1.8	0
10	The predictive value of RS time for short term mortality in patients with acute pulmonary embolism. Journal of Electrocardiology, 2020, 62, 94-99.	0.9	3
11	Reply to the Letter to the Editor Entitled "Comment on â€~Association of Serum Osmolarity With Contrast-Induced Nephropathy in Patients With ST-Segment Elevation Myocardial Infarction'― Angiology, 2020, 71, 671-672.	1.8	0
12	The prognostic value of the serum albumin level for longâ€ŧerm prognosis in patients with acute pulmonary embolism. Clinical Respiratory Journal, 2020, 14, 578-585.	1.6	13
13	Unusual response to Hisâ€refractory atrial premature complex: What is the mechanism?. Journal of Cardiovascular Electrophysiology, 2020, 31, 1232-1234.	1.7	1
14	Relationship between the Severity of Coronary Artery Disease and Catheter-Associated Urethral Stricture in Patients with Acute Coronary Syndrome. The Journal of Tehran Heart Center, 2020, 15, 113-118.	0.3	0
15	Early detection strain/strain rate and time to strain/strain rate abnormalities for left atrial mechanical function in hypertensive patients. Acta Cardiologica, 2019, 74, 141-151.	0.9	6
16	In-hospital and long-term prognoses of patients with a mid-range ejection fraction after an ST-segment myocardial infarction. Acta Cardiologica, 2019, 74, 351-358.	0.9	7
17	Gender disparities in heart failure with mid-range and preserved ejection fraction: results from APOLLON study. Anatolian Journal of Cardiology, 2019, 21, 242-252.	0.9	4
18	Association of Serum Osmolarity With Contrast-Induced Nephropathy in Patients With ST-Segment Elevation Myocardial Infarction. Angiology, 2019, 70, 627-632.	1.8	17

#	Article	IF	CITATIONS
19	Prolonged P wave peak time is associated with the severity of coronary artery disease in patients with non-ST segment elevation myocardial infarction. Journal of Electrocardiology, 2019, 55, 138-143.	0.9	11
20	Assessment of the relationship between preprocedural C-reactive protein/albumin ratio and stent restenosis in patients with ST-segment elevation myocardial infarction. Revista Portuguesa De Cardiologia, 2019, 38, 269-277.	0.5	44
21	The reasons of poor lipid target attainment for secondary prevention in real life practice: Results from EPHESUS. International Journal of Clinical Practice, 2019, 73, 1-9.	1.7	15
22	Prognostic efficacy of C-reactive protein/albumin ratio in ST elevation myocardial infarction. Scandinavian Cardiovascular Journal, 2019, 53, 83-90.	1.2	70
23	Assessment of the relationship between C-reactive protein-to-albumin ratio and slow coronary flow in patients with stable angina pectoris. Coronary Artery Disease, 2019, 30, 505-510.	0.7	11
24	A novel ECG parameter for diagnosis of acute pulmonary embolism: RS time. American Journal of Emergency Medicine, 2019, 37, 1230-1236.	1.6	11
25	The association of PRECISE-DAPT score with development of contrast-induced nephropathy in patients with ST-elevation myocardial infarction undergoing primary percutaneous coronary intervention. Cardiovascular Intervention and Therapeutics, 2019, 34, 207-215.	2.3	26
26	The Predictive Value of PRECISE-DAPT Score for In-Hospital Mortality in Patients With ST-Elevation Myocardial Infarction Undergoing Primary Percutaneous Coronary Intervention. Angiology, 2019, 70, 440-447.	1.8	28
27	The C-Reactive Protein to Albumin Ratio Predicts Acute Kidney Injury in Patients With ST-Segment Elevation Myocardial Infarction Undergoing Primary Percutaneous Coronary Intervention. Heart Lung and Circulation, 2019, 28, 1638-1645.	0.4	36
28	A simple score for the prediction of stent thrombosis in patients with ST elevation myocardial infarction: TIMI risk index. Journal of Cardiovascular and Thoracic Research, 2019, 11, 182-188.	0.9	2
29	Relationship between Câ€reactive protein/albumin ratio and coronary artery disease severity in patients with stable angina pectoris. Journal of Clinical Laboratory Analysis, 2018, 32, e22457.	2.1	61
30	Comparison of SYNTAX score II efficacy with SYNTAX score and TIMI risk score for predicting in-hospital and long-term mortality in patients with ST segment elevation myocardial infarction. International Journal of Cardiovascular Imaging, 2018, 34, 1165-1175.	1.5	20
31	Association between BNP levels and new-onset atrial fibrillation. Herz, 2018, 43, 548-554.	1.1	10
32	Propensity score matching analysis of the impact of Syntax score and Syntax score <scp>II</scp> on new onset atrial fibrillation development in patients with <scp>ST</scp> segment elevation myocardial infarction. Annals of Noninvasive Electrocardiology, 2018, 23, e12504.	1.1	9
33	Fragmented QRS may predict new onset atrial fibrillation in patients with ST-segment elevation myocardial infarction. Journal of Electrocardiology, 2018, 51, 27-32.	0.9	14
34	Successful treatment of massive thrombosis in different locations with prolonged thrombolytic therapy: A life-saving intervention. American Journal of Emergency Medicine, 2018, 36, 1722.e1-1722.e3.	1.6	1
35	The relationship between fragmented QRS complexes and syntax II scores in patients with ST-segment elevation myocardial infarction. Journal of Electrocardiology, 2018, 51, 825-829.	0.9	4
36	Association of Syntax Score II with Contrast-induced Nephropathy and Hemodialysis Requirement in Patients with ST Segment Elevation Myocardial Infarction Undergoing Primary Percutaneous Coronary Intervention. Korean Circulation Journal, 2018, 48, 59.	1.9	15

#	Article	IF	CITATIONS
37	Comparison of syntax score and syntax score II to predict "no reflow phenomenon―in patients with ST-segment elevation myocardial infarction. International Journal of Cardiovascular Imaging, 2017, 33, 1883-1889.	1.5	14
38	Coronary thrombosis in three coronary arteries due to whey protein. American Journal of Emergency Medicine, 2017, 35, 664.e3-664.e4.	1.6	4
39	A Simple and Inexpensive Option for Nonsurgical Septal Reduction in Hypertrophic Obstructive Cardiomyopathy. JACC: Cardiovascular Interventions, 2016, 9, e101-e102.	2.9	1
40	Simple and inexpensive way for the treatment of guidewire-induced distal coronary perforation: subcutaneous fat tissue embolization. Anatolian Journal of Cardiology, 2016, 16, E23-E24.	0.9	1