

Manju Mamtani

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2428558/publications.pdf>

Version: 2024-02-01

15
papers

341
citations

840776

11
h-index

996975

15
g-index

15
all docs

15
docs citations

15
times ranked

939
citing authors

#	ARTICLE	IF	CITATIONS
1	Human Plasma Lipidome Is Pleiotropically Associated With Cardiovascular Risk Factors and Death. <i>Circulation: Cardiovascular Genetics</i> , 2014, 7, 854-863.	5.1	56
2	Genome- and epigenome-wide association study of hypertriglyceridemic waist in Mexican American families. <i>Clinical Epigenetics</i> , 2016, 8, 6.	4.1	52
3	Lipidomic risk score independently and cost-effectively predicts risk of future type 2 diabetes: results from diverse cohorts. <i>Lipids in Health and Disease</i> , 2016, 15, 67.	3.0	44
4	Waist Circumference Independently Associates with the Risk of Insulin Resistance and Type 2 Diabetes in Mexican American Families. <i>PLoS ONE</i> , 2013, 8, e59153.	2.5	40
5	Plasma dihydroceramide species associate with waist circumference in Mexican American families. <i>Obesity</i> , 2014, 22, 950-956.	3.0	32
6	Prevalence of β -thalassaemia in subcastes of Indian Sindhis: Results from a two-phase survey. <i>Public Health</i> , 2007, 121, 193-198.	2.9	24
7	Degree of hyperglycemia independently associates with hospital mortality and length of stay in critically ill, nondiabetic patients: Results from the ANZICS CORE binational registry. <i>Journal of Critical Care</i> , 2020, 55, 149-156.	2.2	16
8	Association of hyperglycaemia with hospital mortality in nondiabetic COVID-19 patients: A cohort study. <i>Diabetes and Metabolism</i> , 2021, 47, 101254.	2.9	16
9	Waist circumference is genetically correlated with incident Type 2 diabetes in Mexican American families. <i>Diabetic Medicine</i> , 2014, 31, 31-35.	2.3	15
10	Bayesian Estimates of the Prevalence of β -Thalassemia Trait in Voluntary Blood Donors of Central India: A Survey. <i>Hemoglobin</i> , 2010, 34, 548-560.	0.8	13
11	Increased waist circumference is independently associated with hypothyroidism in Mexican Americans: replicative evidence from two large, population-based studies. <i>BMC Endocrine Disorders</i> , 2014, 14, 46.	2.2	12
12	Genetic correlation of the plasma lipidome with type 2 diabetes, prediabetes and insulin resistance in Mexican American families. <i>BMC Genetics</i> , 2017, 18, 48.	2.7	10
13	Soluble Forms of Intercellular and Vascular Cell Adhesion Molecules Independently Predict Progression to Type 2 Diabetes in Mexican American Families. <i>PLoS ONE</i> , 2016, 11, e0151177.	2.5	6
14	Genetic association of anthropometric traits with type 2 diabetes in ethnically endogamous Sindhi families. <i>PLoS ONE</i> , 2021, 16, e0257390.	2.5	3
15	Association of Urinary Phthalates with Self-Reported Eye Affliction/Retinopathy in Individuals with Diabetes: National Health and Nutrition Examination Survey, 2001-2010. <i>Journal of Diabetes Research</i> , 2016, 2016, 1-10.	2.3	2