

Jarosław Skłodowski

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/242998/publications.pdf>

Version: 2024-02-01

13
papers

149
citations

1307594

7
h-index

1199594

12
g-index

13
all docs

13
docs citations

13
times ranked

136
citing authors

#	ARTICLE	IF	CITATIONS
1	Responses of ground beetles (Coleoptera, Carabidae) to tree retention groups of various sizes support leaving them in clear-cut areas. <i>Forest Ecology and Management</i> , 2021, 493, 119261.	3.2	4
2	Litter-dwelling beetles (Insecta: Coleoptera) can survive in clear-cutting during subsequent soil ploughing. <i>Agricultural and Forest Entomology</i> , 2020, 22, 50-60.	1.3	1
3	Survival of carabids after windthrow of pine forest depends on the presence of broken tree crowns. <i>Scandinavian Journal of Forest Research</i> , 2020, 35, 10-19.	1.4	8
4	Consequences for millipedes (Myriapoda, Diplopoda) of transforming a primeval forest into a managed forest – A case study from Białowieża (Poland). <i>Forest Ecology and Management</i> , 2018, 409, 593-600.	3.2	3
5	Carabids benefit more from pine stands with added understory or second story of broad-leaved trees favored by climate change than from one-storied pine stands. <i>European Journal of Forest Research</i> , 2018, 137, 745-757.	2.5	12
6	Divergence of soil microarthropod (Hexapoda: Collembola) recovery patterns during natural regeneration and regeneration by planting of windthrown pine forests. <i>Forest Ecology and Management</i> , 2018, 429, 414-424.	3.2	5
7	Manual soil preparation and piles of branches can support ground beetles (Coleoptera, carabidae) better than four different mechanical soil treatments in a clear-cut area of a closed-canopy pine forest in northern Poland. <i>Scandinavian Journal of Forest Research</i> , 2017, 32, 123-133.	1.4	20
8	Three phases of changes in carabid assemblages during secondary succession in a pine forest disturbed by windthrow – results from the first 10 years of observations. <i>Insect Conservation and Diversity</i> , 2017, 10, 449-461.	3.0	13
9	Effects of Topsoil Preparation and Broad-leaved Tree Mixture on Carabid Beetles in Afforested Fallow Plots. <i>Restoration Ecology</i> , 2014, 22, 13-21.	2.9	11
10	Consequence of the transformation of a primeval forest into a managed forest for carabid beetles (Coleoptera: Carabidae) - a case study from Białowieża (Poland). <i>European Journal of Entomology</i> , 2014, 111, 639-648.	1.2	29
11	Ground beetle (Coleoptera, Carabidae) assemblages inhabiting Scots pine stands of Puszcza Piska Forest: six-year responses to a tornado impact. <i>ZooKeys</i> , 2011, 100, 371-392.	1.1	17
12	Anthropogenic transformation of ground beetle assemblages (Coleoptera: Carabidae) in Białowieża Forest, Poland: from primeval forests to managed woodlands of various ages. <i>Entomologica Fennica</i> , 2006, 17, 296-314.	0.6	26
13	Four phases of changes in carabid assemblages during secondary succession in a pine forest disturbed by windthrow. <i>ARPHA Conference Abstracts</i> , 0, 2, .	0.0	0