Shinya Tanaka

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/2431676/publications.pdf

Version: 2024-02-01

		567144	477173
33	867	15	29
papers	citations	h-index	g-index
34	34	34	1283
all docs	docs citations	times ranked	citing authors

#	Article	IF	CITATIONS
1	Low-intensity resistance training with blood flow restriction improves vascular endothelial function and peripheral blood circulation in healthy elderly people. European Journal of Applied Physiology, 2016, 116, 749-757.	1.2	119
2	Gait speed has comparable prognostic capability to six-minute walk distance in older patients with cardiovascular disease. European Journal of Preventive Cardiology, 2018, 25, 212-219.	0.8	92
3	Quadriceps Strength as a Predictor of Mortality in Coronary Artery Disease. American Journal of Medicine, 2015, 128, 1212-1219.	0.6	85
4	Utility of SARC-F for Assessing Physical Function in Elderly Patients With Cardiovascular Disease. Journal of the American Medical Directors Association, 2017, 18, 176-181.	1.2	79
5	Prognostic Value of Psoas Muscle Area and Density in Patients Who Undergo Cardiovascular Surgery. Canadian Journal of Cardiology, 2017, 33, 1652-1659.	0.8	71
6	Complementary Role of Arm Circumference to Body Mass Index inÂRiskÂStratification in Heart Failure. JACC: Heart Failure, 2016, 4, 265-273.	1.9	46
7	Prognostic Usefulness of Arm and Calf Circumference in Patients ≥65ÂYears of Age With Cardiovascular Disease. American Journal of Cardiology, 2017, 119, 186-191.	0.7	41
8	Six-Minute Walk Distance Is an Independent Predictor of Hospital Readmission in Patients With Chronic Heart Failure. International Heart Journal, 2014, 55, 331-336.	0.5	33
9	Stretching Exercises Improve Vascular Endothelial Dysfunction Through Attenuation of Oxidative Stress in Chronic Heart Failure Patients With an Implantable Cardioverter Defibrillator. Journal of Cardiopulmonary Rehabilitation and Prevention, 2017, 37, 130-138.	1.2	33
10	Incremental Value of Objective Frailty Assessment to Predict Mortality in Elderly Patients Hospitalized for Heart Failure. Journal of Cardiac Failure, 2018, 24, 723-732.	0.7	32
11	Short-Term Change in Gait Speed and Clinical Outcomes in Older Patients With Acute Heart Failure. Circulation Journal, 2019, 83, 1860-1867.	0.7	27
12	Prognostic value of sarcopenic obesity estimated by computed tomography in patients with cardiovascular disease and undergoing surgery. Journal of Cardiology, 2019, 74, 273-278.	0.8	20
13	Prevalence and prognosis of respiratory muscle weakness in heart failure patients with preserved ejection fraction. Respiratory Medicine, 2020, 161, 105834.	1.3	19
14	Prevalence and prognostic value of the coexistence of anaemia and frailty in older patients with heart failure. ESC Heart Failure, 2021, 8, 625-633.	1.4	19
15	Preoperative paraspinous muscle sarcopenia and physical performance as prognostic indicators in nonâ€smallâ€cell lung cancer. Journal of Cachexia, Sarcopenia and Muscle, 2021, 12, 646-656.	2.9	16
16	A Single Session of Neuromuscular Electrical Stimulation Enhances Vascular Endothelial Function and Peripheral Blood Circulation in Patients With Acute Myocardial Infarction. International Heart Journal, 2016, 57, 676-681.	0.5	15
17	Safety of neuromuscular electrical stimulation in patients implanted with cardioverter defibrillators. Journal of Electrocardiology, 2016, 49, 99-101.	0.4	15
18	Short physical performance battery discriminates clinical outcomes in hospitalized patients aged 75 years and over. Archives of Gerontology and Geriatrics, 2020, 90, 104155.	1.4	15

#	Article	IF	CITATIONS
19	Changes in Respiratory Muscle Strength Following Cardiac Rehabilitation for Prognosis in Patients with Heart Failure. Journal of Clinical Medicine, 2020, 9, 952.	1.0	14
20	Effects of Acute Phase Intensive Electrical Muscle Stimulation in Frail Elderly Patients With Acute Heart Failure (ACTIVEâ€EMS): Rationale and protocol for a multicenter randomized controlled trial. Clinical Cardiology, 2017, 40, 1189-1196.	0.7	11
21	Multidomain Frailty in Heart Failure: Current Status and Future Perspectives. Current Heart Failure Reports, 2021, 18, 107-120.	1.3	11
22	Effects of electrical muscle stimulation on physical function in frail older patients with acute heart failure: a randomized controlled trial. European Journal of Preventive Cardiology, 2022, 29, e286-e288.	0.8	10
23	SARCâ€F questionnaire identifies physical limitations and predicts post discharge outcomes in elderly patients with cardiovascular disease. JCSM Clinical Reports, 2018, 3, 1-11.	0.5	7
24	Excessive SBP elevation during moderate exercise discriminates patients at high risk of developing left ventricular hypertrophy from hypertensive patients. Journal of Hypertension, 2018, 36, 1291-1298.	0.3	6
25	Office-Based Physical Assessment in Patients Aged 75 Years and Older with Cardiovascular Disease. Gerontology, 2019, 65, 128-135.	1.4	6
26	Prognostic value of pupil area for all ause mortality in patients with heart failure. ESC Heart Failure, 2020, 7, 3067-3074.	1.4	5
27	Low ankle brachial index is associated with the magnitude of impaired walking endurance in patients with heart failure. International Journal of Cardiology, 2016, 224, 400-405.	0.8	4
28	Muscle Weakness Is Associated With an Increase of Left Ventricular Mass Through Excessive Blood Pressure Elevation During Exercise in Patients With Hypertension. International Heart Journal, 2017, 58, 551-556.	0.5	4
29	Pupillary Light Reflex as a New Prognostic Marker in Patients With Heart Failure. Journal of Cardiac Failure, 2019, 25, 156-163.	0.7	4
30	Efficacy and Safety of Acute Phase Intensive Electrical Muscle Stimulation in Frail Older Patients with Acute Heart Failure: Results from the ACTIVE-EMS Trial. Journal of Cardiovascular Development and Disease, 2022, 9, 99.	0.8	4
31	Effects of electrical muscle stimulation in frail elderly patients during haemodialysis (DIAL): rationale and protocol for a crossover randomised controlled trial. BMJ Open, 2019, 9, e025389.	0.8	3
32	Association between chronic kidney disease and physical activity level in patients with ischemic heart disease. Renal Replacement Therapy, 2017, 3, .	0.3	0
33	Decreased level of serum carnitine might lead to arteriosclerosis progression via the accumulation of advanced glycation end products in maintenance hemodialysis patients. Renal Replacement Therapy, 2017, 3, .	0.3	0