

Martin Lipp

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2436152/publications.pdf>

Version: 2024-02-01

151
papers

28,249
citations

15001

68
h-index

9346

148
g-index

152
all docs

152
docs citations

152
times ranked

28068
citing authors

#	ARTICLE	IF	CITATIONS
1	S1P Signalling Differentially Affects Migration of Peritoneal B Cell Populations In Vitro and Influences the Production of Intestinal IgA In Vivo. International Journal of Molecular Sciences, 2018, 19, 391.	1.8	20
2	Cytotoxic T cells modulate inflammation and endogenous opioid analgesia in chronic arthritis. Journal of Neuroinflammation, 2017, 14, 30.	3.1	38
3	The transcriptional coactivator Bob1 promotes the development of follicular T helper cells via Bcl6. EMBO Journal, 2016, 35, 881-898.	3.5	44
4	Dysregulated development of IL-17A and IL-21-expressing follicular helper T cells and increased germinal center formation in the absence of ROR γ t. FASEB Journal, 2016, 30, 761-774.	0.2	24
5	MicroRNA-34a promotes genomic instability by a broad suppression of genome maintenance mechanisms downstream of the oncogene KSHV-vGPCR. Oncotarget, 2016, 7, 10414-10432.	0.8	15
6	Suppression of Peripheral Pain by Blockade of Voltage-Gated Calcium 2.2 Channels in Nociceptors Induces RANKL and Impairs Recovery From Inflammatory Arthritis in a Mouse Model. Arthritis and Rheumatology, 2015, 67, 1657-1667.	2.9	11
7	Potent anti-tumor response by targeting B cell maturation antigen (BCMA) in a mouse model of multiple myeloma. Molecular Oncology, 2015, 9, 1348-1358.	2.1	27
8	The Homeostatic Chemokine CCL21 Predicts Mortality in Aortic Stenosis Patients and Modulates Left Ventricular Remodeling. PLoS ONE, 2014, 9, e112172.	1.1	21
9	Follicular regulatory T cells control humoral autoimmunity via NFAT2-regulated CXCR5 expression. Journal of Experimental Medicine, 2014, 211, 545-561.	4.2	147
10	B-cell-intrinsic STAT6 signaling controls germinal center formation. European Journal of Immunology, 2014, 44, 2130-2138.	1.6	63
11	Immunotherapy of B-cell non-Hodgkin lymphoma by targeting the chemokine receptor CXCR5 in a preclinical mouse model. International Journal of Cancer, 2014, 135, 2623-2632.	2.3	20
12	Increased levels of CCR7 ligands in carotid atherosclerosis: different effects in macrophages and smooth muscle cells. Cardiovascular Research, 2014, 102, 148-156.	1.8	37
13	Access to Follicular Dendritic Cells Is a Pivotal Step in Murine Chronic Lymphocytic Leukemia B-cell Activation and Proliferation. Cancer Discovery, 2014, 4, 1448-1465.	7.7	60
14	Dendritic cell-mediated survival signals in 1/4-Myc B-cell lymphoma depend on the transcription factor C/EBP β . Nature Communications, 2014, 5, 5057.	5.8	17
15	Transition from an autoimmune-prone state to fatal autoimmune disease in CCR7 and ROR γ t double-deficient mice is dependent on gut microbiota. Journal of Autoimmunity, 2013, 47, 58-72.	3.0	13
16	Immunotherapy of B-Cell Lymphoma with an Engineered Bispecific Antibody Targeting CD19 and CD5. Antibodies, 2013, 2, 338-352.	1.2	4
17	A chronic model of arthritis supported by a strain-specific periarticular lymph node in BALB/c mice. Nature Communications, 2013, 4, 1644.	5.8	18
18	CCR7 deficiency causes diarrhea associated with altered ion transport in colonocytes in the absence of overt colitis. Mucosal Immunology, 2012, 5, 377-387.	2.7	8

#	ARTICLE	IF	CITATIONS
19	HER2/neu DNA vaccination by intradermal gene delivery in a mouse tumor model. <i>Oncolmunology</i> , 2012, 1, 1537-1545.	2.1	30
20	CCR7 with S1P1 Signaling through AP-1 for Migration of Foxp3+ Regulatory T-Cells Controls Autoimmune Exocrinopathy. <i>American Journal of Pathology</i> , 2012, 180, 199-208.	1.9	23
21	CCL19 as an adjuvant for intradermal gene gun immunization in a Her2/neu mouse tumor model: improved vaccine efficacy and a role for B cells as APC. <i>Cancer Gene Therapy</i> , 2012, 19, 880-887.	2.2	27
22	CCR9+ Macrophages Are Required for Acute Liver Inflammation in Mouse Models of Hepatitis. <i>Gastroenterology</i> , 2012, 142, 366-376.	0.6	72
23	The Homeostatic Chemokine CCL21 Predicts Mortality and May Play a Pathogenic Role in Heart Failure. <i>PLoS ONE</i> , 2012, 7, e33038.	1.1	33
24	CCL21 (SLC) improves tumor protection by a DNA vaccine in a Her2/neu mouse tumor model. <i>Cancer Gene Therapy</i> , 2012, 19, 69-76.	2.2	34
25	CCL19 (ELC) improves TH1-polarized immune responses and protective immunity in a murine Her2/neu DNA vaccination model. <i>Journal of Gene Medicine</i> , 2012, 14, 128-137.	1.4	20
26	The role of CCR7 in allergic airway inflammation induced by house dust mite exposure. <i>Cellular Immunology</i> , 2012, 275, 24-32.	1.4	22
27	Luminal CD4+ T Cells Penetrate Gut Epithelial Monolayers and Egress From Lamina Propria to Blood Circulation. <i>Gastroenterology</i> , 2011, 141, 2130-2139.e11.	0.6	9
28	Manifestation of Spontaneous and Early Autoimmune Gastritis in CCR7-Deficient Mice. <i>American Journal of Pathology</i> , 2011, 179, 754-765.	1.9	20
29	Cooperative function of CCR7 and lymphotoxin in the formation of a lymphoma-permissive niche within murine secondary lymphoid organs. <i>Blood</i> , 2011, 118, 1020-1033.	0.6	57
30	Steady state migratory RelB ⁺ langerin ⁺ dermal dendritic cells mediate peripheral induction of antigen-specific CD4 ⁺ CD25 ⁺ Foxp3 ⁺ regulatory T cells. <i>European Journal of Immunology</i> , 2011, 41, 1420-1434.	1.6	76
31	Lack of CCR7 induces pulmonary hypertension involving perivascular leukocyte infiltration and inflammation. <i>American Journal of Physiology - Lung Cellular and Molecular Physiology</i> , 2011, 301, L50-L59.	1.3	28
32	Sphingosine-1-phosphate receptor 4 (S1P ₄) deficiency profoundly affects dendritic cell function and T _H 17 cell differentiation in a murine model. <i>FASEB Journal</i> , 2011, 25, 4024-4036.	0.2	104
33	Coordinated Regulation of Lymph Node Vascular Stromal Growth First by CD11c+ Cells and Then by T and B Cells. <i>Journal of Immunology</i> , 2011, 187, 5558-5567.	0.4	109
34	Lack of Chemokine Signaling through CXCR5 Causes Increased Mortality, Ventricular Dilatation and Deranged Matrix during Cardiac Pressure Overload. <i>PLoS ONE</i> , 2011, 6, e18668.	1.1	30
35	Shaping Up Adaptive Immunity: The Impact of CCR7 and CXCR5 on Lymphocyte Trafficking. <i>Microcirculation</i> , 2010, 10, 325-334.	1.0	37
36	The chemokine receptor CXCR5 is pivotal for ectopic mucosa-associated lymphoid tissue neogenesis in chronic <i>Helicobacter pylori</i> -induced inflammation. <i>Journal of Molecular Medicine</i> , 2010, 88, 1169-1180.	1.7	57

#	ARTICLE	IF	CITATIONS
37	In Situ Patrolling of Regulatory T Cells Is Essential for Protecting Autoimmune Exocrinopathy. PLoS ONE, 2010, 5, e8588.	1.1	18
38	Shaping of terminal megakaryocyte differentiation and proplatelet development by sphingosine-1-phosphate receptor S1P ₄ . FASEB Journal, 2010, 24, 4701-4710.	0.2	75
39	Shaping of terminal megakaryocyte differentiation and proplatelet development by sphingosine-1-phosphate receptor S1P ₄ . FASEB Journal, 2010, 24, 4701-4710.	0.2	10
40	Adaptive peripheral immune response increases proliferation of neural precursor cells in the adult hippocampus. FASEB Journal, 2009, 23, 3121-3128.	0.2	69
41	CCR7 Ligands Are Required for Development of Experimental Autoimmune Encephalomyelitis through Generating IL-23-Dependent Th17 Cells. Journal of Immunology, 2009, 183, 2513-2521.	0.4	69
42	CCR7-deficient mice develop atypically persistent germinal centers in response to thymus-independent type 2 antigens. Journal of Leukocyte Biology, 2009, 85, 409-417.	1.5	6
43	CCR7-mediated migration of developing thymocytes to the medulla is essential for negative selection to tissue-restricted antigens. Proceedings of the National Academy of Sciences of the United States of America, 2009, 106, 17129-17133.	3.3	109
44	Identification of a chemokine receptor profile characteristic for mediastinal large B-cell lymphoma. International Journal of Cancer, 2009, 125, 2367-2374.	2.3	19
45	CCR7 signalling as an essential regulator of CNS infiltration in T-cell leukaemia. Nature, 2009, 459, 1000-1004.	13.7	227
46	Chemokine CXCL13 is essential for lymph node initiation and is induced by retinoic acid and neuronal stimulation. Nature Immunology, 2009, 10, 1193-1199.	7.0	266
47	Impaired Effector Memory T-Cell Regulation Facilitates Graft Versus Host Disease in CCR7-Deficient Bone Marrow Transplant Chimeras. Transplantation, 2009, 88, 631-639.	0.5	6
48	CCR7 regulates lymphocyte egress and recirculation through body cavities. Journal of Leukocyte Biology, 2009, 87, 671-682.	1.5	32
49	Thymocyte-Dendritic Cell Interactions near Sources of CCR7 Ligands in the Thymic Cortex. Journal of Immunology, 2008, 181, 7014-7023.	0.4	56
50	Mycobacterium tuberculosis Triggers Formation of Lymphoid Structure in Murine Lungs. Journal of Infectious Diseases, 2007, 195, 46-54.	1.9	132
51	CCR7 is required for the in vivo function of CD4 ⁺ CD25 ⁺ regulatory T cells. Journal of Experimental Medicine, 2007, 204, 735-745.	4.2	282
52	CCR7 deficiency causes ectopic lymphoid neogenesis and disturbed mucosal tissue integrity. Blood, 2007, 109, 886-895.	0.6	54
53	Chemokines CCL19 and CCL21 promote activation-induced cell death of antigen-responding T cells. Blood, 2007, 109, 449-456.	0.6	31
54	CXCR5- and CCR7-dependent lymphoid neogenesis in a murine model of chronic antigen-induced arthritis. Arthritis and Rheumatism, 2007, 56, 3271-3283.	6.7	97

#	ARTICLE	IF	CITATIONS
55	The germinal center response is impaired in the absence of T cell-expressed CXCR5. <i>European Journal of Immunology</i> , 2007, 37, 100-109.	1.6	107
56	Distinctive role of CCR7 in migration and functional activity of naive- and effector/memory-like Treg subsets. <i>European Journal of Immunology</i> , 2007, 37, 1575-1583.	1.6	142
57	Enhanced tumorigenicity of fibroblasts transformed with human herpesvirus 8 chemokine receptor vGPCR by successive passage in nude and immunocompetent mice. <i>Oncogene</i> , 2007, 26, 5702-5712.	2.6	6
58	CCL19 (ELC) as an adjuvant for DNA vaccination: induction of a TH1-type T-cell response and enhancement of antitumor immunity. <i>Cancer Gene Therapy</i> , 2007, 14, 523-532.	2.2	29
59	Secondary lymphoid tissue chemokine (SLC/CCL21)/CCR7 signaling regulates fibrocytes in renal fibrosis. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2006, 103, 14098-14103.	3.3	247
60	CCR7-Dependent Cortex-to-Medulla Migration of Positively Selected Thymocytes Is Essential for Establishing Central Tolerance. <i>Immunity</i> , 2006, 24, 165-177.	6.6	260
61	Coordination between CCR7- and CCR9-mediated chemokine signals in prevascular fetal thymus colonization. <i>Blood</i> , 2006, 108, 2531-2539.	0.6	175
62	Selectin Ligand-Independent Priming and Maintenance of T Cell Immunity during Airborne Tuberculosis. <i>Journal of Immunology</i> , 2006, 176, 1131-1140.	0.4	31
63	Follicular B helper T cell activity is confined to CXCR5 ^{hi} ICOS ^{hi} CD4 T cells and is independent of CD57 expression. <i>European Journal of Immunology</i> , 2006, 36, 1892-1903.	1.6	307
64	Salmonella typhimurium infection triggers dendritic cells and macrophages to adopt distinct migration patterns in vivo. <i>European Journal of Immunology</i> , 2006, 36, 2939-2950.	1.6	25
65	CCR7 Is Critically Important for Migration of Dendritic Cells in Intestinal Lamina Propria to Mesenteric Lymph Nodes. <i>Journal of Immunology</i> , 2006, 176, 803-810.	0.4	381
66	The role of CCL21 in recruitment of T-precursor cells to fetal thymus. <i>Blood</i> , 2005, 105, 31-39.	0.6	126
67	BCA-1/CXCL13 expression is associated with CXCR5-positive B-cell cluster formation in acute renal transplant rejection. <i>Kidney International</i> , 2005, 67, 1616-1621.	2.6	65
68	Chemokine receptor CCR7 required for T lymphocyte exit from peripheral tissues. <i>Nature Immunology</i> , 2005, 6, 889-894.	7.0	434
69	The ratio between dendritic cells and T _H 1 cells determines the outcome of their encounter: Proliferation versus deletion. <i>European Journal of Immunology</i> , 2005, 35, 2851-2863.	1.6	55
70	Chemokine-Controlled Migration in Lymphoid Organogenesis and the Systemic Organization of Immunity. <i>Journal of Immunology</i> , 2005, 175, 251-273.		0
71	CXCR5-Dependent Seeding of Follicular Niches by B and Th Cells Augments Antiviral B Cell Responses. <i>Journal of Immunology</i> , 2005, 175, 7109-7116.	0.4	68
72	Differential requirements for the chemokine receptor CCR7 in T cell activation during <i>Listeria monocytogenes</i> infection. <i>Journal of Experimental Medicine</i> , 2005, 201, 1447-1457.	4.2	60

#	ARTICLE	IF	CITATIONS
73	Requirements for Follicular Exclusion and Competitive Elimination of Autoantigen-Binding B Cells. <i>Journal of Immunology</i> , 2004, 172, 4700-4708.	0.4	80
74	A Novel Model for Lymphocytic Infiltration of the Thyroid Gland Generated by Transgenic Expression of the CC Chemokine CCL21. <i>Journal of Immunology</i> , 2004, 173, 4791-4798.	0.4	81
75	Impact of CCR7 on Priming and Distribution of Antiviral Effector and Memory CTL. <i>Journal of Immunology</i> , 2004, 173, 6684-6693.	0.4	87
76	Distinct and overlapping roles of CXCR5 and CCR7 in B-1 cell homing and early immunity against bacterial pathogens. <i>Journal of Leukocyte Biology</i> , 2004, 76, 709-718.	1.5	43
77	Lymphoid organogenesis: getting the green light from ROR γ t. <i>Nature Immunology</i> , 2004, 5, 12-14.	7.0	19
78	Induced recruitment of NK cells to lymph nodes provides IFN- γ for TH1 priming. <i>Nature Immunology</i> , 2004, 5, 1260-1265.	7.0	1,225
79	Subversion of effector CD8+ T α cell differentiation in acute hepatitis C virus infection: exploring the immunological mechanisms. <i>European Journal of Immunology</i> , 2004, 34, 427-437.	1.6	68
80	The chemokine receptor CCR7 controls lymph node-dependent cytotoxic T cell priming in alloimmune responses. <i>European Journal of Immunology</i> , 2004, 34, 461-470.	1.6	51
81	CCR7 Signals Are Essential for Cortical-Medulla Migration of Developing Thymocytes. <i>Journal of Experimental Medicine</i> , 2004, 200, 493-505.	4.2	349
82	HCMV-encoded chemokine receptor US28 employs multiple routes for internalization. <i>Biochemical and Biophysical Research Communications</i> , 2004, 322, 42-49.	1.0	35
83	All Roads Lead to Rome. <i>Immunity</i> , 2004, 20, 244-246.	6.6	17
84	ICOS+ T α cells produce distinct cytokines in different mucosal immune responses. <i>European Journal of Immunology</i> , 2003, 33, 392-401.	1.6	45
85	The sphingosine 1-phosphate receptor S1P4 regulates cell shape and motility via coupling to Giand G12/13. <i>Journal of Cellular Biochemistry</i> , 2003, 89, 507-519.	1.2	117
86	The impact of CCR7 and CXCR5 on lymphoid organ development and systemic immunity. <i>Immunological Reviews</i> , 2003, 195, 117-135.	2.8	234
87	Concerted action of the chemokine and lymphotoxin system in secondary lymphoid-organ development. <i>Current Opinion in Immunology</i> , 2003, 15, 217-224.	2.4	59
88	Positioning of follicular dendritic cells within the spleen controls prion neuroinvasion. <i>Nature</i> , 2003, 425, 957-962.	13.7	195
89	Role of Homeostatic Chemokine and Sphingosine-1-Phosphate Receptors in the Organization of Lymphoid Tissue. <i>Annals of the New York Academy of Sciences</i> , 2003, 987, 107-116.	1.8	19
90	Regulation of Dendritic Cell Migration to the Draining Lymph Node. <i>Journal of Experimental Medicine</i> , 2003, 198, 615-621.	4.2	806

#	ARTICLE	IF	CITATIONS
91	Cooperating Mechanisms of CXCR5 and CCR7 in Development and Organization of Secondary Lymphoid Organs. <i>Journal of Experimental Medicine</i> , 2003, 197, 1199-1204.	4.2	167
92	Differentiation of Effector/Memory $\gamma\delta$ T Cells and Migratory Routes in Lymph Nodes or Inflammatory Sites. <i>Journal of Experimental Medicine</i> , 2003, 198, 391-397.	4.2	300
93	Shaping Up Adaptive Immunity: The Impact of CCR7 and CXCR5 on Lymphocyte Trafficking. <i>Microcirculation</i> , 2003, 10, 325-334.	1.0	57
94	Expression of B-cell-attracting chemokine 1 (CXCL13) by malignant lymphocytes and vascular endothelium in primary central nervous system lymphoma. <i>Blood</i> , 2003, 101, 815-821.	0.6	182
95	Hochmaligne B-Zell-Lymphome. , 2003, , 456-478.		0
96	Surface Expression and Endocytosis of the Human Cytomegalovirus-encoded Chemokine Receptor US28 Is Regulated by Agonist-independent Phosphorylation. <i>Journal of Biological Chemistry</i> , 2002, 277, 45122-45128.	1.6	64
97	Roles of SLC/CCL21 and CCR7 in Human Kidney for Mesangial Proliferation, Migration, Apoptosis, and Tissue Homeostasis. <i>Journal of Immunology</i> , 2002, 168, 4301-4307.	0.4	83
98	Secondary Lymphoid Tissue Chemokine (CCL21) Activates CXCR3 to Trigger a Ca^{2+} Current and Chemotaxis in Murine Microglia. <i>Journal of Immunology</i> , 2002, 168, 3221-3226.	0.4	138
99	Monocyte-Expressed Urokinase Regulates Human Vascular Smooth Muscle Cell Migration in a Coculture Model. <i>Biological Chemistry</i> , 2002, 383, 217-21.	1.2	24
100	Regulation of E2F1-Dependent Gene Transcription and Apoptosis by the ETS-Related Transcription Factor GABP β 1. <i>Molecular and Cellular Biology</i> , 2002, 22, 2147-2158.	1.1	25
101	Chemokine Requirements for B Cell Entry to Lymph Nodes and Peyer's Patches. <i>Journal of Experimental Medicine</i> , 2002, 196, 65-75.	4.2	479
102	Up-regulation of the chemokine receptor CCR7 in classical but not in lymphocyte-predominant Hodgkin disease correlates with distinct dissemination of neoplastic cells in lymphoid organs. <i>Blood</i> , 2002, 99, 1109-1116.	0.6	98
103	Role for CCR7 Ligands in the Emigration of Newly Generated T Lymphocytes from the Neonatal Thymus. <i>Immunity</i> , 2002, 16, 205-218.	6.6	216
104	Balanced responsiveness to chemoattractants from adjacent zones determines B-cell position. <i>Nature</i> , 2002, 416, 94-99.	13.7	506
105	Systemic immunoregulatory and pathogenic functions of homeostatic chemokine receptors. <i>Journal of Leukocyte Biology</i> , 2002, 72, 1-8.	1.5	60
106	CC Chemokine Receptor 7-dependent and -independent Pathways for Lymphocyte Homing. <i>Journal of Experimental Medicine</i> , 2001, 194, 1875-1881.	4.2	121
107	The T cell chemokine receptor CCR7 is internalized on stimulation with ELC, but not with SLC. <i>European Journal of Immunology</i> , 2001, 31, 3291-3297.	1.6	118
108	The Poliovirus Receptor CD155 Mediates Cell-to-Matrix Contacts by Specifically Binding to Vitronectin. <i>Virology</i> , 2001, 285, 218-227.	1.1	84

#	ARTICLE	IF	CITATIONS
109	Involvement of inhibitory NKRs in the survival of a subset of memory-phenotype CD8+ T cells. <i>Nature Immunology</i> , 2001, 2, 430-435.	7.0	153
110	Skewed maturation of memory HIV-specific CD8 T lymphocytes. <i>Nature</i> , 2001, 410, 106-111.	13.7	910
111	Aberrant High Expression of B Lymphocyte Chemokine (Blc/Cxcl13) by C11b+Cd11c+ Dendritic Cells in Murine Lupus and Preferential Chemotaxis of B1 Cells towards Blc. <i>Journal of Experimental Medicine</i> , 2001, 193, 1393-1402.	4.2	149
112	Tumorigenesis induced by the HHV8-encoded chemokine receptor requires ligand modulation of high constitutive activity. <i>Journal of Clinical Investigation</i> , 2001, 108, 1789-1796.	3.9	95
113	Phosphorylation of Oct-2 at sites located in the POU domain induces differential down-regulation of Oct-2 DNA-binding ability. <i>Biochemical Journal</i> , 2000, 347, 29.	1.7	7
114	Phosphorylation of Oct-2 at sites located in the POU domain induces differential down-regulation of Oct-2 DNA-binding ability. <i>Biochemical Journal</i> , 2000, 347, 29-35.	1.7	18
115	A chemokine-driven positive feedback loop organizes lymphoid follicles. <i>Nature</i> , 2000, 406, 309-314.	13.7	1,103
116	Sphingosine-1-phosphate is a ligand for the G protein-coupled receptor EDG-6. <i>Blood</i> , 2000, 95, 2624-2629.	0.6	176
117	A Shift in the Phenotype of Melan-A-Specific CTL Identifies Melanoma Patients with an Active Tumor-Specific Immune Response. <i>Journal of Immunology</i> , 2000, 165, 6644-6652.	0.4	128
118	Cxc Chemokine Receptor 5 Expression Defines Follicular Homing T Cells with B Cell Helper Function. <i>Journal of Experimental Medicine</i> , 2000, 192, 1553-1562.	4.2	1,094
119	Follicular B Helper T Cells Express Cxc Chemokine Receptor 5, Localize to B Cell Follicles, and Support Immunoglobulin Production. <i>Journal of Experimental Medicine</i> , 2000, 192, 1545-1552.	4.2	1,284
120	Urokinase Stimulates Human Vascular Smooth Muscle Cell Migration via a Phosphatidylinositol 3-Kinase-Tyk2 Interaction. <i>Journal of Biological Chemistry</i> , 2000, 275, 39466-39473.	1.6	74
121	Identification of a Nuclear Respiratory Factor-1 Binding Site within the Core Promoter of the human polio virus receptor/CD155 Gene. <i>Journal of Biological Chemistry</i> , 2000, 275, 12453-12462.	1.6	36
122	CCR6 Mediates Dendritic Cell Localization, Lymphocyte Homeostasis, and Immune Responses in Mucosal Tissue. <i>Immunity</i> , 2000, 12, 495-503.	6.6	478
123	CXCR5-deficient mice develop functional germinal centers in the splenic T cell zone. <i>European Journal of Immunology</i> , 2000, 30, 560-567.	1.6	91
124	Sphingosine-1-phosphate is a ligand for the G protein-coupled receptor EDG-6. <i>Blood</i> , 2000, 95, 2624-2629.	0.6	0
125	Identification and Characterization of the cis-Acting Elements of the Human CD155 Gene Core Promoter. <i>Journal of Biological Chemistry</i> , 1999, 274, 1791-1800.	1.6	21
126	Compromised Ox40 Function in Cd28-Deficient Mice Is Linked with Failure to Develop Cxc Chemokine Receptor 5 ⁺ Positive Cd4 Cells and Germinal Centers. <i>Journal of Experimental Medicine</i> , 1999, 190, 1115-1122.	4.2	247

#	ARTICLE	IF	CITATIONS
127	Two subsets of memory T lymphocytes with distinct homing potentials and effector functions. <i>Nature</i> , 1999, 401, 708-712.	13.7	5,333
128	Distinct patterns and kinetics of chemokine production regulate dendritic cell function. <i>European Journal of Immunology</i> , 1999, 29, 1617-1625.	1.6	588
129	Switch in chemokine receptor expression upon TCR stimulation reveals novel homing potential for recently activated T cells. <i>European Journal of Immunology</i> , 1999, 29, 2037-2045.	1.6	348
130	CCR7 Coordinates the Primary Immune Response by Establishing Functional Microenvironments in Secondary Lymphoid Organs. <i>Cell</i> , 1999, 99, 23-33.	13.5	2,122
131	The murine chemokine receptor CXCR4 is tightly regulated during T cell development and activation. <i>Journal of Leukocyte Biology</i> , 1999, 66, 996-1004.	1.5	46
132	A Lymphoid Tissue-Specific Receptor, EDG6, with Potential Immune Modulatory Functions Mediated by Extracellular Lysophospholipids. <i>Current Topics in Microbiology and Immunology</i> , 1999, 246, 131-137.	0.7	21
133	EDG6, a Novel G-Protein-Coupled Receptor Related to Receptors for Bioactive Lysophospholipids, Is Specifically Expressed in Lymphoid Tissue. <i>Genomics</i> , 1998, 53, 164-169.	1.3	222
134	Downstream Activation of a TATA-less Promoter by Oct-2, Bob1, and NF- κ B Directs Expression of the Homing Receptor BLR1 to Mature B Cells. <i>Journal of Biological Chemistry</i> , 1998, 273, 28831-28836.	1.6	63
135	The Promoters for Human and Monkey Poliovirus Receptors. <i>Journal of Biological Chemistry</i> , 1997, 272, 5579-5586.	1.6	35
136	Abnormal Expression of the B-Cell Homing Chemokine Receptor BLR1 During the Progression of Acquired Immunodeficiency Syndrome. <i>Blood</i> , 1997, 90, 520-525.	0.6	42
137	Analyzing cytotoxic T lymphocyte activity: a simple and reliable flow cytometry-based assay. <i>Journal of Immunological Methods</i> , 1997, 204, 135-142.	0.6	57
138	Abnormal Expression of the B-Cell Homing Chemokine Receptor BLR1 During the Progression of Acquired Immunodeficiency Syndrome. <i>Blood</i> , 1997, 90, 520-525.	0.6	7
139	A Putative Chemokine Receptor, BLR1, Directs B Cell Migration to Defined Lymphoid Organs and Specific Anatomic Compartments of the Spleen. <i>Cell</i> , 1996, 87, 1037-1047.	13.5	1,059
140	A κ v murine leukemia virus enhances lymphomagenesis in myc-kappa transgenic and in wild-type mice. <i>Virology</i> , 1995, 206, 93-99.	1.1	9
141	A versatile flow cytometry-based assay for the determination of short- and long-term natural killer cell activity. <i>Journal of Immunological Methods</i> , 1995, 185, 209-216.	0.6	68
142	Breakpoints of burkitt's lymphoma t(8;22) translocations map within a distance of 300 kb downstream of MYC. <i>Genes Chromosomes and Cancer</i> , 1994, 9, 282-287.	1.5	80
143	The hepatitis B virusPreS2/St transactivator utilizes AP-1 and other transcription factors for transactivation. <i>Hepatology</i> , 1994, 19, 23-31.	3.6	33
144	Cloning and chromosomal organization of a gene encoding a putative amino-acid permease from <i>Saccharomyces cerevisiae</i> . <i>Gene</i> , 1994, 143, 129-133.	1.0	24

#	ARTICLE	IF	CITATIONS
145	The G protein-coupled receptor BLR1 is involved in murine B cell differentiation and is also expressed in neuronal tissues. <i>European Journal of Immunology</i> , 1993, 23, 2532-2539.	1.6	65
146	Identification of a protein from <i>Saccharomyces cerevisiae</i> with E2F-like DNA-binding and transactivating properties. <i>FEBS Letters</i> , 1993, 321, 153-158.	1.3	7
147	Evaluation of the Utility of Interphase Cytogenetics to Detect Residual Cells with a Malignant Genotype in Mixed Cell Populations: A Burkitt Lymphoma Model. <i>DNA and Cell Biology</i> , 1993, 12, 637-643.	0.9	9
148	Specific metaphase and interphase detection of the breakpoint region in 8q24 of burkitt lymphoma cells by triple-color fluorescence in situ hybridization. <i>Genes Chromosomes and Cancer</i> , 1992, 4, 69-74.	1.5	87
149	Differentiation-specific expression of a novel G protein-coupled receptor from Burkitt's lymphoma. <i>European Journal of Immunology</i> , 1992, 22, 2795-2799.	1.6	137
150	A novel divergently transcribed human histone H2A/H2B gene pair. <i>DNA Sequence</i> , 1991, 1, 409-413.	0.7	16
151	t(2;8) variant translocation in burkitt's lymphoma: Mapping of chromosomal breakpoints by in situ hybridization. <i>International Journal of Cancer</i> , 1989, 44, 261-265.	2.3	4