

Mercedes Barbero

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2438225/publications.pdf>

Version: 2024-02-01

8
papers

213
citations

1478505

6
h-index

1720034

7
g-index

8
all docs

8
docs citations

8
times ranked

294
citing authors

| # | ARTICLE | IF | CITATIONS |
|---|--|-----|-----------|
| 1 | Measurement of arsenic compounds in littoral zone algae from the Western Mediterranean Sea. Occurrence of arsenobetaine. <i>Chemosphere</i> , 2010, 81, 867-875. | 8.2 | 52 |
| 2 | Determination of Water-Soluble Arsenic Compounds in Commercial Edible Seaweed by LC-ICPMS. <i>Journal of Agricultural and Food Chemistry</i> , 2011, 59, 12963-12968. | 5.2 | 50 |
| 3 | Occurrence of arsenic species in algae and freshwater plants of an extreme arid region in northern Chile, the Loa River Basin. <i>Chemosphere</i> , 2013, 90, 556-564. | 8.2 | 41 |
| 4 | Occurrence of inorganic arsenic in edible Shiitake (<i>Lentinula edodes</i>) products. <i>Food Chemistry</i> , 2014, 158, 207-215. | 8.2 | 41 |
| 5 | Notes on Some Corticolous and Lignicolous <i>Buellia</i> Species From the Iberian Peninsula. <i>Lichenologist</i> , 2000, 32, 105-128. | 0.8 | 22 |
| 6 | <i>Rinodina Algarvensis</i> , A New Saxicolous Sorediate Species from Portugal Containing the Stictic Acid Complex. <i>Lichenologist</i> , 1996, 28, 1-8. | 0.8 | 6 |
| 7 | <i>Neofuscelia halei</i> sp. nov. (Lichen-Forming Ascomycota) from Spain and the Canary Islands. <i>Bryologist</i> , 1993, 96, 355. | 0.6 | 1 |
| 8 | <i>Rinodina Algarvensis</i> , A New Saxicolous Sorediate Species from Portugal Containing the Stictic Acid Complex. <i>Lichenologist</i> , 1996, 28, 1. | 0.8 | 0 |