

# Ming Zhong

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2443330/publications.pdf>

Version: 2024-02-01

24  
papers

503  
citations

516710

16  
h-index

677142

22  
g-index

24  
all docs

24  
docs citations

24  
times ranked

634  
citing authors

#	ARTICLE	IF	CITATIONS
1	Preoperative and postoperative predictors of long-term outcome after endovascular treatment of poor-grade aneurysmal subarachnoid hemorrhage. <i>Journal of Neurosurgery</i> , 2016, 126, 1764-1771.	1.6	43
2	Sinomenine enhances microglia M2 polarization and attenuates inflammatory injury in intracerebral hemorrhage. <i>Journal of Neuroimmunology</i> , 2016, 299, 28-34.	2.3	40
3	Aneurysm rebleeding after poor-grade aneurysmal subarachnoid hemorrhage: Predictors and impact on clinical outcomes. <i>Journal of the Neurological Sciences</i> , 2016, 371, 62-66.	0.6	39
4	Stent-assisted coiling versus coiling alone of poor-grade ruptured intracranial aneurysms: a multicenter study. <i>Journal of NeuroInterventional Surgery</i> , 2017, 9, 165-168.	3.3	36
5	Predicting Long-Term Outcomes After Poor-Grade Aneurysmal Subarachnoid Hemorrhage Using Decision Tree Modeling. <i>Neurosurgery</i> , 2020, 87, 523-529.	1.1	32
6	Factors and outcomes associated with ultra-early surgery for poor-grade aneurysmal subarachnoid haemorrhage: a multicentre retrospective analysis. <i>BMJ Open</i> , 2015, 5, e007410-e007410.	1.9	31
7	Larger size ratio associated with the rupture of very small (≤3mm) anterior communicating artery aneurysms. <i>Journal of NeuroInterventional Surgery</i> , 2017, 9, 278-282.	3.3	30
8	Endovascular Coiling versus Surgical Clipping for Poor-Grade Ruptured Intracranial Aneurysms: Postoperative Complications and Clinical Outcome in a Multicenter Poor-Grade Aneurysm Study. <i>American Journal of Neuroradiology</i> , 2016, 37, 873-878.	2.4	28
9	Predicting intraprocedural rupture and thrombus formation during coiling of ruptured anterior communicating artery aneurysms. <i>Journal of NeuroInterventional Surgery</i> , 2017, 9, 370-375.	3.3	27
10	Stent-assisted coiling versus coiling alone of ruptured anterior communicating artery aneurysms: A single-center experience. <i>Clinical Neurology and Neurosurgery</i> , 2016, 144, 96-100.	1.4	21
11	Variation in Patient Characteristics and Outcomes Between Early and Delayed Surgery in Poor-Grade Aneurysmal Subarachnoid Hemorrhage. <i>Neurosurgery</i> , 2016, 78, 224-231.	1.1	20
12	Primary decompressive craniectomy for poor-grade middle cerebral artery aneurysms with associated intracerebral hemorrhage. <i>Clinical Neurology and Neurosurgery</i> , 2015, 133, 1-5.	1.4	19
13	Complications and outcomes after early surgical treatment for poor-grade ruptured intracranial aneurysms: A multicenter retrospective cohort. <i>International Journal of Surgery</i> , 2015, 23, 57-61.	2.7	19
14	Smoking Associated with Increased Aneurysm Size in Patients with Anterior Communicating Artery Aneurysms. <i>World Neurosurgery</i> , 2016, 87, 155-161.	1.3	18
15	A Multicenter prospective study of poor-grade aneurysmal subarachnoid hemorrhage (AMPAS): observational registry study. <i>BMC Neurology</i> , 2014, 14, 86.	1.8	17
16	Endovascular Coiling versus Surgical Clipping of Very Small Ruptured Anterior Communicating Artery Aneurysms. <i>World Neurosurgery</i> , 2019, 126, e1246-e1250.	1.3	17
17	Sex differences in aneurysm morphologies and clinical outcomes in ruptured anterior communicating artery aneurysms: a retrospective study. <i>BMJ Open</i> , 2016, 6, e009920.	1.9	16
18	Expression change of interleukin-8 gene in rabbit basilar artery after subarachnoid hemorrhage. <i>Neuroscience Bulletin</i> , 2007, 23, 151-155.	2.9	15

#	ARTICLE	IF	CITATIONS
19	Predictors of good functional outcomes and mortality in patients with severe rebleeding after aneurysmal subarachnoid hemorrhage. <i>Clinical Neurology and Neurosurgery</i> , 2016, 144, 28-32.	1.4	10
20	Comparison of Aggressive Surgical Treatment and Palliative Treatment in Elderly Patients with Poor-Grade Intracranial Aneurysmal Subarachnoid Hemorrhage. <i>BioMed Research International</i> , 2018, 2018, 1-8.	1.9	10
21	A Multicenter Analysis of Computed Tomography Angiography Alone Versus Digital Subtraction Angiography for the Surgical Treatment of Poor-Grade Aneurysmal Subarachnoid Hemorrhage. <i>World Neurosurgery</i> , 2016, 91, 106-111.	1.3	5
22	Change in Urine Albumin-to-Creatinine Ratio and Risk of Diabetic Peripheral Neuropathy in Type 2 Diabetes: A Retrospective Cohort Study. <i>Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy</i> , 2021, Volume 14, 1763-1772.	2.4	4
23	China Intracranial Aneurysm Project (CIAP): protocol for a prospective cohort study of interventional treatment and craniotomy for unruptured aneurysms. <i>BMJ Open</i> , 2018, 8, e019333.	1.9	3
24	Predictors of unfavorable outcome in stent-assisted coiling for symptomatic unruptured intracranial spontaneous vertebral artery dissecting aneurysms (uis-VADAs): results from a multicenter study. <i>Journal of NeuroInterventional Surgery</i> , 2022, 14, 1008-1013.	3.3	3