

Ahmed A Attallah

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/244357/publications.pdf>

Version: 2024-02-01

16
papers

223
citations

1039880

9
h-index

996849

15
g-index

16
all docs

16
docs citations

16
times ranked

302
citing authors

#	ARTICLE	IF	CITATIONS
1	Krüppel-like factor 6-mediated loss of BCAA catabolism contributes to kidney injury in mice and humans. Proceedings of the National Academy of Sciences of the United States of America, 2021, 118, .	3.3	34
2	Loss of proximal tubular transcription factor Krüppel-like factor 15 exacerbates kidney injury through loss of fatty acid oxidation. Kidney International, 2021, 100, 1250-1267.	2.6	28
3	Fibro-1± score as a simple and useful non-invasive test for predicting significant liver fibrosis in chronic hepatitis C patients. Arab Journal of Gastroenterology, 2011, 12, 74-79.	0.4	21
4	A simple diagnostic index comprising epithelial membrane antigen and fibronectin for hepatocellular carcinoma. Annals of Hepatology, 2015, 14, 869-880.	0.6	21
5	Fibro-check: a combination of direct and indirect markers for liver fibrosis staging in chronic hepatitis C patients. Annals of Hepatology, 2015, 14, 225-233.	0.6	20
6	Interferon-gamma is associated with hepatic dysfunction in fibrosis, cirrhosis, and hepatocellular carcinoma. Journal of Immunoassay and Immunochemistry, 2016, 37, 597-610.	0.5	20
7	GPC-HCC model: a combination of glypican-3 with other routine parameters improves the diagnostic efficacy in hepatocellular carcinoma. Tumor Biology, 2016, 37, 12571-12577.	0.8	19
8	Circulating levels and clinical implications of epithelial membrane antigen and cytokeratin-1 in women with breast cancer: can their ratio improve the results?. Tumor Biology, 2014, 35, 10737-10745.	0.8	14
9	Simplified HCC-ART score for highly sensitive detection of small-sized and early-stage hepatocellular carcinoma in the widely used Okuda, CLIP, and BCLC staging systems. International Journal of Clinical Oncology, 2017, 22, 332-339.	1.0	10
10	HCC-DETECT: a combination of nuclear, cytoplasmic, and oncofetal proteins as biomarkers for hepatocellular carcinoma. Tumor Biology, 2015, 36, 7667-7674.	0.8	8
11	Circulating levels of collagen III and MMP-1 in patients with chronic hepatitis C co-infected with hepatitis B virus. British Journal of Biomedical Science, 2017, 74, 95-100.	1.2	7
12	Fibro-check: a combination of direct and indirect markers for liver fibrosis staging in chronic hepatitis C patients. Annals of Hepatology, 2015, 14, 225-33.	0.6	7
13	Impact of Hepatitis C Virus/Schistosoma mansoni Coinfection on the Circulating Levels of HCV-NS4 Protein and Extracellular-Matrix Deposition in Patients with Different Hepatic Fibrosis Stages. American Journal of Tropical Medicine and Hygiene, 2016, 95, 1044-1050.	0.6	5
14	Combined use of Epithelial Membrane Antigen and Nuclear Matrix Protein 52 as Sensitive Biomarkers for Detection of Bladder Cancer. International Journal of Biological Markers, 2015, 30, 407-413.	0.7	4
15	Perinatal transmission of hepatitis C antigens: envelope 1, envelope 2 and non-structural 4. Infectious Diseases, 2015, 47, 568-574.	1.4	4
16	Levels of Schistosoma mansoni Circulating Antigen in Chronic Hepatitis C Patients with Different Stages of Liver Fibrosis. Journal of Immunoassay and Immunochemistry, 2016, 37, 316-330.	0.5	1