
## Marek Dariusz Kasielski

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/2446547/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Body mass index is negatively associated with telomere length: a collaborative cross-sectional meta-analysis of 87 observational studies. American Journal of Clinical Nutrition, 2018, 108, 453-475.	2.2	137
2	Anti-MCV and anti-CCP antibodies—diagnostic and prognostic value in children with juvenile idiopathic arthritis (JIA). Clinical Rheumatology, 2016, 35, 2699-2706.	1.0	13
3	The relationship between peripheral blood mononuclear cells telomere length and diet - unexpected effect of red meat. Nutrition Journal, 2015, 15, 68.	1.5	22
4	Reactive oxygen species and serum antioxidant defense in juvenile idiopathic arthritis. Clinical Rheumatology, 2015, 34, 451-456.	1.0	19
5	Myocardial oxidative stress in patients with active infective endocarditis. International Journal of Cardiology, 2013, 167, 270-276.	0.8	21
6	Uric Acid but Not Apple Polyphenols Is Responsible for the Rise of Plasma Antioxidant Activity after Apple Juice Consumption in Healthy Subjects. Journal of the American College of Nutrition, 2010, 29, 397-406.	1.1	44
7	No evidence of enhanced oxidant production in blood obtained from patients with obstructive sleep apnea. Journal of Negative Results in BioMedicine, 2008, 7, 10.	1.4	10
8	Decreased H2O2 in exhaled breath condensate during pregnancy—Feasible effect of 17β-estradiol. Respiratory Physiology and Neurobiology, 2008, 162, 152-159.	0.7	6
9	Elevated resting and agonist-induced whole blood chemiluminescence in patients with active infective endocarditis. Interactive Cardiovascular and Thoracic Surgery, 2008, 8, 12-16.	0.5	8
10	Increased hydrogen peroxide concentration in the exhaled breath condensate of stable COPD patients after nebulized N-acetylcysteine. Pulmonary Pharmacology and Therapeutics, 2007, 20, 281-289.	1.1	24
11	Increased Exhaled H2O2 and Impaired Lung Function in Patients Undergoing Bioincompatible Hemodialysis. International Journal of Artificial Organs, 2007, 30, 879-888.	0.7	10
12	Increased whole blood chemiluminescence in patients with chronic renal failure independent of hemodialysis treatment. Archivum Immunologiae Et Therapiae Experimentalis, 2006, 54, 347-355.	1.0	12
13	Increased hydrogen peroxide in the exhaled breath of uraemic patients unaffected by haemodialysis. Nephrology Dialysis Transplantation, 2004, 19, 158-163.	0.4	20
14	Effect of inhaled N-acetylcysteine on hydrogen peroxide exhalation in healthy subjects. Pulmonary Pharmacology and Therapeutics, 2004, 17, 155-162.	1.1	30
15	Elevated exhalation of hydrogen peroxide and thiobarbituric acid reactive substances in patients with community acquired pneumonia. Respiratory Medicine, 2004, 98, 669-676.	1.3	47
16	Elevated exhalation of hydrogen peroxide in patients with systemic sclerosis. European Journal of Clinical Investigation, 2003, 33, 274-279.	1.7	26
17	Exhaled hydrogen peroxide correlates with the release of reactive oxygen species by blood phagocytes in healthy subjects. Respiratory Medicine, 2003, 97, 718-725.	1.3	40
18	Long-term administration of N-acetylcysteine decreases hydrogen peroxide exhalation in subjects with chronic obstructive pulmonary disease. Respiratory Medicine, 2001, 95, 448-456.	1.3	159

#	Article	IF	CITATIONS
19	Inhaled glucocorticosteroids decrease hydrogen peroxide level in expired air condensate in asthmatic patients. Respiratory Medicine, 2000, 94, 416-421.	1.3	62
20	Increased content of thiobarbituric acid-reactive substances and hydrogen peroxide in the expired breath condensate of patients with stable chronic obstructive pulmonary disease: no significant effect of cigarette smoking. Respiratory Medicine, 1999, 93, 389-396.	1.3	161