

# Teresa Victor

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2448121/publications.pdf>

Version: 2024-02-01

24  
papers

2,117  
citations

516710

16  
h-index

642732

23  
g-index

24  
all docs

24  
docs citations

24  
times ranked

3439  
citing authors

#	ARTICLE	IF	CITATIONS
1	Functional magnetic resonance imaging data for the association between polygenic risk scores for neuroticism and reward-punishment processing. <i>Data in Brief</i> , 2022, 42, 108014.	1.0	0
2	Polygenic risk for neuroticism moderates response to gains and losses in amygdala and caudate: Evidence from a clinical cohort. <i>Journal of Affective Disorders</i> , 2021, 293, 124-132.	4.1	5
3	Evaluating the resource allocation index as a potential fMRI-based biomarker for substance use disorder. <i>Drug and Alcohol Dependence</i> , 2020, 216, 108211.	3.2	1
4	Women with Major Depressive Disorder, Irrespective of Comorbid Anxiety Disorders, Show Blunted Bilateral Frontal Responses during Win and Loss Anticipation. <i>Journal of Affective Disorders</i> , 2020, 273, 157-166.	4.1	4
5	Web-Based Graphic Representation of the Life Course of Mental Health: Cross-Sectional Study Across the Spectrum of Mood, Anxiety, Eating, and Substance Use Disorders. <i>JMIR Mental Health</i> , 2020, 7, e16919.	3.3	9
6	EEG Microstates Temporal Dynamics Differentiate Individuals with Mood and Anxiety Disorders From Healthy Subjects. <i>Frontiers in Human Neuroscience</i> , 2019, 13, 56.	2.0	54
7	Machine Learning Analysis of the Relationships Between Gray Matter Volume and Childhood Trauma in a Transdiagnostic Community-Based Sample. <i>Biological Psychiatry: Cognitive Neuroscience and Neuroimaging</i> , 2019, 4, 734-742.	1.5	11
8	Interoception and Mental Health: A Roadmap. <i>Biological Psychiatry: Cognitive Neuroscience and Neuroimaging</i> , 2018, 3, 501-513.	1.5	524
9	A Nonlinear Simulation Framework Supports Adjusting for Age When Analyzing BrainAGE. <i>Frontiers in Aging Neuroscience</i> , 2018, 10, 317.	3.4	183
10	Tulsa 1000: a naturalistic study protocol for multilevel assessment and outcome prediction in a large psychiatric sample. <i>BMJ Open</i> , 2018, 8, e016620.	1.9	88
11	Comparison of two different analysis approaches for DTI free-water corrected and uncorrected maps in the study of white matter microstructural integrity in individuals with depression. <i>Human Brain Mapping</i> , 2017, 38, 4690-4702.	3.6	30
12	Co-altered functional networks and brain structure in unmedicated patients with bipolar and major depressive disorders. <i>Brain Structure and Function</i> , 2017, 222, 4051-4064.	2.3	77
13	Real-Time Functional Magnetic Resonance Imaging Amygdala Neurofeedback Changes Positive Information Processing in Major Depressive Disorder. <i>Biological Psychiatry</i> , 2017, 82, 578-586.	1.3	92
14	Sex differences in neural responses to subliminal sad and happy faces in healthy individuals: Implications for depression. <i>Journal of Neuroscience Research</i> , 2017, 95, 703-710.	2.9	13
15	Aberrant decision-making and drug addiction – how strong is the evidence?. <i>Current Opinion in Behavioral Sciences</i> , 2017, 13, 25-33.	3.9	34
16	The Effect of Mineralocorticoid and Glucocorticoid Receptor Antagonism on Autobiographical Memory Recall and Amygdala Response to Implicit Emotional Stimuli. <i>International Journal of Neuropsychopharmacology</i> , 2016, 19, pyw036.	2.1	9
17	Relationship between neurotoxic kynurenine metabolites and reductions in right medial prefrontal cortical thickness in major depressive disorder. <i>Brain, Behavior, and Immunity</i> , 2016, 53, 39-48.	4.1	136
18	Resting-state functional network connectivity in prefrontal regions differs between unmedicated patients with bipolar and major depressive disorders. <i>Journal of Affective Disorders</i> , 2016, 190, 483-493.	4.1	102

#	ARTICLE	IF	CITATIONS
19	Activation of the kynurenine pathway is associated with striatal volume in major depressive disorder. <i>Psychoneuroendocrinology</i> , 2015, 62, 54-58.	2.7	80
20	Reduction of kynurenic acid to quinolinic acid ratio in both the depressed and remitted phases of major depressive disorder. <i>Brain, Behavior, and Immunity</i> , 2015, 46, 55-59.	4.1	162
21	Neuroprotective kynurenine metabolite indices are abnormally reduced and positively associated with hippocampal and amygdalar volume in bipolar disorder. <i>Psychoneuroendocrinology</i> , 2015, 52, 200-211.	2.7	106
22	Putative Neuroprotective and Neurotoxic Kynurenine Pathway Metabolites Are Associated with Hippocampal and Amygdalar Volumes in Subjects with Major Depressive Disorder. <i>Neuropsychopharmacology</i> , 2015, 40, 463-471.	5.4	199
23	Inflammation and neurological disease-related genes are differentially expressed in depressed patients with mood disorders and correlate with morphometric and functional imaging abnormalities. <i>Brain, Behavior, and Immunity</i> , 2013, 31, 161-171.	4.1	127
24	Changes in the neural correlates of implicit emotional face processing during antidepressant treatment in major depressive disorder. <i>International Journal of Neuropsychopharmacology</i> , 2013, 16, 2195-2208.	2.1	71