

Christoph Riethmüller

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2450402/publications.pdf>

Version: 2024-02-01

9
papers

253
citations

1307594

7
h-index

1474206

9
g-index

9
all docs

9
docs citations

9
times ranked

363
citing authors

#	ARTICLE	IF	CITATIONS
1	Corneocyte Nanotexture as Biomarker for Individual Susceptibility to Skin Irritants. <i>Annals of Work Exposures and Health</i> , 2021, 65, 201-205.	1.4	3
2	Changes in Skin Barrier Function after Repeated Exposition to Phospholipid-Based Surfactants and Sodium Dodecyl Sulfate In Vivo and Corneocyte Surface Analysis by Atomic Force Microscopy. <i>Pharmaceutics</i> , 2021, 13, 436.	4.5	13
3	Filaggrin Expression and Processing Deficiencies Impair Corneocyte Surface Texture and Stiffness in Mice. <i>Journal of Investigative Dermatology</i> , 2020, 140, 615-623.e5.	0.7	28
4	Actinic keratosis and surrounding skin exhibit changes in corneocyte surface topography and decreased levels of filaggrin degradation products. <i>Experimental Dermatology</i> , 2020, 29, 462-466.	2.9	2
5	Specific barrier response profiles after experimentally induced skin irritation in vivo. <i>Contact Dermatitis</i> , 2018, 79, 59-66.	1.4	29
6	Assessing the skin barrier via corneocyte morphometry. <i>Experimental Dermatology</i> , 2018, 27, 923-930.	2.9	20
7	Effect of allergens and irritants on levels of natural moisturizing factor and corneocyte morphology. <i>Contact Dermatitis</i> , 2017, 76, 287-295.	1.4	27
8	Filaggrin breakdown products determine corneocyte conformation in patients with atopic dermatitis. <i>Journal of Allergy and Clinical Immunology</i> , 2015, 136, 1573-1580.e2.	2.9	93
9	Nanotopography follows force in TGF- β 1 stimulated epithelium. <i>Nanotechnology</i> , 2010, 21, 265102.	2.6	38