

Yung H Wong

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/2453691/yung-h-wong-publications-by-year.pdf>

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

74
papers

1,865
citations

22
h-index

40
g-index

75
ext. papers

2,036
ext. citations

5.4
avg, IF

4.7
L-index

#	Paper	IF	Citations
74	Modeling the Heterodimer Interfaces of Melatonin Receptors. <i>Frontiers in Cellular Neuroscience</i> , 2021 , 15, 725296	6.1	
73	Re-examining the Dissociation Model of G protein activation from the perspective of G $\beta\gamma$ signaling. <i>FEBS Journal</i> , 2021 , 288, 2490-2501	5.7	3
72	CTCF and EGR1 suppress breast cancer cell migration through transcriptional control of Nm23-H1. <i>Scientific Reports</i> , 2021 , 11, 491	4.9	6
71	Molecular basis defining the selectivity of substituted isoquinolinones for the melatonin MT receptor. <i>Biochemical Pharmacology</i> , 2020 , 177, 114020	6	3
70	Age Associated Decrease of MT-1 Melatonin Receptor in Human Dermal Skin Fibroblasts Impairs Protection Against UV-Induced DNA Damage. <i>International Journal of Molecular Sciences</i> , 2020 , 21,	6.3	15
69	Mutations at the dimer interface and surface residues of Nm23-H1 metastasis suppressor affect its expression and function. <i>Molecular and Cellular Biochemistry</i> , 2020 , 474, 95-112	4.2	
68	GPCRs in Autocrine and Paracrine Regulations. <i>Frontiers in Endocrinology</i> , 2019 , 10, 428	5.7	10
67	Small Molecules as Drugs to Upregulate Metastasis Suppressors in Cancer Cells. <i>Current Medicinal Chemistry</i> , 2019 , 26, 5876-5899	4.3	6
66	Role of G Protein-Coupled Receptors in the Regulation of Structural Plasticity and Cognitive Function. <i>Molecules</i> , 2017 , 22,	4.8	23
65	RGS19 upregulates Nm23-H1/2 metastasis suppressors by transcriptional activation via the cAMP/PKA/CREB pathway. <i>Oncotarget</i> , 2017 , 8, 69945-69960	3.3	10
64	Promoting axon regeneration in the adult CNS by modulation of the melanopsin/GPCR signaling. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2016 , 113, 1937-42	11.5	68
63	Differential Regulation of CXCL8 Production by Different G Protein Subunits with Synergistic Stimulation by Gi- and Gq-Regulated Pathways. <i>Journal of Molecular Biology</i> , 2016 , 428, 3869-84	6.5	4
62	Regulator of G protein signaling 20 enhances cancer cell aggregation, migration, invasion and adhesion. <i>Cellular Signalling</i> , 2016 , 28, 1663-72	4.9	25
61	Metastasis suppressors Nm23H1 and Nm23H2 differentially regulate neoplastic transformation and tumorigenesis. <i>Cancer Letters</i> , 2015 , 361, 207-17	9.9	16
60	Injured adult retinal axons with Pten and Socs3 co-deletion reform active synapses with suprachiasmatic neurons. <i>Neurobiology of Disease</i> , 2015 , 73, 366-76	7.5	35
59	Activator of G protein signaling 3 forms a complex with resistance to inhibitors of cholinesterase-8A without promoting nucleotide exchange on G $\beta\gamma$ (13). <i>Molecular and Cellular Biochemistry</i> , 2015 , 401, 27-38	4.2	0
58	Regulatory functions of Nm23-H2 in tumorigenesis: insights from biochemical to clinical perspectives. <i>Naunyn-Schmiedeberg's Archives of Pharmacology</i> , 2015 , 388, 243-56	3.4	9

57	An intact helical domain is required for G α 14 to stimulate phospholipase C β <i>BMC Structural Biology</i> , 2015 , 15, 18	2.7	3
56	Association of activated G α to the tumor suppressor Fhit is enhanced by phospholipase C β <i>BMC Cancer</i> , 2015 , 15, 775	4.8	1
55	Angiotensin-[1-12] interacts with angiotensin type I receptors. <i>Neuropharmacology</i> , 2014 , 81, 267-73	5.5	8
54	Synthesis and functional characterization of substituted isoquinolinones as MT2-selective melatoninergic ligands. <i>PLoS ONE</i> , 2014 , 9, e113638	3.7	3
53	G β -mediated activation of protein kinase D exhibits subunit specificity and requires G β -responsive phospholipase C α isoforms. <i>Cell Communication and Signaling</i> , 2013 , 11, 22	7.5	10
52	Activation of G α subunits up-regulates the expression of the tumor suppressor Fhit. <i>Cellular Signalling</i> , 2013 , 25, 2440-52	4.9	3
51	RGS19 inhibits Ras signaling through Nm23H1/2-mediated phosphorylation of the kinase suppressor of Ras. <i>Cellular Signalling</i> , 2013 , 25, 1064-74	4.9	23
50	Regulator of G protein signaling 19 suppresses Ras-induced neoplastic transformation and tumorigenesis. <i>Cancer Letters</i> , 2013 , 339, 33-41	9.9	18
49	Neuronal functions of activators of G protein signaling. <i>NeuroSignals</i> , 2013 , 21, 259-71	1.9	4
48	A molecular and chemical perspective in defining melatonin receptor subtype selectivity. <i>International Journal of Molecular Sciences</i> , 2013 , 14, 18385-406	6.3	22
47	Elevated expression of RGS19 impairs the responsiveness of stress-activated protein kinases to serum. <i>Molecular and Cellular Biochemistry</i> , 2012 , 362, 159-68	4.2	5
46	Activation of Ras-dependent signaling pathways by G(14) -coupled receptors requires the adaptor protein TPR1. <i>Journal of Cellular Biochemistry</i> , 2012 , 113, 3486-97	4.7	9
45	CCR1-mediated STAT3 tyrosine phosphorylation and CXCL8 expression in THP-1 macrophage-like cells involve pertussis toxin-insensitive G(14/16) signaling and IL-6 release. <i>Journal of Immunology</i> , 2012 , 189, 5266-76	5.3	22
44	Pharmacokinetics, oral bioavailability and metabolism of a novel isoquinolinone-based melatonin receptor agonist in rats. <i>Xenobiotica</i> , 2012 , 42, 1138-50	2	4
43	3-Methoxyphenylpropyl amides as novel receptor subtype-selective melatoninergic ligands: characterization of physicochemical and pharmacokinetic properties. <i>Xenobiotica</i> , 2011 , 41, 35-45	2	9
42	Activation of STAT3 by specific G α subunits and multiple G β gamma dimers. <i>International Journal of Biochemistry and Cell Biology</i> , 2010 , 42, 1052-9	5.6	23
41	G α 16 activates Ras by forming a complex with tetratricopeptide repeat 1 (TPR1) and Son of Sevenless (SOS). <i>Cellular Signalling</i> , 2010 , 22, 1448-58	4.9	16
40	CCR1-mediated activation of Nuclear Factor-kappaB in THP-1 monocytic cells involves Pertussis Toxin-insensitive G α (14) and G α (16) signaling cascades. <i>Journal of Leukocyte Biology</i> , 2009 , 86, 1319-29	6.5	12

39	Gbeta3 forms distinct dimers with specific Ggamma subunits and preferentially activates the beta3 isoform of phospholipase C. <i>Cellular Signalling</i> , 2009 , 21, 737-44	4.9	26
38	The RhoA-specific guanine nucleotide exchange factor p63RhoGEF binds to activated Galpha(16) and inhibits the canonical phospholipase Cbeta pathway. <i>Cellular Signalling</i> , 2009 , 21, 1317-25	4.9	13
37	Multiple Gi proteins participate in nerve growth factor-induced activation of c-Jun N-terminal kinases in PC12 cells. <i>Neurochemical Research</i> , 2009 , 34, 1101-12	4.6	12
36	Regulation of mTOR and p70 S6 kinase by the muscarinic M4 receptor in PC12 cells. <i>Cell Biology International</i> , 2009 , 33, 230-8	4.5	7
35	G protein signaling controls the differentiation of multiple cell lineages. <i>BioFactors</i> , 2009 , 35, 232-8	6.1	12
34	Prostacyclin receptor-induced STAT3 phosphorylation in human erythroleukemia cells is mediated via Galpha(s) and Galpha(16) hybrid signaling. <i>Cellular Signalling</i> , 2008 , 20, 2095-106	4.9	9
33	Activation of the human FPRL-1 receptor promotes Ca ²⁺ mobilization in U87 astrocytoma cells. <i>Neurochemical Research</i> , 2008 , 33, 125-33	4.6	5
32	Mutations on the Switch III region and the alpha3 helix of Galpha16 differentially affect receptor coupling and regulation of downstream effectors. <i>Journal of Molecular Signaling</i> , 2008 , 3, 17	1	6
31	Formyl peptide receptor like 1 differentially requires mitogen-activated protein kinases for the induction of glial fibrillary acidic protein and interleukin-1alpha in human U87 astrocytoma cells. <i>Cellular Signalling</i> , 2007 , 19, 2106-17	4.9	24
30	Dopaminergic and adrenergic toxicities on SK-N-MC human neuroblastoma cells are mediated through G protein signaling and oxidative stress. <i>Apoptosis: an International Journal on Programmed Cell Death</i> , 2007 , 12, 167-79	5.4	24
29	Activation of muscarinic M4 receptor augments NGF-induced pro-survival Akt signaling in PC12 cells. <i>Cellular Signalling</i> , 2006 , 18, 285-93	4.9	27
28	Transcriptional activation of c-Fos by constitutively active Galpha(16)QL through a STAT1-dependent pathway. <i>Cellular Signalling</i> , 2006 , 18, 2143-53	4.9	9
27	CKBM stimulates MAPKs but inhibits LPS-induced IFN-gamma in lymphocytes. <i>Phytotherapy Research</i> , 2006 , 20, 725-31	6.7	9
26	Activation of delta-, kappa-, and mu-opioid receptors induces phosphorylation of tuberlin in transfected HEK 293 cells and native cells. <i>Biochemical and Biophysical Research Communications</i> , 2005 , 334, 838-44	3.4	13
25	Integration of G protein signals by extracellular signal-regulated protein kinases in SK-N-MC neuroepithelioma cells. <i>Journal of Neurochemistry</i> , 2005 , 94, 1457-70	6	28
24	Gq-mediated activation of c-Jun N-terminal kinase by the gastrin-releasing peptide-preferring bombesin receptor is inhibited upon costimulation of the Gs-coupled dopamine D1 receptor in COS-7 cells. <i>Molecular Pharmacology</i> , 2005 , 68, 1354-64	4.3	17
23	Epidermal growth factor differentially augments G(i)-mediated stimulation of c-Jun N-terminal kinase activity. <i>British Journal of Pharmacology</i> , 2004 , 142, 635-46	8.6	11
22	Replacement of the alpha5 helix of Galpha16 with Galphas-specific sequences enhances promiscuity of Galpha16 toward Gs-coupled receptors. <i>Cellular Signalling</i> , 2004 , 16, 51-62	4.9	14

21	Differential chemokine activation of CC chemokine receptor 1-regulated pathways: ligand selective activation of Galpha 14-coupled pathways. <i>European Journal of Immunology</i> , 2004 , 34, 785-795	6.1	36
20	The beta6/alpha5 regions of Galphai2 and GalphaoA increase the promiscuity of Galpha16 but are insufficient for pertussis toxin-catalyzed ADP-ribosylation. <i>European Journal of Pharmacology</i> , 2003 , 473, 105-15	5.3	3
19	BML-190 and AM251 act as inverse agonists at the human cannabinoid CB2 receptor: signalling via cAMP and inositol phosphates. <i>FEBS Letters</i> , 2003 , 536, 157-60	3.8	24
18	Melatonin mt1 and MT2 receptors stimulate c-Jun N-terminal kinase via pertussis toxin-sensitive and -insensitive G proteins. <i>Cellular Signalling</i> , 2002 , 14, 249-57	4.9	119
17	Differential Coupling of μ and δ Opioid Receptors to G β 6-Mediated Stimulation of Phospholipase C. <i>Journal of Neurochemistry</i> , 2002 , 70, 2203-2211	6	58
16	Chimeric Galphaq subunits can distinguish the long form of the Xenopus Mel1c melatonin receptor from the mammalian mt1 and MT2 melatonin receptors. <i>Journal of Pineal Research</i> , 2001 , 30, 171-9	10.4	10
15	Galpha(14) links a variety of G(i)- and G(s)-coupled receptors to the stimulation of phospholipase C. <i>British Journal of Pharmacology</i> , 2001 , 132, 1431-40	8.6	49
14	Prostacyclin receptor-independent inhibition of phospholipase C activity by non-prostanoid prostacyclin mimetics. <i>British Journal of Pharmacology</i> , 2001 , 134, 1375-84	8.6	22
13	G(z) signaling: emerging divergence from G(i) signaling. <i>Oncogene</i> , 2001 , 20, 1615-25	9.2	66
12	Opioid-induced adenylyl cyclase supersensitization in human embryonic kidney 293 cells requires pertussis toxin-sensitive G proteins other than G(i1) and G(i3). <i>Neuroscience Letters</i> , 2001 , 299, 25-8	3.3	21
11	Role of extracellular signal-regulated kinases in opioid-induced adenylyl cyclase superactivation in human embryonic kidney 293 cells. <i>Neuroscience Letters</i> , 2001 , 316, 13-6	3.3	12
10	Regulation of adenylyl cyclase, ERK1/2, and CREB by Gz following acute and chronic activation of the delta-opioid receptor. <i>Journal of Neurochemistry</i> , 2000 , 74, 1685-93	6	30
9	Molecular mechanisms and regulation of opioid receptor signaling. <i>Annual Review of Pharmacology and Toxicology</i> , 2000 , 40, 389-430	17.9	528
8	Preactivation permits subsequent stimulation of phospholipase C by G(i)-coupled receptors. <i>Molecular Pharmacology</i> , 2000 , 57, 700-8	4.3	59
7	Stimulation of phospholipase C by the cloned mu, delta and kappa opioid receptors via chimeric G alpha(q) mutants. <i>European Journal of Neuroscience</i> , 1999 , 11, 383-8	3.5	12
6	Regulation of calcium influx and phospholipase C activity by indoleamines in dinoflagellate <i>Cryptocodinium cohnii</i> . <i>Journal of Pineal Research</i> , 1998 , 24, 152-61	10.4	15
5	Chimeric Galpha(q) mutants harboring the last five carboxy-terminal residues of Galpha(i2) or Galpha(o) are resistant to pertussis toxin-catalyzed ADP-ribosylation. <i>FEBS Letters</i> , 1998 , 441, 67-70	3.8	12
4	The calcitonin gene-related peptide-induced acetylcholinesterase synthesis in cultured chick myotubes is mediated by cyclic AMP. <i>Journal of Neurochemistry</i> , 1998 , 71, 152-60	6	36

3	Pertussis toxin-insensitive signaling of the ORL1 receptor: coupling to G α and G16 proteins. <i>Journal of Neurochemistry</i> , 1998 , 71, 2203-10	6	40
2	Functional role of amino-terminal serine16 and serine27 of G alphaZ in receptor and effector coupling. <i>Journal of Neurochemistry</i> , 1997 , 68, 2514-22	6	8
1	ADP-ribosylation with pertussis toxin modulates the GTP-sensitive opioid ligand binding in digitonin-soluble extracts of rat brain membranes. <i>Journal of Neurochemistry</i> , 1988 , 51, 114-21	6	11