

# Daisuke Yasuda

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/245389/publications.pdf>

Version: 2024-02-01

11  
papers

187  
citations

1040056

9  
h-index

1372567

10  
g-index

11  
all docs

11  
docs citations

11  
times ranked

295  
citing authors

#	ARTICLE	IF	CITATIONS
1	Lysophosphatidic Acid Receptor 4 Activation Augments Drug Delivery in Tumors by Tightening Endothelial Cell-Cell Contact. <i>Cell Reports</i> , 2017, 20, 2072-2086.	6.4	29
2	Helix 8 of leukotriene B <sub>4</sub> type 2 receptor is required for the folding to pass the quality control in the endoplasmic reticulum. <i>FASEB Journal</i> , 2009, 23, 1470-1481.	0.5	28
3	Amino Acid Residues Critical for Endoplasmic Reticulum Export and Trafficking of Platelet-activating Factor Receptor. <i>Journal of Biological Chemistry</i> , 2010, 285, 5931-5940.	3.4	27
4	Identification of optineurin as an interleukin-1 receptor-associated kinase 1-binding protein and its role in regulation of MyD88-dependent signaling. <i>Journal of Biological Chemistry</i> , 2017, 292, 17250-17257.	3.4	22
5	The G $\alpha$ 12/13-coupled receptor LPA4 limits proper adipose tissue expansion and remodeling in diet-induced obesity. <i>JCI Insight</i> , 2018, 3, .	5.0	22
6	The lysophosphatidic acid receptor LPA4 regulates hematopoiesis-supporting activity of bone marrow stromal cells. <i>Scientific Reports</i> , 2015, 5, 11410.	3.3	20
7	The atypical N-glycosylation motif, Asn-Cys-Cys, in human GPR109A is required for normal cell surface expression and intracellular signaling. <i>FASEB Journal</i> , 2015, 29, 2412-2422.	0.5	13
8	Characterization of brassinosteroid-regulated proteins in a nuclear-enriched fraction of Arabidopsis suspension-cultured cells. <i>Plant Physiology and Biochemistry</i> , 2011, 49, 985-995.	5.8	12
9	Specific ligands as pharmacological chaperones: The transport of misfolded G-protein coupled receptors to the cell surface. <i>IUBMB Life</i> , 2010, 62, 453-459.	3.4	11
10	Amino Acid Residues of G-Protein-Coupled Receptors Critical for Endoplasmic Reticulum Export and Trafficking. <i>Methods in Enzymology</i> , 2013, 521, 203-216.	1.0	3
11	Helix 8 for ER Export of Leukotriene B <sub>4</sub> type 2 Receptor (BLT2). <i>FASEB Journal</i> , 2010, 24, 1b87.	0.5	0