Hideo Nakanishi

List of Publications by Year
in descending order

21	651	933264 10	940416 16
papers	citations	h-index	g-index

21	21	21	821
all docs	docs citations	times ranked	citing authors

#	Article	IF	Citations
1	Microvascular Density in Glaucomatous Eyes With Hemifield Visual Field Defects: An Optical Coherence Tomography Angiography Study. American Journal of Ophthalmology, 2016, 168, 237-249.	1.7	204
2	Microstructure of Peripapillary Atrophy and Subsequent Visual Field Progression in Treated Primary Open-Angle Glaucoma. Ophthalmology, 2016, 123, 542-551.	2.5	61
3	Lamina Cribrosa Defects and Optic Disc Morphology in Primary Open Angle Glaucoma with High Myopia. PLoS ONE, 2014, 9, e115313.	1.1	53
4	Joint Effect of Cigarette Smoking andCFHandLOC387715/HTRA1Polymorphisms on Polypoidal Choroidal Vasculopathy. , 2010, 51, 6183.		39
5	Significance ofC2/CFBVariants in Age-Related Macular Degeneration and Polypoidal Choroidal Vasculopathy in a Japanese Population. , 2012, 53, 794.		37
6	Sensitivity and specificity for detecting early glaucoma in eyes with high myopia from normative database of macular ganglion cell complex thickness obtained from normal non-myopic or highly myopic Asian eyes. Graefe's Archive for Clinical and Experimental Ophthalmology, 2015, 253, 1143-1152.	1.0	36
7	Association of 15q14 and 15q25 with High Myopia in Japanese. , 2011, 52, 4853.		34
8	Genetic Variants in Pigment Epithelium-Derived Factor Influence Response of Polypoidal Choroidal Vasculopathy to Photodynamic Therapy. Ophthalmology, 2011, 118, 1408-1415.	2.5	24
9	Single-Nucleotide Polymorphisms in the Promoter Region of Matrix Metalloproteinase-1, -2, and -3 in Japanese with High Myopia. , 2010, 51, 4432.		23
10	Association of ARMS2 Genotype With Bilateral Involvement of Exudative Age-Related Macular Degeneration. American Journal of Ophthalmology, 2012, 154, 542-548.e1.	1.7	22
11	Influence of high myopia on outcomes of trabeculectomy with mitomycin C in patients with primary open-angle glaucoma. Japanese Journal of Ophthalmology, 2016, 60, 446-453.	0.9	17
12	Visualization of the Lamina Cribrosa Microvasculature in Normal and Glaucomatous Eyes: A Swept-source Optical Coherence Tomography Angiography Study. Journal of Glaucoma, 2018, 27, 1032-1035.	0.8	17
13	Retinal Blood Flow Velocity Change in Parafoveal Capillary after Topical Tafluprost Treatment in Eyes with Primary Open-angle Glaucoma. Scientific Reports, 2017, 7, 5019.	1.6	16
14	Insulin-like growth factor 1 is not associated with high myopia in a large Japanese cohort. Molecular Vision, 2013, 19, 1074-81.	1.1	16
15	Bilateral papillomacular retinoschisis and macular detachment accompanied by focal lamina cribrosa defect in glaucomatous eyes. Japanese Journal of Ophthalmology, 2014, 58, 435-442.	0.9	13
16	Association of Bruchâ€™s membrane opening and optic disc morphology to axial length and visual field defects in eyes with primary open-angle glaucoma. Graefe's Archive for Clinical and Experimental Ophthalmology, 2018, 256, 599-610.	1.0	12
17	Structural dissociation of optic disc margin components with optic disc tilting: a spectral domain optical coherence tomography study. Graefe's Archive for Clinical and Experimental Ophthalmology, 2016, 254, 343-349.	1.0	11
18	Morphological changes after trabeculectomy in highly myopic eyes with high intraocular pressure by using swept-source optical coherence tomography. American Journal of Ophthalmology Case Reports, 2016, 3, 54-60.	0.4	7

#	Article	IF	Citations
19	Baerveldt or Ahmed glaucoma valve implantation with pars plana tube insertion in Japanese eyes with neovascular glaucoma: 1-year outcomes. Clinical Ophthalmology, 2018, Volume 12, 2439-2449.	0.9	7
20	Long Term Follow-Up of Prototheca keratitis: A Case Report . International Medical Case Reports Journal, 2020, Volume 13, 503-506.	0.3	2
21	Time course of changes in anterior chamber structures after Nd. JCRS Online Case Reports, 2020, 8, e00024.	0.1	0