
## Hideo Nakanishi

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/2470633/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Time course of changes in anterior chamber structures after Nd. JCRS Online Case Reports, 2020, 8, e00024.	0.2	0
2	<p>Long Term Follow-Up of <em>Prototheca</em> keratitis: A Case Report</p> . International Medical Case Reports Journal, 2020, Volume 13, 503-506.	0.8	2
3	Association of Bruch's membrane opening and optic disc morphology to axial length and visual field defects in eyes with primary open-angle glaucoma. Graefe's Archive for Clinical and Experimental Ophthalmology, 2018, 256, 599-610.	1.9	12
4	Baerveldt or Ahmed glaucoma valve implantation with pars plana tube insertion in Japanese eyes with neovascular glaucoma: 1-year outcomes. Clinical Ophthalmology, 2018, Volume 12, 2439-2449.	1.8	7
5	Visualization of the Lamina Cribrosa Microvasculature in Normal and Glaucomatous Eyes: A Swept-source Optical Coherence Tomography Angiography Study. Journal of Glaucoma, 2018, 27, 1032-1035.	1.6	17
6	Retinal Blood Flow Velocity Change in Parafoveal Capillary after Topical Tafluprost Treatment in Eyes with Primary Open-angle Glaucoma. Scientific Reports, 2017, 7, 5019.	3.3	16
7	Morphological changes after trabeculectomy in highly myopic eyes with high intraocular pressure by using swept-source optical coherence tomography. American Journal of Ophthalmology Case Reports, 2016, 3, 54-60.	0.7	7
8	Influence of high myopia on outcomes of trabeculectomy with mitomycin C in patients with primary open-angle glaucoma. Japanese Journal of Ophthalmology, 2016, 60, 446-453.	1.9	17
9	Microvascular Density in Glaucomatous Eyes With Hemifield Visual Field Defects: An Optical Coherence Tomography Angiography Study. American Journal of Ophthalmology, 2016, 168, 237-249.	3.3	204
10	Structural dissociation of optic disc margin components with optic disc tilting: a spectral domain optical coherence tomography study. Graefe's Archive for Clinical and Experimental Ophthalmology, 2016, 254, 343-349.	1.9	11
11	Microstructure of Peripapillary Atrophy and Subsequent Visual Field Progression in Treated Primary Open-Angle Glaucoma. Ophthalmology, 2016, 123, 542-551.	5.2	61
12	Sensitivity and specificity for detecting early glaucoma in eyes with high myopia from normative database of macular ganglion cell complex thickness obtained from normal non-myopic or highly myopic Asian eyes. Graefe's Archive for Clinical and Experimental Ophthalmology, 2015, 253, 1143-1152.	1.9	36
13	Lamina Cribrosa Defects and Optic Disc Morphology in Primary Open Angle Glaucoma with High Myopia. PLoS ONE, 2014, 9, e115313.	2.5	53
14	Bilateral papillomacular retinoschisis and macular detachment accompanied by focal lamina cribrosa defect in glaucomatous eyes. Japanese Journal of Ophthalmology, 2014, 58, 435-442.	1.9	13
15	Insulin-like growth factor 1 is not associated with high myopia in a large Japanese cohort. Molecular Vision, 2013, 19, 1074-81.	1.1	16
16	Significance of <i>C2</i> / <i>CFB</i> Variants in Age-Related Macular Degeneration and Polypoidal Choroidal Vasculopathy in a Japanese Population. , 2012, 53, 794.		37
17	Association of ARMS2 Genotype With Bilateral Involvement of Exudative Age-Related Macular Degeneration. American Journal of Ophthalmology, 2012, 154, 542-548.e1.	3.3	22
18	Genetic Variants in Pigment Epithelium-Derived Factor Influence Response of Polypoidal Choroidal Vasculopathy to Photodynamic Therapy. Ophthalmology, 2011, 118, 1408-1415.	5.2	24

#	Article	IF	CITATIONS
19	Association of 15q14 and 15q25 with High Myopia in Japanese. , 2011, 52, 4853.		34
20	Joint Effect of Cigarette Smoking and <i>CFH</i> and <i>LOC387715/HTRA1</i> Polymorphisms on Polypoidal Choroidal Vasculopathy. , 2010, 51, 6183.		39
21	Single-Nucleotide Polymorphisms in the Promoter Region of Matrix Metalloproteinase-1, -2, and -3 in Japanese with High Myopia. , 2010, 51, 4432.		23