

# Fang-Jie Zhao

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2473172/publications.pdf>

Version: 2024-02-01

368  
papers

45,896  
citations

863

117  
h-index

2375

198  
g-index

377  
all docs

377  
docs citations

377  
times ranked

23164  
citing authors

#	ARTICLE	IF	CITATIONS
1	Dietary cadmium exposure, risks to human health and mitigation strategies. <i>Critical Reviews in Environmental Science and Technology</i> , 2023, 53, 939-963.	6.6	37
2	Oxidation of organoarsenicals and antimonite by a novel flavin monooxygenase widely present in soil bacteria. <i>Environmental Microbiology</i> , 2022, 24, 752-761.	1.8	26
3	Toxic metals and metalloids: Uptake, transport, detoxification, phytoremediation, and crop improvement for safer food. <i>Molecular Plant</i> , 2022, 15, 27-44.	3.9	131
4	What is a plant nutrient? Changing definitions to advance science and innovation in plant nutrition. <i>Plant and Soil</i> , 2022, 476, 11-23.	1.8	38
5	The relative contributions of root uptake and remobilization to the loading of Cd and As into rice grains: Implications in simultaneously controlling grain Cd and As accumulation using a segmented water management strategy. <i>Environmental Pollution</i> , 2022, 293, 118497.	3.7	47
6	Soil amendments with ZnSO <sub>4</sub> or MnSO <sub>4</sub> are effective at reducing Cd accumulation in rice grain: An application of the voltaic cell principle. <i>Environmental Pollution</i> , 2022, 294, 118650.	3.7	11
7	Functional characterization of the methylarsenite-inducible arsRM operon from <i>Noviherbaspirillum denitrificans</i> HC18. <i>Environmental Microbiology</i> , 2022, , .	1.8	6
8	Glutathione Is Involved in the Reduction of Methylarsenate to Generate Antibiotic Methylarsenite in <i>Enterobacter</i> sp. Strain CZ-1. <i>Applied and Environmental Microbiology</i> , 2022, 88, aem0246721.	1.4	4
9	Elucidating heterogeneous iron biomineralization patterns in a denitrifying As(III)-oxidizing bacterium: implications for arsenic immobilization. <i>Environmental Science: Nano</i> , 2022, 9, 1076-1090.	2.2	5
10	The vacuolar transporter OsNRAMP2 mediates Fe remobilization during germination and affects Cd distribution to rice grain. <i>Plant and Soil</i> , 2022, 476, 79-95.	1.8	12
11	Widespread Occurrence of the Highly Toxic Dimethylated Monothioarsenate (DMMTA) in Rice Globally. <i>Environmental Science &amp; Technology</i> , 2022, 56, 3575-3586.	4.6	27
12	ArsZ from <i>Ensifer adhaerens</i> ST2 is a novel methylarsenite oxidase. <i>Environmental Microbiology</i> , 2022, 24, 3013-3021.	1.8	6
13	China national food safety standards of cadmium in staple foods: Issues and thinking. <i>Chinese Science Bulletin</i> , 2022, 67, 3252-3260.	0.4	2
14	The Vacuolar Molybdate Transporter OsMOT1;2 Controls Molybdenum Remobilization in Rice. <i>Frontiers in Plant Science</i> , 2022, 13, 863816.	1.7	1
15	Variation in cadmium accumulation and speciation within the same population of the hyperaccumulator <i>Noccaea caerulescens</i> grown in a moderately contaminated soil. <i>Plant and Soil</i> , 2022, 475, 379-394.	1.8	7
16	Significant Nutritional Gaps in Tibetan Adults Living in Agricultural Counties Along Yarlung Zangbo River. <i>Frontiers in Nutrition</i> , 2022, 9, 845026.	1.6	5
17	Prevalent and highly mobile antibiotic resistance genes in commercial organic fertilizers. <i>Environment International</i> , 2022, 162, 107157.	4.8	21
18	Molybdenum: More than an essential element. <i>Journal of Experimental Botany</i> , 2022, 73, 1766-1774.	2.4	25

#	ARTICLE	IF	CITATIONS
19	Exploring Key Soil Parameters Relevant to Arsenic and Cadmium Accumulation in Rice Grain in Southern China. <i>Soil Systems</i> , 2022, 6, 36.	1.0	4
20	Suppression of methanogenesis in paddy soil increases dimethylarsenate accumulation and the incidence of straighthead disease in rice. <i>Soil Biology and Biochemistry</i> , 2022, 169, 108689.	4.2	14
21	Local and Systemic Response to Heterogeneous Sulfate Resupply after Sulfur Deficiency in Rice. <i>International Journal of Molecular Sciences</i> , 2022, 23, 6203.	1.8	3
22	The roles of membrane transporters in arsenic uptake, translocation and detoxification in plants. <i>Critical Reviews in Environmental Science and Technology</i> , 2021, 51, 2449-2484.	6.6	51
23	Producing Cd-safe rice grains in moderately and seriously Cd-contaminated paddy soils. <i>Chemosphere</i> , 2021, 267, 128893.	4.2	25
24	Stable isotope fractionation of cadmium in the soil-rice-human continuum. <i>Science of the Total Environment</i> , 2021, 761, 143262.	3.9	28
25	Reducing cadmium bioavailability and accumulation in vegetable by an alkalizing bacterial strain. <i>Science of the Total Environment</i> , 2021, 758, 143596.	3.9	18
26	Cadmium Inhibits Lateral Root Emergence in Rice by Disrupting OsPIN-Mediated Auxin Distribution and the Protective Effect of OsHMA3. <i>Plant and Cell Physiology</i> , 2021, 62, 166-177.	1.5	21
27	Cadmium speciation and release kinetics in a paddy soil as affected by soil amendments and flooding-draining cycle. <i>Environmental Pollution</i> , 2021, 268, 115944.	3.7	27
28	High-Affinity Sulfate Transporter Sultr1;2 Is a Major Transporter for Cr(VI) Uptake in Plants. <i>Environmental Science &amp; Technology</i> , 2021, 55, 1576-1584.	4.6	41
29	The Voltaic Effect as a Novel Mechanism Controlling the Remobilization of Cadmium in Paddy Soils during Drainage. <i>Environmental Science &amp; Technology</i> , 2021, 55, 1750-1758.	4.6	59
30	Sulfate addition and rising temperature promote arsenic methylation and the formation of methylated thioarsenates in paddy soils. <i>Soil Biology and Biochemistry</i> , 2021, 154, 108129.	4.2	38
31	DNA hypomethylation in tetraploid rice potentiates stress-responsive gene expression for salt tolerance. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2021, 118, .	3.3	44
32	A molecular switch in sulfur metabolism to reduce arsenic and enrich selenium in rice grain. <i>Nature Communications</i> , 2021, 12, 1392.	5.8	48
33	Dynamics of Dimethylated Monothioarsenate (DMMTA) in Paddy Soils and Its Accumulation in Rice Grains. <i>Environmental Science &amp; Technology</i> , 2021, 55, 8665-8674.	4.6	25
34	Free Radicals Produced from the Oxidation of Ferrous Sulfides Promote the Remobilization of Cadmium in Paddy Soils During Drainage. <i>Environmental Science &amp; Technology</i> , 2021, 55, 9845-9853.	4.6	63
35	Food Consumption and Dietary Patterns of Local Adults Living on the Tibetan Plateau: Results from 14 Countries along the Yarlung Tsangpo River. <i>Nutrients</i> , 2021, 13, 2444.	1.7	17
36	Future research needs for environmental science in China. <i>Geography and Sustainability</i> , 2021, , .	1.9	3

#	ARTICLE	IF	CITATIONS
37	Two-year and multi-site field trials to evaluate soil amendments for controlling cadmium accumulation in rice grain. <i>Environmental Pollution</i> , 2021, 289, 117918.	3.7	20
38	Phylogenomics reveals the basis of adaptation of <i>Pseudorhizobium</i> species to extreme environments and supports a taxonomic revision of the genus. <i>Systematic and Applied Microbiology</i> , 2021, 44, 126165.	1.2	33
39	Univariate and Multivariate QTL Analyses Reveal Covariance Among Mineral Elements in the Rice Ionome. <i>Frontiers in Genetics</i> , 2021, 12, 638555.	1.1	10
40	Targeted expression of the arsenate reductase HAC1 identifies cell type specificity of arsenic metabolism and transport in plant roots. <i>Journal of Experimental Botany</i> , 2021, 72, 415-425.	2.4	12
41	<sc>ArsV</sc> and <sc>ArsW</sc> provide synergistic resistance to the antibiotic methylarsenite. <i>Environmental Microbiology</i> , 2021, 23, 7550-7562.	1.8	11
42	Demethylation of the Antibiotic Methylarsenite is Coupled to Denitrification in Anoxic Paddy Soil. <i>Environmental Science &amp; Technology</i> , 2021, 55, 15484-15494.	4.6	13
43	Natural variation in the promoter of <i>OsHMA3</i> contributes to differential grain cadmium accumulation between <i>Indica</i> and <i>Japonica</i> rice. <i>Journal of Integrative Plant Biology</i> , 2020, 62, 314-329.	4.1	72
44	Protein phosphatase 2A alleviates cadmium toxicity by modulating ethylene production in <sc><i>Arabidopsis thaliana</i></sc>. <i>Plant, Cell and Environment</i> , 2020, 43, 1008-1022.	2.8	13
45	Binding and adsorption energy of Cd in soils and its environmental implication for Cd bioavailability. <i>Soil Science Society of America Journal</i> , 2020, 84, 472-482.	1.2	10
46	<i>N</i>-Hydroxyarylamine <i>O</i>-Acetyltransferases Catalyze Acetylation of 3-Amino-4-Hydroxyphenylarsonic Acid in the 4-Hydroxy-3-Nitrobenzenearsonic Acid Transformation Pathway of <i>Enterobacter</i> sp. Strain CZ-1. <i>Applied and Environmental Microbiology</i> , 2020, 86, .	1.4	9
47	Arsenic and cadmium accumulation in rice and mitigation strategies. <i>Plant and Soil</i> , 2020, 446, 1-21.	1.8	327
48	Genetic mapping of ionomic quantitative trait loci in rice grain and straw reveals OsMOT1;1 as the putative causal gene for a molybdenum QTL qMo8. <i>Molecular Genetics and Genomics</i> , 2020, 295, 391-407.	1.0	20
49	Mutation in <i>OsCADT1</i> enhances cadmium tolerance and enriches selenium in rice grain. <i>New Phytologist</i> , 2020, 226, 838-850.	3.5	45
50	Overexpression of Rice <i>OsHMA3</i> in Wheat Greatly Decreases Cadmium Accumulation in Wheat Grains. <i>Environmental Science &amp; Technology</i> , 2020, 54, 10100-10108.	4.6	72
51	<sc>OsNRAMP1</sc> transporter contributes to cadmium and manganese uptake in rice. <i>Plant, Cell and Environment</i> , 2020, 43, 2476-2491.	2.8	191
52	Breeding for low cadmium barley by introgression of a Sukkula-like transposable element. <i>Nature Food</i> , 2020, 1, 489-499.	6.2	44
53	Chemical Speciation and Distribution of Cadmium in Rice Grain and Implications for Bioavailability to Humans. <i>Environmental Science &amp; Technology</i> , 2020, 54, 12072-12080.	4.6	46
54	Overexpression of the manganese/cadmium transporter OsNRAMP5 reduces cadmium accumulation in rice grain. <i>Journal of Experimental Botany</i> , 2020, 71, 5705-5715.	2.4	75

#	ARTICLE	IF	CITATIONS
55	Microbe mediated immobilization of arsenic in the rice rhizosphere after incorporation of silica impregnated biochar composites. <i>Journal of Hazardous Materials</i> , 2020, 398, 123096.	6.5	46
56	OASTL-A1 functions as a cytosolic cysteine synthase and affects arsenic tolerance in rice. <i>Journal of Experimental Botany</i> , 2020, 71, 3678-3689.	2.4	19
57	Nitrite Accumulation Is Required for Microbial Anaerobic Iron Oxidation, but Not for Arsenite Oxidation, in Two Heterotrophic Denitrifiers. <i>Environmental Science &amp; Technology</i> , 2020, 54, 4036-4045.	4.6	33
58	Dimethylarsinic acid is the causal agent inducing rice straighthead disease. <i>Journal of Experimental Botany</i> , 2020, 71, 5631-5644.	2.4	40
59	The within-field spatial variation in rice grain Cd concentration is determined by soil redox status and pH during grain filling. <i>Environmental Pollution</i> , 2020, 261, 114151.	3.7	55
60	QTL pyramiding for producing nutritious and safe rice grains. <i>Journal of Integrative Plant Biology</i> , 2020, 62, 264-268.	4.1	4
61	Localized Intensification of Arsenic Release within the Emergent Rice Rhizosphere. <i>Environmental Science &amp; Technology</i> , 2020, 54, 3138-3147.	4.6	34
62	Increased arsenic mobilization in the rice rhizosphere is mediated by iron-reducing bacteria. <i>Environmental Pollution</i> , 2020, 263, 114561.	3.7	35
63	Strategies to manage the risk of heavy metal(loid) contamination in agricultural soils. <i>Frontiers of Agricultural Science and Engineering</i> , 2020, 7, 333.	0.9	11
64	Variation in the BrHMA3 coding region controls natural variation in cadmium accumulation in Brassica rapa vegetables. <i>Journal of Experimental Botany</i> , 2019, 70, 5865-5878.	2.4	36
65	Special section on soil and human health “An editorial. <i>European Journal of Soil Science</i> , 2019, 70, 859-861.	1.8	2
66	Dynamics of metal(loid) resistance genes driven by succession of bacterial community during manure composting. <i>Environmental Pollution</i> , 2019, 255, 113276.	3.7	16
67	Water management impacts the soil microbial communities and total arsenic and methylated arsenicals in rice grains. <i>Environmental Pollution</i> , 2019, 247, 736-744.	3.7	68
68	Biotransformation of arsenic-containing roxarsone by an aerobic soil bacterium <i>Enterobacter</i> sp. CZ-1. <i>Environmental Pollution</i> , 2019, 247, 482-487.	3.7	28
69	Sulfate-reducing bacteria and methanogens are involved in arsenic methylation and demethylation in paddy soils. <i>ISME Journal</i> , 2019, 13, 2523-2535.	4.4	122
70	Microbial sulfate reduction decreases arsenic mobilization in flooded paddy soils with high potential for microbial Fe reduction. <i>Environmental Pollution</i> , 2019, 251, 952-960.	3.7	61
71	Engineering Crops without Genome Integration Using Nanotechnology. <i>Trends in Plant Science</i> , 2019, 24, 574-577.	4.3	48
72	Epigenetic regulation of sulfur homeostasis in plants. <i>Journal of Experimental Botany</i> , 2019, 70, 4171-4182.	2.4	28

#	ARTICLE	IF	CITATIONS
73	Cadmium contamination in agricultural soils of China and the impact on food safety. <i>Environmental Pollution</i> , 2019, 249, 1038-1048.	3.7	395
74	Producing cadmium-free Indica rice by overexpressing OsHMA3. <i>Environment International</i> , 2019, 126, 619-626.	4.8	110
75	The C-type ATP-Binding Cassette Transporter OsABCC7 Is Involved in the Root-to-Shoot Translocation of Arsenic in Rice. <i>Plant and Cell Physiology</i> , 2019, 60, 1525-1535.	1.5	48
76	Map-based cloning of a new total loss-of-function allele of OsHMA3 causes high cadmium accumulation in rice grain. <i>Journal of Experimental Botany</i> , 2019, 70, 2857-2871.	2.4	57
77	Iron and Manganese (Oxyhydro)oxides, Rather than Oxidation of Sulfides, Determine Mobilization of Cd during Soil Drainage in Paddy Soil Systems. <i>Environmental Science &amp; Technology</i> , 2019, 53, 2500-2508.	4.6	236
78	Natural variation in a molybdate transporter controls grain molybdenum concentration in rice. <i>New Phytologist</i> , 2019, 221, 1983-1997.	3.5	44
79	SpHMA1 is a chloroplast cadmium exporter protecting photochemical reactions in the Cd hyperaccumulator <i>Sedum plumbizincicola</i> . <i>Plant, Cell and Environment</i> , 2019, 42, 1112-1124.	2.8	49
80	Minimizing experimental artefacts in synchrotron-based X-ray analyses of Fe speciation in tissues of rice plants. <i>Journal of Synchrotron Radiation</i> , 2019, 26, 1272-1279.	1.0	7
81	Cadmium Phytoremediation: Call Rice CAL1. <i>Molecular Plant</i> , 2018, 11, 640-642.	3.9	31
82	Soil and human health. <i>European Journal of Soil Science</i> , 2018, 69, 158-158.	1.8	2
83	Microbe mediated arsenic release from iron minerals and arsenic methylation in rhizosphere controls arsenic fate in soil-rice system after straw incorporation. <i>Environmental Pollution</i> , 2018, 236, 598-608.	3.7	118
84	Geographical variations of cadmium and arsenic concentrations and arsenic speciation in Chinese rice. <i>Environmental Pollution</i> , 2018, 238, 482-490.	3.7	148
85	Risk of Silver Transfer from Soil to the Food Chain Is Low after Long-Term (20 Years) Field Applications of Sewage Sludge. <i>Environmental Science &amp; Technology</i> , 2018, 52, 4901-4909.	4.6	39
86	Dissecting the components controlling root-to-shoot arsenic translocation in <i>Arabidopsis thaliana</i> . <i>New Phytologist</i> , 2018, 217, 206-218.	3.5	56
87	Antibiotics and antibiotic resistance from animal manures to soil: a review. <i>European Journal of Soil Science</i> , 2018, 69, 181-195.	1.8	291
88	Arsenic methylation by a novel ArsM As(III) S-adenosylmethionine methyltransferase that requires only two conserved cysteine residues. <i>Molecular Microbiology</i> , 2018, 107, 265-276.	1.2	42
89	Nramp5 expression and functionality likely explain higher cadmium uptake in rice than in wheat and maize. <i>Plant and Soil</i> , 2018, 433, 377-389.	1.8	111
90	ARSENATE INDUCED CHLOROSIS 1/ TRANSLOCON AT THE OUTER ENVELOPE MEMBRANE OF CHLOROPLASTS 132 Protects Chloroplasts from Arsenic Toxicity. <i>Plant Physiology</i> , 2018, 178, 1568-1583.	2.3	18

#	ARTICLE	IF	CITATIONS
91	Genome-Wide Association Studies Reveal the Genetic Basis of Ionic Variation in Rice. <i>Plant Cell</i> , 2018, 30, 2720-2740.	3.1	164
92	Engineered silver nanoparticles in terrestrial environments: a meta-analysis shows that the overall environmental risk is small. <i>Environmental Science: Nano</i> , 2018, 5, 2531-2544.	2.2	25
93	OsWRKY28 Regulates Phosphate and Arsenate Accumulation, Root System Architecture and Fertility in Rice. <i>Frontiers in Plant Science</i> , 2018, 9, 1330.	1.7	61
94	Effective methods to reduce cadmium accumulation in rice grain. <i>Chemosphere</i> , 2018, 207, 699-707.	4.2	170
95	Long-term effects of manure and chemical fertilizers on soil antibiotic resistome. <i>Soil Biology and Biochemistry</i> , 2018, 122, 111-119.	4.2	98
96	Dietary cadmium intake from rice and vegetables and potential health risk: A case study in Xiangtan, southern China. <i>Science of the Total Environment</i> , 2018, 639, 271-277.	3.9	231
97	OsATX1 Interacts with Heavy Metal P1B-Type ATPases and Affects Copper Transport and Distribution. <i>Plant Physiology</i> , 2018, 178, 329-344.	2.3	96
98	Decreasing arsenic accumulation in rice by overexpressing <i>OsNIP1;1</i> and <i>OsNIP3;3</i> through disrupting arsenite radial transport in roots. <i>New Phytologist</i> , 2018, 219, 641-653.	3.5	122
99	Particle-specific toxicity and bioavailability of cerium oxide (CeO <sub>2</sub> ) nanoparticles to <i>Arabidopsis thaliana</i> . <i>Journal of Hazardous Materials</i> , 2017, 322, 292-300.	6.5	90
100	Mineral Availability as a Key Regulator of Soil Carbon Storage. <i>Environmental Science &amp; Technology</i> , 2017, 51, 4960-4969.	4.6	167
101	OsHAC4 is critical for arsenate tolerance and regulates arsenic accumulation in rice. <i>New Phytologist</i> , 2017, 215, 1090-1101.	3.5	156
102	OsPTR7 (OsNPF8.1), a Putative Peptide Transporter in Rice, is Involved in Dimethylarsenate Accumulation in Rice Grain. <i>Plant and Cell Physiology</i> , 2017, 58, 904-913.	1.5	65
103	Predicting Cadmium Safety Thresholds in Soils Based on Cadmium Uptake by Chinese Cabbage. <i>Pedosphere</i> , 2017, 27, 475-481.	2.1	33
104	Soil Environment and Pollution Remediation. <i>Pedosphere</i> , 2017, 27, 387-388.	2.1	9
105	The Nodulin 26-like intrinsic membrane protein OsNIP3;2 is involved in arsenite uptake by lateral roots in rice. <i>Journal of Experimental Botany</i> , 2017, 68, 3007-3016.	2.4	84
106	Heavy metal ATPase 3 (HMA3) confers cadmium hypertolerance on the cadmium/zinc hyperaccumulator <i>Sedum plumbizincicola</i> . <i>New Phytologist</i> , 2017, 215, 687-698.	3.5	191
107	Historical trends in iodine and selenium in soil and herbage at the Park Grass Experiment, Rothamsted Research, UK. <i>Soil Use and Management</i> , 2017, 33, 252-262.	2.6	15
108	Nitrate Stimulates Anaerobic Microbial Arsenite Oxidation in Paddy Soils. <i>Environmental Science &amp; Technology</i> , 2017, 51, 4377-4386.	4.6	95



#	ARTICLE	IF	CITATIONS
109	Genotypic and Environmental Variations in Grain Cadmium and Arsenic Concentrations Among a Panel of High Yielding Rice Cultivars. <i>Rice</i> , 2017, 10, 9.	1.7	124
110	Arsenic methylation by a genetically engineered Rhizobium-legume symbiont. <i>Plant and Soil</i> , 2017, 416, 259-269.	1.8	48
111	Heavy metal concentrations and arsenic speciation in animal manure composts in China. <i>Waste Management</i> , 2017, 64, 333-339.	3.7	158
112	Characterizing the uptake, accumulation and toxicity of silver sulfide nanoparticles in plants. <i>Environmental Science: Nano</i> , 2017, 4, 448-460.	2.2	85
113	Determining the fate of selenium in wheat biofortification: an isotopically labelled field trial study. <i>Plant and Soil</i> , 2017, 420, 61-77.	1.8	24
114	Control of arsenic mobilization in paddy soils by manganese and iron oxides. <i>Environmental Pollution</i> , 2017, 231, 37-47.	3.7	145
115	Microbial Processes Mediating the Evolution of Methylarsine Gases from Dimethylarsenate in Paddy Soils. <i>Environmental Science &amp; Technology</i> , 2017, 51, 13190-13198.	4.6	12
116	Allelic Variation of NtNramp5 Associated with Cultivar Variation in Cadmium Accumulation in Tobacco. <i>Plant and Cell Physiology</i> , 2017, 58, 1583-1593.	1.5	41
117	Bacterial community and arsenic functional genes diversity in arsenic contaminated soils from different geographic locations. <i>PLoS ONE</i> , 2017, 12, e0176696.	1.1	40
118	<i>Arsenicibacter rosenii</i> gen. nov., sp. nov., an efficient arsenic methylating and volatilizing bacterium isolated from an arsenic-contaminated paddy soil. <i>International Journal of Systematic and Evolutionary Microbiology</i> , 2017, 67, 3186-3191.	0.8	11
119	A novel pathway of arsenate detoxification. <i>Molecular Microbiology</i> , 2016, 100, 928-930.	1.2	21
120	OsCLT1, a CRT-like transporter 1, is required for glutathione homeostasis and arsenic tolerance in rice. <i>New Phytologist</i> , 2016, 211, 658-670.	3.5	75
121	Arsenic Methylation in <i>Arabidopsis thaliana</i> Expressing an Algal Arsenite Methyltransferase Gene Increases Arsenic Phytotoxicity. <i>Journal of Agricultural and Food Chemistry</i> , 2016, 64, 2674-2681.	2.4	39
122	Nanotechnology: A New Opportunity in Plant Sciences. <i>Trends in Plant Science</i> , 2016, 21, 699-712.	4.3	690
123	The role of OsPT8 in arsenate uptake and varietal difference in arsenate tolerance in rice. <i>Journal of Experimental Botany</i> , 2016, 67, 6051-6059.	2.4	158
124	OsHAC1;1 and OsHAC1;2 Function as Arsenate Reductases and Regulate Arsenic Accumulation. <i>Plant Physiology</i> , 2016, 172, 1708-1719.	2.3	200
125	Long-Term Impact of Field Applications of Sewage Sludge on Soil Antibiotic Resistome. <i>Environmental Science &amp; Technology</i> , 2016, 50, 12602-12611.	4.6	97
126	Isolation and Characterization of an Aluminum-resistant Mutant in Rice. <i>Rice</i> , 2016, 9, 60.	1.7	15



#	ARTICLE	IF	CITATIONS
127	Changes in antibiotic concentrations and antibiotic resistome during commercial composting of animal manures. <i>Environmental Pollution</i> , 2016, 219, 182-190.	3.7	166
128	Efficient Arsenic Methylation and Volatilization Mediated by a Novel Bacterium from an Arsenic-Contaminated Paddy Soil. <i>Environmental Science &amp; Technology</i> , 2016, 50, 6389-6396.	4.6	86
129	A loss-of-function allele of <i>OsHMA3</i> associated with high cadmium accumulation in shoots and grain of <i>Japonica</i> rice cultivars. <i>Plant, Cell and Environment</i> , 2016, 39, 1941-1954.	2.8	168
130	Aluminium alleviates fluoride toxicity in tea ( <i>Camellia sinensis</i> ). <i>Plant and Soil</i> , 2016, 402, 179-190.	1.8	42
131	The shift of the microbial community in activated sludge with calcium treatment and its implication to sludge settleability. <i>Bioresource Technology</i> , 2016, 207, 11-18.	4.8	46
132	Environmental factors influencing aluminium accumulation in tea ( <i>Camellia sinensis</i> L.). <i>Plant and Soil</i> , 2016, 400, 223-230.	1.8	16
133	Phytotoxicity and detoxification mechanism differ among inorganic and methylated arsenic species in <i>Arabidopsis thaliana</i> . <i>Plant and Soil</i> , 2016, 401, 243-257.	1.8	47
134	Concentrations of metals and metalloids in soils that have the potential to lead to exceedance of maximum limit concentrations of contaminants in food and feed. <i>Soil Use and Management</i> , 2015, 31, 34-45.	2.6	21
135	Diversity and Abundance of Arsenic Biotransformation Genes in Paddy Soils from Southern China. <i>Environmental Science &amp; Technology</i> , 2015, 49, 4138-4146.	4.6	195
136	Distribution of the stable isotopes <sup>57</sup> Fe and <sup>68</sup> Zn in grain tissues of various wheat lines differing in their phytate content. <i>Plant and Soil</i> , 2015, 396, 73-83.	1.8	22
137	The role of nodes in arsenic storage and distribution in rice. <i>Journal of Experimental Botany</i> , 2015, 66, 3717-3724.	2.4	99
138	Genetically Engineering <i>Bacillus subtilis</i> with a Heat-Resistant Arsenite Methyltransferase for Bioremediation of Arsenic-Contaminated Organic Waste. <i>Applied and Environmental Microbiology</i> , 2015, 81, 6718-6724.	1.4	68
139	Iron and zinc isotope fractionation during uptake and translocation in rice ( <i>Oryza sativa</i> ) grown in oxic and anoxic soils. <i>Comptes Rendus - Geoscience</i> , 2015, 347, 397-404.	0.4	37
140	Anaerobic Arsenite Oxidation by an Autotrophic Arsenite-Oxidizing Bacterium from an Arsenic-Contaminated Paddy Soil. <i>Environmental Science &amp; Technology</i> , 2015, 49, 5956-5964.	4.6	121
141	Genome-wide transcriptomic and phylogenetic analyses reveal distinct aluminum-tolerance mechanisms in the aluminum-accumulating species buckwheat ( <i>Fagopyrum tataricum</i> ). <i>BMC Plant Biology</i> , 2015, 15, 16.	1.6	48
142	Arsenic Methylation and Volatilization by Arsenite <i>S</i> -Adenosylmethionine Methyltransferase in <i>Pseudomonas alcaligenes</i> NBRC14159. <i>Applied and Environmental Microbiology</i> , 2015, 81, 2852-2860.	1.4	84
143	Accumulation and phytotoxicity of perfluorooctanoic acid in the model plant species <i>Arabidopsis thaliana</i> . <i>Environmental Pollution</i> , 2015, 206, 560-566.	3.7	52
144	Soil Contamination in China: Current Status and Mitigation Strategies. <i>Environmental Science &amp; Technology</i> , 2015, 49, 750-759.	4.6	1,488

#	ARTICLE	IF	CITATIONS
145	Transcriptional and physiological analyses identify a regulatory role for hydrogen peroxide in the lignin biosynthesis of copper-stressed rice roots. <i>Plant and Soil</i> , 2015, 387, 323-336.	1.8	79
146	Soil pollution and soil organisms: an overview of research progress and perspectives. <i>Acta Ecologica Sinica</i> , 2015, 35, .	0.0	1
147	Genome-wide Association Mapping Identifies a New Arsenate Reductase Enzyme Critical for Limiting Arsenic Accumulation in Plants. <i>PLoS Biology</i> , 2014, 12, e1002009.	2.6	227
148	Genome Wide Association Mapping of Grain Arsenic, Copper, Molybdenum and Zinc in Rice ( <i>Oryza</i> ) Tj ETQq0 0 0 rgBT /Overlock 10 Tf 5	1.1	228
149	OsNRAMP5 contributes to manganese translocation and distribution in rice shoots. <i>Journal of Experimental Botany</i> , 2014, 65, 4849-4861.	2.4	211
150	Combined NanoSIMS and synchrotron X-ray fluorescence reveal distinct cellular and subcellular distribution patterns of trace elements in rice tissues. <i>New Phytologist</i> , 2014, 201, 104-115.	3.5	157
151	Silicon has opposite effects on the accumulation of inorganic and methylated arsenic species in rice. <i>Plant and Soil</i> , 2014, 376, 423-431.	1.8	73
152	Lead in rice: Analysis of baseline lead levels in market and field collected rice grains. <i>Science of the Total Environment</i> , 2014, 485-486, 428-434.	3.9	78
153	Imaging element distribution and speciation in plant cells. <i>Trends in Plant Science</i> , 2014, 19, 183-192.	4.3	138
154	Earth Abides Arsenic Biotransformations. <i>Annual Review of Earth and Planetary Sciences</i> , 2014, 42, 443-467.	4.6	423
155	Iron Bioavailability in Two Commercial Cultivars of Wheat: Comparison between Wholegrain and White Flour and the Effects of Nicotianamine and 2-Deoxymugineic Acid on Iron Uptake into Caco-2 Cells. <i>Journal of Agricultural and Food Chemistry</i> , 2014, 62, 10320-10325.	2.4	60
156	Impact of agronomic practices on arsenic accumulation and speciation in rice grain. <i>Environmental Pollution</i> , 2014, 194, 217-223.	3.7	104
157	Distribution and Speciation of Iron and Zinc in Grain of Two Wheat Genotypes. <i>Journal of Agricultural and Food Chemistry</i> , 2014, 62, 708-716.	2.4	70
158	Effects of Nitrogen on the Distribution and Chemical Speciation of Iron and Zinc in Pearling Fractions of Wheat Grain. <i>Journal of Agricultural and Food Chemistry</i> , 2014, 62, 4738-4746.	2.4	50
159	Understanding Elemental Uptake in Plants Using High Resolution SIMS and Complementary Techniques. <i>Microscopy and Microanalysis</i> , 2014, 20, 1316-1317.	0.2	0
160	High resolution SIMS analysis of arsenic in rice. <i>Surface and Interface Analysis</i> , 2013, 45, 309-311.	0.8	12
161	Use of Synchrotron-Based Techniques to Elucidate Metal Uptake and Metabolism in Plants. <i>Advances in Agronomy</i> , 2013, , 1-82.	2.4	70
162	Methylated Arsenic Species in Rice: Geographical Variation, Origin, and Uptake Mechanisms. <i>Environmental Science &amp; Technology</i> , 2013, 47, 3957-3966.	4.6	276

#	ARTICLE	IF	CITATIONS
163	Assessing the contributions of lateral roots to element uptake in rice using an auxin-related lateral root mutant. <i>Plant and Soil</i> , 2013, 372, 125-136.	1.8	26
164	Variation in Rice Cadmium Related to Human Exposure. <i>Environmental Science &amp; Technology</i> , 2013, 47, 5613-5618.	4.6	365
165	Historical arsenic contamination of soil due to long-term phosphate fertiliser applications. <i>Environmental Pollution</i> , 2013, 180, 259-264.	3.7	59
166	Arsenic Methylation in Soils and Its Relationship with Microbial <i>arsM</i> Abundance and Diversity, and As Speciation in Rice. <i>Environmental Science &amp; Technology</i> , 2013, 47, 7147-7154.	4.6	166
167	Proteomic analysis of copper stress responses in the roots of two rice ( <i>Oryza sativa</i> L.) varieties differing in Cu tolerance. <i>Plant and Soil</i> , 2013, 366, 647-658.	1.8	78
168	Adaptation to acidic soil is achieved by increased numbers of <i>cis</i> -acting elements regulating <i>ALMT1</i> expression in <i>Holcus lanatus</i> . <i>Plant Journal</i> , 2013, 76, 10-23.	2.8	60
169	Biogeochemistry of Arsenic in Paddy Environments. , 2012, , 71-101.		13
170	Heterologous expression of the yeast arsenite efflux system ACR3 improves <i>Arabidopsis thaliana</i> tolerance to arsenic stress. <i>New Phytologist</i> , 2012, 194, 716-723.	3.5	66
171	Pathways and Relative Contributions to Arsenic Volatilization from Rice Plants and Paddy Soil. <i>Environmental Science &amp; Technology</i> , 2012, 46, 8090-8096.	4.6	131
172	Selenium speciation in soil extracts using LC-ICP-MS. <i>International Journal of Environmental Analytical Chemistry</i> , 2012, 92, 222-236.	1.8	32
173	Arsenic & Rice. , 2012, , .		92
174	Identifying the species of copper that are toxic to plant roots in alkaline nutrient solutions. <i>Plant and Soil</i> , 2012, 361, 317-327.	1.8	14
175	Contrasting effects of dwarfing alleles and nitrogen availability on mineral concentrations in wheat grain. <i>Plant and Soil</i> , 2012, 360, 93-107.	1.8	25
176	Beneficial Elements. , 2012, , 249-269.		70
177	Function of Nutrients. , 2012, , 191-248.		383
178	Knocking Out ACR2 Does Not Affect Arsenic Redox Status in <i>Arabidopsis thaliana</i> : Implications for As Detoxification and Accumulation in Plants. <i>PLoS ONE</i> , 2012, 7, e42408.	1.1	34
179	Elemental imaging at the nanoscale: NanoSIMS and complementary techniques for element localisation in plants. <i>Analytical and Bioanalytical Chemistry</i> , 2012, 402, 3263-3273.	1.9	131
180	Methylated arsenic species in plants originate from soil microorganisms. <i>New Phytologist</i> , 2012, 193, 665-672.	3.5	312

#	ARTICLE	IF	CITATIONS
181	Variation in grain arsenic assessed in a diverse panel of rice ( <i>Oryza sativa</i> ) grown in multiple sites. <i>New Phytologist</i> , 2012, 193, 650-664.	3.5	126
182	Floral, climatic and soil pH controls on leaf ash content in China's terrestrial plants. <i>Global Ecology and Biogeography</i> , 2012, 21, 376-382.	2.7	36
183	Phytochelatin play a key role in arsenic accumulation and tolerance in the aquatic macrophyte <i>Wolffia globosa</i> . <i>Environmental Pollution</i> , 2012, 165, 18-24.	3.7	47
184	Localisation of iron in wheat grain using high resolution secondary ion mass spectrometry. <i>Journal of Cereal Science</i> , 2012, 55, 183-187.	1.8	59
185	Arsenic translocation in rice investigated using radioactive <sup>73</sup> As tracer. <i>Plant and Soil</i> , 2012, 350, 413-420.	1.8	66
186	Risk from Arsenic in Rice Grain. , 2012, , 31-50.		6
187	The Physiology of Arsenic in Rice. , 2012, , 103-138.		4
188	Strategies for Producing Low Arsenic Rice. , 2012, , 139-151.		3
189	Arsenic in Other Crops. , 2012, , 153-166.		0
190	Assessing the Labile Arsenic Pool in Contaminated Paddy Soils by Isotopic Dilution Techniques and Simple Extractions. <i>Environmental Science &amp; Technology</i> , 2011, 45, 4262-4269.	4.6	75
191	High-Resolution Secondary Ion Mass Spectrometry Reveals the Contrasting Subcellular Distribution of Arsenic and Silicon in Rice Roots. <i>Plant Physiology</i> , 2011, 156, 913-924.	2.3	122
192	Spatial distribution of arsenic and temporal variation of its concentration in rice. <i>New Phytologist</i> , 2011, 189, 200-209.	3.5	121
193	Evaluation of an electrostatic toxicity model for predicting Ni <sup>2+</sup> toxicity to barley root elongation in hydroponic cultures and in soils. <i>New Phytologist</i> , 2011, 192, 414-427.	3.5	23
194	Long-term impacts of zinc and copper enriched sewage sludge additions on bacterial, archaeal and fungal communities in arable and grassland soils. <i>Soil Biology and Biochemistry</i> , 2011, 43, 932-941.	4.2	65
195	Evidence for a role of phytochelatin in regulating arsenic accumulation in rice grain. <i>Environmental and Experimental Botany</i> , 2011, 71, 416-416.	2.0	75
196	The role of root hairs in cadmium acquisition by barley. <i>Environmental Pollution</i> , 2011, 159, 408-415.	3.7	38
197	The dynamics of arsenic in four paddy fields in the Bengal delta. <i>Environmental Pollution</i> , 2011, 159, 947-953.	3.7	95
198	Phytoremediation of arsenic contaminated paddy soils with <i>Pteris vittata</i> markedly reduces arsenic uptake by rice. <i>Environmental Pollution</i> , 2011, 159, 3739-3743.	3.7	98

#	ARTICLE	IF	CITATIONS
199	Selenium concentration and speciation in biofortified flour and bread: Retention of selenium during grain biofortification, processing and production of Se-enriched food. <i>Food Chemistry</i> , 2011, 126, 1771-1778.	4.2	110
200	Recent developments in modifying crops and agronomic practice to improve human health. <i>Food Policy</i> , 2011, 36, S94-S101.	2.8	64
201	Investigating the Contribution of the Phosphate Transport Pathway to Arsenic Accumulation in Rice $\hat{A}$ . <i>Plant Physiology</i> , 2011, 157, 498-508.	2.3	299
202	Development of a Real-Time PCR Assay for Detection and Quantification of <i>Rhizobium leguminosarum</i> Bacteria and Discrimination between Different Biovars in Zinc-Contaminated Soil. <i>Applied and Environmental Microbiology</i> , 2011, 77, 4626-4633.	1.4	24
203	The aromatic/arginine selectivity filter of NIP aquaporins plays a critical role in substrate selectivity for silicon, boron, and arsenic. <i>Journal of Experimental Botany</i> , 2011, 62, 4391-4398.	2.4	229
204	Arsenic as a Food Chain Contaminant: Mechanisms of Plant Uptake and Metabolism and Mitigation Strategies. <i>Annual Review of Plant Biology</i> , 2010, 61, 535-559.	8.6	1,023
205	Measurement of zinc stable isotope ratios in biogeochemical matrices by double-spike MC-ICPMS and determination of the isotope ratio pool available for plants from soil. <i>Analytical and Bioanalytical Chemistry</i> , 2010, 398, 3115-3125.	1.9	95
206	Rice is more efficient in arsenite uptake and translocation than wheat and barley. <i>Plant and Soil</i> , 2010, 328, 27-34.	1.8	277
207	Soil factors affecting selenium concentration in wheat grain and the fate and speciation of Se fertilisers applied to soil. <i>Plant and Soil</i> , 2010, 332, 19-30.	1.8	84
208	Impact of sulphur fertilisation on crop response to selenium fertilisation. <i>Plant and Soil</i> , 2010, 332, 31-40.	1.8	70
209	Selenium biofortification of high-yielding winter wheat ( <i>Triticum aestivum</i> L.) by liquid or granular Se fertilisation. <i>Plant and Soil</i> , 2010, 332, 5-18.	1.8	242
210	Relative impact of soil, metal source and metal concentration on bacterial community structure and community tolerance. <i>Soil Biology and Biochemistry</i> , 2010, 42, 1408-1417.	4.2	19
211	Predicting molybdenum toxicity to higher plants: Influence of soil properties. <i>Environmental Pollution</i> , 2010, 158, 3095-3102.	3.7	61
212	Predicting molybdenum toxicity to higher plants: Estimation of toxicity threshold values. <i>Environmental Pollution</i> , 2010, 158, 3085-3094.	3.7	60
213	The role of the rice aquaporin Lsi1 in arsenite efflux from roots. <i>New Phytologist</i> , 2010, 186, 392-399.	3.5	196
214	Evidence for the mechanisms of zinc uptake by rice using isotope fractionation. <i>Plant, Cell and Environment</i> , 2010, 33, 370-381.	2.8	107
215	Influence of Sulfur Deficiency on the Expression of Specific Sulfate Transporters and the Distribution of Sulfur, Selenium, and Molybdenum in Wheat. <i>Plant Physiology</i> , 2010, 153, 327-336.	2.3	151
216	Complexation of Arsenite with Phytochelatins Reduces Arsenite Efflux and Translocation from Roots to Shoots in <i>Arabidopsis</i> . <i>Plant Physiology</i> , 2010, 152, 2211-2221.	2.3	206

#	ARTICLE	IF	CITATIONS
217	Arsenic Speciation in Phloem and Xylem Exudates of Castor Bean. <i>Plant Physiology</i> , 2010, 154, 1505-1513.	2.3	104
218	Accumulation, Distribution, and Speciation of Arsenic in Wheat Grain. <i>Environmental Science &amp; Technology</i> , 2010, 44, 5464-5468.	4.6	86
219	Arsenic Bioavailability to Rice Is Elevated in Bangladeshi Paddy Soils. <i>Environmental Science &amp; Technology</i> , 2010, 44, 8515-8521.	4.6	139
220	Arsenic Influence on Genetic Variation in Grain Trace-Element Nutrient Content in Bengal Delta Grown Rice. <i>Environmental Science &amp; Technology</i> , 2010, 44, 8284-8288.	4.6	29
221	Arsenic Shoot-Grain Relationships in Field Grown Rice Cultivars. <i>Environmental Science &amp; Technology</i> , 2010, 44, 1471-1477.	4.6	54
222	Selenium Speciation in Soil and Rice: Influence of Water Management and Se Fertilization. <i>Journal of Agricultural and Food Chemistry</i> , 2010, 58, 11837-11843.	2.4	118
223	NanoSIMS analysis of arsenic and selenium in cereal grain. <i>New Phytologist</i> , 2010, 185, 434-445.	3.5	126
224	Distinguishing Diffusional and Plant Control of Cd and Ni Uptake by Hyperaccumulator and Nonhyperaccumulator Plants. <i>Environmental Science &amp; Technology</i> , 2010, 44, 6636-6641.	4.6	61
225	Biofortification and phytoremediation. <i>Current Opinion in Plant Biology</i> , 2009, 12, 373-380.	3.5	277
226	Variation in mineral micronutrient concentrations in grain of wheat lines of diverse origin. <i>Journal of Cereal Science</i> , 2009, 49, 290-295.	1.8	423
227	Sulphur nutrition differentially affects the distribution of asparagine in wheat grain. <i>Journal of Cereal Science</i> , 2009, 50, 407-409.	1.8	31
228	Physiological responses to Cd and Zn in two Cd/Zn hyperaccumulating <i>Thlaspi</i> species. <i>Environmental and Experimental Botany</i> , 2009, 66, 479-486.	2.0	54
229	Effect of nitrogen form on the rhizosphere dynamics and uptake of cadmium and zinc by the hyperaccumulator <i>Thlaspi caerulescens</i> . <i>Plant and Soil</i> , 2009, 318, 205-215.	1.8	131
230	Identification and functional analysis of two ZIP metal transporters of the hyperaccumulator <i>Thlaspi caerulescens</i> . <i>Plant and Soil</i> , 2009, 325, 79-95.	1.8	51
231	Arsenite transport in plants. <i>Cellular and Molecular Life Sciences</i> , 2009, 66, 2329-2339.	2.4	127
232	Arsenic uptake and metabolism in plants. <i>New Phytologist</i> , 2009, 181, 777-794.	3.5	973
233	Arsenic uptake and speciation in the rootless duckweed <i>Wolffia globosa</i> . <i>New Phytologist</i> , 2009, 182, 421-428.	3.5	111
234	Arsenite efflux is not enhanced in the arsenate-tolerant phenotype of <i>Holcus lanatus</i> . <i>New Phytologist</i> , 2009, 183, 340-348.	3.5	53

#	ARTICLE	IF	CITATIONS
235	Response to the Comment by Van Geen and Duxbury. <i>Environmental Science &amp; Technology</i> , 2009, 43, 3972-3973.	4.6	3
236	Phytotoxicity and bioavailability of cobalt to plants in a range of soils. <i>Chemosphere</i> , 2009, 75, 979-986.	4.2	127
237	Selenium in higher plants: understanding mechanisms for biofortification and phytoremediation. <i>Trends in Plant Science</i> , 2009, 14, 436-442.	4.3	486
238	Mitigation of Arsenic Accumulation in Rice with Water Management and Silicon Fertilization. <i>Environmental Science &amp; Technology</i> , 2009, 43, 3778-3783.	4.6	356
239	Environmental and Genetic Control of Arsenic Accumulation and Speciation in Rice Grain: Comparing a Range of Common Cultivars Grown in Contaminated Sites Across Bangladesh, China, and India. <i>Environmental Science &amp; Technology</i> , 2009, 43, 8381-8386.	4.6	146
240	Identification of Low Inorganic and Total Grain Arsenic Rice Cultivars from Bangladesh. <i>Environmental Science &amp; Technology</i> , 2009, 43, 6070-6075.	4.6	151
241	The Rice Aquaporin Lsi1 Mediates Uptake of Methylated Arsenic Species. <i>Plant Physiology</i> , 2009, 150, 2071-2080.	2.3	350
242	Geographical Variation in Total and Inorganic Arsenic Content of Polished (White) Rice. <i>Environmental Science &amp; Technology</i> , 2009, 43, 1612-1617.	4.6	673
243	METHODS FOR THE ANALYSIS OF SELENIUM AND OTHER MINERALS. , 2009, , 95-111.		0
244	Evidence of decreasing mineral density in wheat grain over the last 160 years. <i>Journal of Trace Elements in Medicine and Biology</i> , 2008, 22, 315-324.	1.5	373
245	Arsenate-induced toxicity: Effects on antioxidative enzymes and DNA damage in <i>Vicia faba</i> . <i>Environmental Toxicology and Chemistry</i> , 2008, 27, 413-419.	2.2	86
246	Selenium uptake, translocation and speciation in wheat supplied with selenate or selenite. <i>New Phytologist</i> , 2008, 178, 92-102.	3.5	593
247	Variation in root-to-shoot translocation of cadmium and zinc among different accessions of the hyperaccumulators <i>Thlaspi caerulescens</i> and <i>Thlaspi praecox</i> . <i>New Phytologist</i> , 2008, 178, 315-325.	3.5	90
248	Highly efficient xylem transport of arsenite in the arsenic hyperaccumulator <i>Pteris vittata</i> . <i>New Phytologist</i> , 2008, 180, 434-441.	3.5	161
249	Historical changes in the concentrations of selenium in soil and wheat grain from the Broadbalk experiment over the last 160 years. <i>Science of the Total Environment</i> , 2008, 389, 532-538.	3.9	44
250	Growing Rice Aerobically Markedly Decreases Arsenic Accumulation. <i>Environmental Science &amp; Technology</i> , 2008, 42, 5574-5579.	4.6	567
251	Atmospheric SO <sub>2</sub> Emissions Since the Late 1800s Change Organic Sulfur Forms in Humic Substance Extracts of Soils. <i>Environmental Science &amp; Technology</i> , 2008, 42, 3550-3555.	4.6	44
252	Arsenic accumulation by the aquatic fern <i>Azolla</i> : Comparison of arsenate uptake, speciation and efflux by <i>A. caroliniana</i> and <i>A. filiculoides</i> . <i>Environmental Pollution</i> , 2008, 156, 1149-1155.	3.7	89



#	ARTICLE	IF	CITATIONS
253	Use of Co speciation and soil properties to explain variation in Co toxicity to root growth of barley ( <i>Hordeum vulgare</i> L.) in different soils. <i>Environmental Pollution</i> , 2008, 156, 883-890.	3.7	43
254	Feasibility of phytoextraction to remediate cadmium and zinc contaminated soils. <i>Environmental Pollution</i> , 2008, 156, 905-914.	3.7	91
255	Does cadmium play a physiological role in the hyperaccumulator <i>Thlaspi caerulescens</i> ?. <i>Chemosphere</i> , 2008, 71, 1276-1283.	4.2	84
256	Transporters of arsenite in rice and their role in arsenic accumulation in rice grain. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2008, 105, 9931-9935.	3.3	1,202
257	Characterization of Cd Translocation and Identification of the Cd Form in Xylem Sap of the Cd-Hyperaccumulator <i>Arabidopsis halleri</i> . <i>Plant and Cell Physiology</i> , 2008, 49, 540-548.	1.5	157
258	Role of Sulfur for Plant Production in Agricultural and Natural Ecosystems. <i>Advances in Photosynthesis and Respiration</i> , 2008, , 417-435.	1.0	27
259	Mutation in Nicotianamine Aminotransferase Stimulated the Fe(II) Acquisition System and Led to Iron Accumulation in Rice. <i>Plant Physiology</i> , 2007, 145, 1647-1657.	2.3	167
260	Expression and functional analysis of metal transporter genes in two contrasting ecotypes of the hyperaccumulator <i>Thlaspi caerulescens</i> . <i>Journal of Experimental Botany</i> , 2007, 58, 1717-1728.	2.4	119
261	Defect Identification in Wheat Grain by Micro-Fluidic Electrophoresis: Sulfur Deficiency and Bug Damage. <i>Cereal Chemistry</i> , 2007, 84, 301-303.	1.1	9
262	Phytotoxicity of nickel in a range of European soils: Influence of soil properties, Ni solubility and speciation. <i>Environmental Pollution</i> , 2007, 145, 596-605.	3.7	150
263	Role of salicylic acid in alleviating oxidative damage in rice roots ( <i>Oryza sativa</i> ) subjected to cadmium stress. <i>Environmental Pollution</i> , 2007, 147, 743-749.	3.7	212
264	Estimates of ambient background concentrations of trace metals in soils for risk assessment. <i>Environmental Pollution</i> , 2007, 148, 221-229.	3.7	80
265	Effect of liming and seasonal variation on lead concentration of tea plant ( <i>Camellia sinensis</i> (L.) O.) Tj ETQq1 1 0.784314 rgBT /Overl 4.2 34	4.2	34
266	Critical Loads of Metals and Other Trace Elements to Terrestrial Environments. <i>Environmental Science &amp; Technology</i> , 2007, 41, 6326-6331.	4.6	35
267	Rapid reduction of arsenate in the medium mediated by plant roots. <i>New Phytologist</i> , 2007, 176, 590-599.	3.5	340
268	Strategies for increasing the selenium content of wheat. <i>Journal of Cereal Science</i> , 2007, 46, 282-292.	1.8	196
269	Effects of soil compaction and irrigation on the concentrations of selenium and arsenic in wheat grains. <i>Science of the Total Environment</i> , 2007, 372, 433-439.	3.9	62
270	Terrestrial Biotic Ligand Model. 2. Application to Ni and Cu Toxicities to Plants, Invertebrates, and Microbes in Soil. <i>Environmental Science &amp; Technology</i> , 2006, 40, 7094-7100.	4.6	164

#	ARTICLE	IF	CITATIONS
271	A Terrestrial Biotic Ligand Model. 1. Development and Application to Cu and Ni Toxicities to Barley Root Elongation in Soils. <i>Environmental Science &amp; Technology</i> , 2006, 40, 7085-7093.	4.6	224
272	Effect of soil characteristics on Cd uptake by the hyperaccumulator <i>Thlaspi caerulescens</i> . <i>Environmental Pollution</i> , 2006, 139, 167-175.	3.7	127
273	Scale and causes of lead contamination in Chinese tea. <i>Environmental Pollution</i> , 2006, 139, 125-132.	3.7	102
274	Field evaluation of Cd and Zn phytoextraction potential by the hyperaccumulators <i>Thlaspi caerulescens</i> and <i>Arabidopsis halleri</i> . <i>Environmental Pollution</i> , 2006, 141, 115-125.	3.7	268
275	Field evaluation of in situ remediation of a heavy metal contaminated soil using lime and red-mud. <i>Environmental Pollution</i> , 2006, 142, 530-539.	3.7	365
276	Cadmium uptake, translocation and tolerance in the hyperaccumulator <i>Arabidopsis halleri</i> . <i>New Phytologist</i> , 2006, 172, 646-654.	3.5	212
277	Synthesis of low molecular weight thiols in response to Cd exposure in <i>Thlaspi caerulescens</i> . <i>Plant, Cell and Environment</i> , 2006, 29, 1422-1429.	2.8	62
278	SOIL FACTORS CONTROLLING THE EXPRESSION OF COPPER TOXICITY TO PLANTS IN A WIDE RANGE OF EUROPEAN SOILS. <i>Environmental Toxicology and Chemistry</i> , 2006, 25, 726.	2.2	159
279	COMPARISON OF SOIL SOLUTION SPECIATION AND DIFFUSIVE GRADIENTS IN THIN-FILMS MEASUREMENT AS AN INDICATOR OF COPPER BIOAVAILABILITY TO PLANTS. <i>Environmental Toxicology and Chemistry</i> , 2006, 25, 733.	2.2	83
280	INFLUENCE OF SOIL PROPERTIES AND AGING ON ARSENIC PHYTOTOXICITY. <i>Environmental Toxicology and Chemistry</i> , 2006, 25, 1663.	2.2	80
281	Sulphur speciation and turnover in soils: evidence from sulphur K-edge XANES spectroscopy and isotope dilution studies. <i>Soil Biology and Biochemistry</i> , 2006, 38, 1000-1007.	4.2	75
282	Effects of sulphur on yield and malting quality of barley. <i>Journal of Cereal Science</i> , 2006, 43, 369-377.	1.8	45
283	Biofortification of UK food crops with selenium. <i>Proceedings of the Nutrition Society</i> , 2006, 65, 169-181.	0.4	378
284	Comparison of root absorption, translocation and tolerance of arsenic in the hyperaccumulator <i>Pteris vittata</i> and the nonhyperaccumulator <i>Pteris tremula</i> . <i>New Phytologist</i> , 2005, 165, 755-761.	3.5	92
285	Isotopic discrimination of zinc in higher plants. <i>New Phytologist</i> , 2005, 165, 703-710.	3.5	219
286	Cadmium hyperaccumulation protects <i>Thlaspi caerulescens</i> from leaf feeding damage by thrips ( <i>Tj ETQq0 0 0 rgBT, Overlock, 10 Tf 50</i> )	3.5	84
287	Subcellular localisation of Cd and Zn in the leaves of a Cd-hyperaccumulating ecotype of <i>Thlaspi caerulescens</i> . <i>Planta</i> , 2005, 220, 731-736.	1.6	217
288	Identification of the form of Cd in the leaves of a superior Cd-accumulating ecotype of <i>Thlaspi caerulescens</i> using <sup>113</sup> Cd-NMR. <i>Planta</i> , 2005, 221, 928-936.	1.6	95

#	ARTICLE	IF	CITATIONS
289	Sulphur fractionation in calcareous soils and bioavailability to plants. <i>Plant and Soil</i> , 2005, 268, 103-109.	1.8	49
290	Speciation and solubility of Cu, Ni and Pb in contaminated soils. <i>European Journal of Soil Science</i> , 2004, 55, 579-590.	1.8	57
291	Research Priorities for Conservation of Metallophyte Biodiversity and their Potential for Restoration and Site Remediation. <i>Restoration Ecology</i> , 2004, 12, 106-116.	1.4	304
292	POLLUTION-INDUCED COMMUNITY TOLERANCE OF SOIL MICROBES IN RESPONSE TO A ZINC GRADIENT. <i>Environmental Toxicology and Chemistry</i> , 2004, 23, 2665.	2.2	43
293	Cadmium and zinc accumulation by the hyperaccumulator <i>Thlaspi caerulescens</i> from soils enriched with insoluble metal compounds. <i>Soil Science and Plant Nutrition</i> , 2004, 50, 511-515.	0.8	13
294	Copper uptake by <i>Elsholtzia splendens</i> and <i>Silene vulgaris</i> and assessment of copper phytoavailability in contaminated soils. <i>Environmental Pollution</i> , 2004, 128, 307-315.	3.7	175
295	Factors affecting the concentrations of lead in British wheat and barley grain. <i>Environmental Pollution</i> , 2004, 131, 461-468.	3.7	59
296	Arsenic hyperaccumulation by <i>Pteris vittata</i> from arsenic contaminated soils and the effect of liming and phosphate fertilisation. <i>Environmental Pollution</i> , 2004, 132, 113-120.	3.7	98
297	A new method for the determination of the $^{34}\text{S}/^{32}\text{S}$ ratio of water-soluble sulphur in soil. <i>International Journal of Environmental Analytical Chemistry</i> , 2004, 84, 323-329.	1.8	2
298	Co-segregation analysis of cadmium and zinc accumulation in <i>Thlaspi caerulescens</i> interecotypic crosses. <i>New Phytologist</i> , 2004, 163, 299-312.	3.5	65
299	Predicting Cadmium Concentrations in Wheat and Barley Grain Using Soil Properties. <i>Journal of Environmental Quality</i> , 2004, 33, 532-541.	1.0	218
300	Predicting Cadmium Concentrations in Wheat and Barley Grain Using Soil Properties. <i>Journal of Environmental Quality</i> , 2004, 33, 532.	1.0	62
301	Determination of the Distribution of Sulphur in Wheat Starchy Endosperm Cells Using Secondary Ion Mass Spectroscopy (SIMS) Combined with Isotope Enhancement. <i>Journal of Cereal Science</i> , 2003, 37, 311-318.	1.8	12
302	Title is missing!. <i>Plant and Soil</i> , 2003, 249, 37-43.	1.8	370
303	Phytoextraction of metals and metalloids from contaminated soils. <i>Current Opinion in Biotechnology</i> , 2003, 14, 277-282.	3.3	908
304	Uptake and distribution of nickel and other metals in the hyperaccumulator <i>Berkheya coddii</i> . <i>New Phytologist</i> , 2003, 158, 279-285.	3.5	135
305	The role of phytochelatins in arsenic tolerance in the hyperaccumulator <i>Pteris vittata</i> . <i>New Phytologist</i> , 2003, 159, 403-410.	3.5	231
306	Stable Sulfur Isotope Ratio Indicates Long-Term Changes in Sulfur Deposition in the Broadbalk Experiment since 1845. <i>Journal of Environmental Quality</i> , 2003, 32, 33-39.	1.0	44

#	ARTICLE	IF	CITATIONS
307	Stable Sulfur Isotope Ratio Indicates Long-Term Changes in Sulfur Deposition in the Broadbalk Experiment since 1845. <i>Journal of Environmental Quality</i> , 2003, 32, 33.	1.0	18
308	Influence of Iron Status on Cadmium and Zinc Uptake by Different Ecotypes of the Hyperaccumulator <i>Thlaspi caerulescens</i> . <i>Plant Physiology</i> , 2002, 128, 1359-1367.	2.3	293
309	Characteristics of cadmium uptake in two contrasting ecotypes of the hyperaccumulator <i>Thlaspi caerulescens</i> . <i>Journal of Experimental Botany</i> , 2002, 53, 535-543.	2.4	328
310	Mechanisms of Arsenic Hyperaccumulation in <i>Pteris vittata</i> . Uptake Kinetics, Interactions with Phosphate, and Arsenic Speciation. <i>Plant Physiology</i> , 2002, 130, 1552-1561.	2.3	548
311	In situ fixation of metals in soils using bauxite residue: chemical assessment. <i>Environmental Pollution</i> , 2002, 118, 435-443.	3.7	297
312	In situ fixation of metals in soils using bauxite residue: biological effects. <i>Environmental Pollution</i> , 2002, 118, 445-452.	3.7	143
313	Evidence of low selenium concentrations in UK bread-making wheat grain. <i>Journal of the Science of Food and Agriculture</i> , 2002, 82, 1160-1165.	1.7	106
314	Arsenic hyperaccumulation by different fern species. <i>New Phytologist</i> , 2002, 156, 27-31.	3.5	361
315	Arsenic distribution and speciation in the fronds of the hyperaccumulator <i>Pteris vittata</i> . <i>New Phytologist</i> , 2002, 156, 195-203.	3.5	285
316	Plant and rhizosphere processes involved in phytoremediation of metal-contaminated soils. , 2002, , 207-214.		16
317	Leaching of heavy metals from contaminated soils using EDTA. <i>Environmental Pollution</i> , 2001, 113, 111-120.	3.7	429
318	Long-term effects of land use and fertiliser treatments on sulphur transformations in soils from the Broadbalk experiment. <i>Soil Biology and Biochemistry</i> , 2001, 33, 1797-1804.	4.2	56
319	USE OF THE ENRICHED STABLE ISOTOPE <sup>34</sup> S TO STUDY SULPHUR UPTAKE AND DISTRIBUTION IN WHEAT. <i>Journal of Plant Nutrition</i> , 2001, 24, 1551-1560.	0.9	20
320	A New Method to Measure Effective Soil Solution Concentration Predicts Copper Availability to Plants. <i>Environmental Science &amp; Technology</i> , 2001, 35, 2602-2607.	4.6	435
321	Phytoremediation of Heavy Metal-Contaminated Soils: Natural Hyperaccumulation versus Chemically Enhanced Phytoextraction. <i>Journal of Environmental Quality</i> , 2001, 30, 1919-1926.	1.0	493
322	Trends in <sup>13</sup> C/ <sup>12</sup> C ratios and C isotope discrimination of wheat since 1845. <i>Oecologia</i> , 2001, 128, 336-342.	0.9	45
323	Physiological evidence for a high-affinity cadmium transporter highly expressed in a <i>Thlaspi caerulescens</i> ecotype. <i>New Phytologist</i> , 2001, 149, 53-60.	3.5	254
324	What's new about cadmium hyperaccumulation?. <i>New Phytologist</i> , 2001, 149, 2-3.	3.5	35

#	ARTICLE	IF	CITATIONS
325	Root exudates of the hyperaccumulator <i>Thlaspi caerulescens</i> do not enhance metal mobilization. <i>New Phytologist</i> , 2001, 151, 613-620.	3.5	174
326	Plant and rhizosphere processes involved in phytoremediation of metal-contaminated soils. <i>Plant and Soil</i> , 2001, 232, 207-214.	1.8	455
327	Using plant analysis to predict yield losses caused by sulphur deficiency. <i>Annals of Applied Biology</i> , 2001, 138, 123-127.	1.3	31
328	Cellular compartmentation of nickel in the hyperaccumulators <i>Alyssum lesbiacum</i> , <i>Alyssum bertolonii</i> and <i>Thlaspi goesingense</i> . <i>Journal of Experimental Botany</i> , 2001, 52, 2291-2300.	2.4	317
329	INTER-LABORATORY COMPARISON OF SULPHUR AND NITROGEN ANALYSIS IN PLANTS AND SOILS. <i>Communications in Soil Science and Plant Analysis</i> , 2001, 32, 685-695.	0.6	16
330	Long-Term Changes in the Extractability and Bioavailability of Zinc and Cadmium after Sludge Application. <i>Journal of Environmental Quality</i> , 2000, 29, 875-883.	1.0	129
331	Cadmium accumulation in populations of <i>Thlaspi caerulescens</i> and <i>Thlaspi goesingense</i> . <i>New Phytologist</i> , 2000, 145, 11-20.	3.5	354
332	Zinc hyperaccumulation and cellular distribution in <i>Arabidopsis halleri</i> . <i>Plant, Cell and Environment</i> , 2000, 23, 507-514.	2.8	307
333	Title is missing!. <i>Plant and Soil</i> , 2000, 225, 95-107.	1.8	90
334	Cellular compartmentation of cadmium and zinc in relation to other elements in the hyperaccumulator <i>Arabidopsis halleri</i> . <i>Planta</i> , 2000, 212, 75-84.	1.6	618
335	Long-Term Effects of Land Use and Fertilizer Treatments on Sulfur Cycling. <i>Journal of Environmental Quality</i> , 2000, 29, 1867-1874.	1.0	55
336	Copper Speciation and Impacts on Bacterial Biosensors in the Pore Water of Copper-Contaminated Soils. <i>Environmental Science &amp; Technology</i> , 2000, 34, 5115-5121.	4.6	150
337	Sulphur accumulation and redistribution in wheat ( <i>Triticum aestivum</i> ): a study using stable sulphur isotope ratios as a tracer system. <i>Plant, Cell and Environment</i> , 1999, 22, 831-839.	2.8	56
338	Sulphur Assimilation and Effects on Yield and Quality of Wheat. <i>Journal of Cereal Science</i> , 1999, 30, 1-17.	1.8	330
339	Variation in the Breadmaking Quality and Rheological Properties of Wheat in Relation to Sulphur Nutrition under Field Conditions. <i>Journal of Cereal Science</i> , 1999, 30, 19-31.	1.8	93
340	Effects of sulphur nutrition on growth and nitrogen fixation of pea ( <i>Pisum sativum</i> L.). <i>Plant and Soil</i> , 1999, 212, 207-217.	1.8	55
341	Responses of breadmaking quality to sulphur in three wheat varieties. <i>Journal of the Science of Food and Agriculture</i> , 1999, 79, 1865-1874.	1.7	48
342	Solubility of zinc and interactions between zinc and phosphorus in the hyperaccumulator <i>Thlaspi caerulescens</i> . <i>Plant, Cell and Environment</i> , 1998, 21, 108-114.	2.8	51

#	ARTICLE	IF	CITATIONS
343	Thiosulphate and tetrathionate oxidation in arable soils. <i>Soil Biology and Biochemistry</i> , 1998, 30, 553-559.	4.2	25
344	Use of Sulfur Isotope Ratios To Determine Anthropogenic Sulfur Signals in a Grassland Ecosystem. <i>Environmental Science &amp; Technology</i> , 1998, 32, 2288-2291.	4.6	45
345	Distribution of Sulfur within Oilseed Rape Leaves in Response to Sulfur Deficiency during Vegetative Growth. <i>Plant Physiology</i> , 1998, 118, 1337-1344.	2.3	167
346	Nitrogen to sulphur ratio in rapeseed and in rapeseed protein and its use in diagnosing sulphur deficiency. <i>Journal of Plant Nutrition</i> , 1997, 20, 549-558.	0.9	48
347	Sulphur nutrition: An important factor for the quality of wheat and rapeseed. <i>Soil Science and Plant Nutrition</i> , 1997, 43, 1137-1142.	0.8	47
348	Uptake and transport of zinc in the hyperaccumulator <i>Thlaspi caerulescens</i> and the non-hyperaccumulator <i>Thlaspi ochroleucum</i> . <i>Plant, Cell and Environment</i> , 1997, 20, 898-906.	2.8	257
349	Title is missing!. <i>Plant and Soil</i> , 1997, 188, 153-159.	1.8	259
350	Title is missing!. <i>Plant and Soil</i> , 1997, 197, 71-78.	1.8	210
351	Sulphur nutrition: An important factor for the quality of wheat and rapeseed. , 1997, , 917-922.		8
352	Method to determine elemental sulphur in soils applied to measure sulphur oxidation. <i>Soil Biology and Biochemistry</i> , 1996, 28, 1083-1087.	4.2	18
353	Sulphur uptake, yield responses and the interactions between nitrogen and sulphur in winter oilseed rape ( <i>Brassica napus</i> ). <i>Journal of Agricultural Science</i> , 1996, 126, 53-62.	0.6	174
354	Responses of two wheat varieties to sulphur addition and diagnosis of sulphur deficiency. <i>Plant and Soil</i> , 1996, 181, 317-327.	1.8	80
355	Changes in the sulphur status of British wheat grain in the last decade, and its geographical distribution. <i>Journal of the Science of Food and Agriculture</i> , 1995, 68, 507-514.	1.7	37
356	A risk assessment of sulphur deficiency in cereals using soil and atmospheric deposition data. <i>Soil Use and Management</i> , 1995, 11, 110-114.	2.6	87
357	The Cadmium Content of British Wheat Grain. <i>Journal of Environmental Quality</i> , 1995, 24, 850-855.	1.0	45
358	Comparison of <i>aqua regia</i> digestion with sodium carbonate fusion for the determination of total phosphorus in soils by inductively coupled plasma atomic emission spectroscopy (ICP). <i>Communications in Soil Science and Plant Analysis</i> , 1995, 26, 1357-1368.	0.6	84
359	Extractable sulphate and organic sulphur in soils and their availability to plants. <i>Plant and Soil</i> , 1994, 164, 243-250.	1.8	126
360	Influence of nitrogen and sulphur on the glucosinolate profile of rapeseed ( <i>brassica napus</i> l). <i>Journal of the Science of Food and Agriculture</i> , 1994, 64, 295-304.	1.7	95

#	ARTICLE	IF	CITATIONS
361	Comparison of three wet digestion methods for the determination of plant sulphur by inductively coupled plasma atomic emission spectroscopy (ICP-AES). <i>Communications in Soil Science and Plant Analysis</i> , 1994, 25, 407-418.	0.6	288
362	Comparison of sulphur uptake by oilseed rape and the soil sulphur status of two adjacent fields with different soil series. <i>Soil Use and Management</i> , 1994, 10, 47-50.	2.6	11
363	Sulphur turnover in the developing pods of single and double low varieties of oilseed rape ( <i>Brassica</i> ) Tj ETQq1 1 0.784314 rgBT /Overl 1.7	1.7	24
364	Influence of sulphur and nitrogen on seed yield and quality of low glucosinolate oilseed rape ( <i>Brassica napus</i> L). <i>Journal of the Science of Food and Agriculture</i> , 1993, 63, 29-37.	1.7	138
365	Sulphur uptake and distribution in double and single low varieties of oilseed rape ( <i>Brassica napus</i> L.). <i>Plant and Soil</i> , 1993, 150, 69-76.	1.8	40
366	Glucosinolate changes in developing pods of single and double low varieties of autumn-sown oilseed rape ( <i>B. napus</i> ). <i>Annals of Applied Biology</i> , 1993, 122, 135-143.	1.3	17
367	The influence of spring nitrogen on yield, yield components and glucosinolate content of autumn-sown oilseed rape ( <i>Brassica napus</i> ). <i>Journal of Agricultural Science</i> , 1993, 120, 219-224.	0.6	43
368	Correction for protein content in the determination of the glucosinolate content of rapeseed by the X-RF method. <i>Journal of the Science of Food and Agriculture</i> , 1992, 58, 431-433.	1.7	22