Shinichi Sato

List of Publications by Year in Descending Order

Source: https://exaly.com/author-pdf/248271/shinichi-sato-publications-by-year.pdf

Version: 2024-04-10

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

210
papers

6,085
h-index

71
g-index

725
ext. papers

7,124
ext. citations

40
h-index

5.47
avg, IF

L-index

#	Paper	IF	Citations
210	Decrease in MAP3Ks expression enhances the cell death caused by hyperthermia <i>International Journal of Hyperthermia</i> , 2022 , 39, 200-208	3.7	1
209	Predictors of Rituximab Effect on Modified Rodnan Skin Score in Systemic Sclerosis: a machine learning analysis of the DESIRES trial <i>Rheumatology</i> , 2022 ,	3.9	1
208	Discordant lymphomas of classic Hodgkin lymphoma and peripheral T-cell lymphoma following dupilumab treatment for atopic dermatitis <i>International Journal of Hematology</i> , 2022 , 1	2.3	O
207	Serum C-X-C Chemokine Ligand 1 Levels in Patients with Systemic Sclerosis: Relationship of Clinical and Laboratory Observations to Anti-CD20 Monoclonal Antibody Administration. <i>Life</i> , 2022 , 12, 646	3	1
206	Association of serum CXCL12 levels with arthropathy in patients with systemic sclerosis. <i>International Journal of Rheumatic Diseases</i> , 2021 , 24, 260-267	2.3	1
205	Endothelial CCR6 expression due to FLI1 deficiency contributes to vasculopathy associated with systemic sclerosis. <i>Arthritis Research and Therapy</i> , 2021 , 23, 283	5.7	1
204	The contribution of LIGHT to the development of systemic sclerosis by modulating IL-6 and Th1 chemokine expression in dermal fibroblasts. <i>Journal of Investigative Dermatology</i> , 2021 ,	4.3	1
203	Interleukin-31 promotes fibrosis and T helper 2 polarization in systemic sclerosis. <i>Nature Communications</i> , 2021 , 12, 5947	17.4	3
202	HMGB1-mediated chromatin remodeling attenuates gene expression for the protection from allergic contact dermatitis. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2021 , 118,	11.5	3
201	Development of a prediction model of treatment response in patients with cutaneous arteritis: Insights from a cohort of 33 patients. <i>Journal of Dermatology</i> , 2021 , 48, 1021-1026	1.6	О
2 00	Association of serum CCL20 levels with pulmonary vascular involvement and primary biliary cholangitis in patients with systemic sclerosis. <i>International Journal of Rheumatic Diseases</i> , 2021 , 24, 711	1 - 738	4
199	Rapidly progressive interstitial lung disease associated with dermatomyositis Longitudinal course of anti-MDA5 antibody titer in two cases. <i>Journal of Cutaneous Immunology and Allergy</i> , 2021 , 4, 78-82	0.3	
198	Serum vasohibin-1 levels: A potential marker of dermal and pulmonary fibrosis in systemic sclerosis. <i>Experimental Dermatology</i> , 2021 , 30, 951-958	4	1
197	Fli1 deficiency suppresses RALDH1 activity of dermal dendritic cells and related induction of regulatory T cells: a possible role in scleroderma. <i>Arthritis Research and Therapy</i> , 2021 , 23, 137	5.7	2
196	Immune checkpoint inhibitor combination therapies very frequently induce secondary adrenal insufficiency. <i>Scientific Reports</i> , 2021 , 11, 11617	4.9	1
195	Dermatomyositis-like eruptions and fasciitis with novel compound heterozygous MEFV mutations: Newly recognized features of a variant of familial Mediterranean fever. <i>Journal of Dermatology</i> , 2021 , 48, 1453-1456	1.6	
194	Safety and efficacy of rituximab in systemic sclerosis (DESIRES): a double-blind, investigator-initiated, randomised, placebo-controlled trial. <i>Lancet Rheumatology, The</i> , 2021 , 3, e489-e4	9 7 .2	20

(2020-2021)

193	Serum S100A12 levels: Possible association with skin sclerosis and interstitial lung disease in systemic sclerosis. <i>Experimental Dermatology</i> , 2021 , 30, 409-415	4	3
192	Fibrosarcomatous dermatofibrosarcoma protuberans in a seven-year-old boy. <i>European Journal of Dermatology</i> , 2021 , 31, 106-107	0.8	
191	A potential contribution of decreased serum galectin-10 levels to systemic inflammation and pulmonary vascular involvement in systemic sclerosis. <i>Experimental Dermatology</i> , 2021 , 30, 959-965	4	1
190	CD147-Cyclophilin a Interactions Promote Proliferation and Survival of Cutaneous T-Cell Lymphoma. <i>International Journal of Molecular Sciences</i> , 2021 , 22,	6.3	5
189	B Cell Depletion Inhibits Fibrosis via Suppression of Profibrotic Macrophage Differentiation in a Mouse Model of Systemic Sclerosis. <i>Arthritis and Rheumatology</i> , 2021 , 73, 2086-2095	9.5	2
188	Expert-Level Distinction of Systemic Sclerosis from Hand Photographs Using Deep Convolutional Neural Networks. <i>Journal of Investigative Dermatology</i> , 2021 , 141, 2536-2539	4.3	Ο
187	Case of systemic sclerosis with multiple primary malignancies in whom anti-RNA polymerase III antibody was detected by immunoprecipitation. <i>Journal of Dermatology</i> , 2020 , 47, e269-e270	1.6	Ο
186	Rapid decrease of serum surfactant protein-D levels predicts the reactivity of rituximab therapy in systemic sclerosis-associated interstitial lung disease. <i>Journal of Dermatology</i> , 2020 , 47, 796-800	1.6	2
185	Decreased serum cathepsin S levels in patients with systemic sclerosis-associated interstitial lung disease. <i>Journal of Dermatology</i> , 2020 , 47, 1027-1032	1.6	2
184	Altered Properties of Endothelial Cells and Mesenchymal Stem Cells Underlying the Development of Scleroderma-like Vasculopathy in KLF5 ;Fli-1 Mice. <i>Arthritis and Rheumatology</i> , 2020 , 72, 2136-2146	9.5	7
183	Combined immunosuppressive therapy provides favorable prognosis and increased risk of cytomegalovirus reactivation in anti-melanoma differentiation-associated gene 5 antibody-positive dermatomyositis. <i>Journal of Dermatology</i> , 2020 , 47, 483-489	1.6	8
182	Subacute thyroiditis in psoriasis patients treated with biologics targeting tumor necrosis factor- and interleukin-17A, a report of two cases. <i>Journal of Cutaneous Immunology and Allergy</i> , 2020 , 3, 33-34	0.3	2
181	Regulation of skin fibrosis by RALDH1-producing dermal dendritic cells via retinoic acid-mediated regulatory T cell induction: A role in scleroderma. <i>Journal of Dermatological Science</i> , 2020 , 97, 125-134	4.3	3
180	Clinical significance of endothelial vasodilatory function evaluated by EndoPAT in patients with systemic sclerosis. <i>Journal of Dermatology</i> , 2020 , 47, 609-614	1.6	3
179	A case of folliculotropic mycosis fungoides successfully treated with topical steroid treatment. Journal of Cancer Research and Therapeutics, 2020 , 16, 196-198	1.2	
178	Tumoral calcinosis in systemic lupus erythematosus associated with fat necrosis. <i>Journal of Dermatology</i> , 2020 , 47, e134-e135	1.6	1
177	Fli1 deficiency induces endothelial adipsin expression, contributing to the onset of pulmonary arterial hypertension in systemic sclerosis. <i>Rheumatology</i> , 2020 , 59, 2005-2015	3.9	3
176	Characteristics of Japanese patients with eosinophilic fasciitis: A brief multicenter study. <i>Journal of Dermatology</i> , 2020 , 47, 1391-1394	1.6	1

175	Association of functional (GA)n microsatellite polymorphism in the FLI1 gene with susceptibility to human systemic sclerosis. <i>Rheumatology</i> , 2020 , 59, 3553-3562	3.9	4
174	YKL-40 Promotes Proliferation of Cutaneous T-Cell Lymphoma Tumor Cells through Extracellular Signal-Regulated Kinase Pathways. <i>Journal of Investigative Dermatology</i> , 2020 , 140, 860-868.e3	4.3	5
173	Increased IL-26 Expression Promotes T Helper Type 17- and T Helper Type 2-Associated Cytokine Production by Keratinocytes in Atopic Dermatitis. <i>Journal of Investigative Dermatology</i> , 2020 , 140, 636	5-644.e2	. 17
172	Characteristics of Pulmonary Arterial Hypertension in Patients with Systemic Sclerosis and Anticentriole Autoantibodies. <i>International Heart Journal</i> , 2020 , 61, 413-418	1.8	4
171	Overlapping systemic sclerosis and sarcoidosis with mutually exclusive disease activities: a case report and analysis of previous studies. <i>European Journal of Dermatology</i> , 2020 , 30, 50-52	0.8	
170	Metabolome Analysis Reveals Dermal Histamine Accumulation in Murine Dermatitis Provoked by Genetic Deletion of P-Glycoprotein and Breast Cancer Resistance Protein. <i>Pharmaceutical Research</i> , 2019 , 36, 158	4.5	3
169	IL-10Broducing regulatory B cells in skin diseases. <i>Journal of Cutaneous Immunology and Allergy</i> , 2019 , 2, 68-74	0.3	2
168	Keratinocyte Proline-Rich Protein Deficiency în Atopic Dermatitis Leads to Barrier Disruption. <i>Journal of Investigative Dermatology</i> , 2019 , 139, 1867-1875.e7	4.3	5
167	Possible association of decreased serum CXCL14 levels with digital ulcers in patients with systemic sclerosis. <i>Journal of Dermatology</i> , 2019 , 46, 584-589	1.6	3
166	Facile fabrication of PEG-coated PLGA microspheres via SPG membrane emulsification for the treatment of scleroderma by ECM degrading enzymes. <i>Colloids and Surfaces B: Biointerfaces</i> , 2019 , 179, 453-461	6	4
165	Cyclophosphamide Pulse Therapy Normalizes Vascular Abnormalities in a Mouse Model of Systemic Sclerosis Vasculopathy. <i>Journal of Investigative Dermatology</i> , 2019 , 139, 1150-1160	4.3	1
164	A potential contribution of decreased galectin-7 expression in stratified epithelia to the development of cutaneous and oesophageal manifestations in systemic sclerosis. <i>Experimental Dermatology</i> , 2019 , 28, 536-542	4	13
163	Novel topical and systemic therapies in atopic dermatitis. <i>Immunological Medicine</i> , 2019 , 42, 84-93	3.7	15
162	Possible therapeutic applicability of galectin-9 in cutaneous T-cell lymphoma. <i>Journal of Dermatological Science</i> , 2019 , 96, 134-142	4.3	4
161	Association of NCF1 polymorphism with systemic lupus erythematosus and systemic sclerosis but not with ANCA-associated vasculitis in a Japanese population. <i>Scientific Reports</i> , 2019 , 9, 16366	4.9	9
160	Rituximab therapy is more effective than cyclophosphamide therapy for Japanese patients with anti-topoisomerase I-positive systemic sclerosis-associated interstitial lung disease. <i>Journal of Dermatology</i> , 2019 , 46, 1006-1013	1.6	24
159	A case of concurrent psoriasis and ulcerative colitis with development of arthritis during adalimumab treatment. <i>European Journal of Dermatology</i> , 2019 , 29, 651-653	0.8	О
158	A case of papuloerythroderma secondary to crusted scabies. <i>Journal of Cutaneous Immunology and Allergy</i> , 2019 , 2, 174-175	0.3	

(2018-2019)

157	Prevention of calpain-dependent degradation of STK38 by MEKK2-mediated phosphorylation. <i>Scientific Reports</i> , 2019 , 9, 16010	4.9	4	
156	Increased expression of aquaporin-1 in dermal fibroblasts and dermal microvascular endothelial cells possibly contributes to skin fibrosis and edema in patients with systemic sclerosis. <i>Journal of Dermatological Science</i> , 2019 , 93, 24-32	4.3	7	
155	Association between serum autotaxin or phosphatidylserine-specific phospholipase A1 levels and melanoma. <i>Journal of Dermatology</i> , 2018 , 45, 571-579	1.6	16	
154	Diagnostic criteria, severity classification and guidelines of localized scleroderma. <i>Journal of Dermatology</i> , 2018 , 45, 755-780	1.6	35	
153	Diagnostic criteria, severity classification and guidelines of systemic sclerosis. <i>Journal of Dermatology</i> , 2018 , 45, 633-691	1.6	16	
152	Serum Soluble CD48 Levels as a Prognostic Marker in Mycosis Fungoides and Sary Syndrome. Journal of Investigative Dermatology, 2018 , 138, 2286-2288	4.3	2	
151	Exacerbated Immune Complex-Mediated Vascular Injury in Mice with Heterozygous Deficiency of Aryl Hydrocarbon Receptor through Upregulation of Fc[Receptor III Expression on Macrophages. <i>Journal of Investigative Dermatology</i> , 2018 , 138, 2195-2204	4.3	1	
150	Impact of a new simplified disability scoring system for adult patients with localized scleroderma. <i>Journal of Dermatology</i> , 2018 , 45, 431-435	1.6	2	
149	Successful treatment with rituximab in a Japanese patient with systemic sclerosis-associated interstitial lung disease resistant to oral steroid and cyclophosphamide. <i>Journal of Dermatology</i> , 2018 , 45, e140-e141	1.6	2	
148	Systemic sclerosis complicated with localized scleroderma-like lesions induced by KBner phenomenon. <i>Journal of Dermatological Science</i> , 2018 , 89, 282-289	4.3	11	
147	Variants at HLA-A, HLA-C, and HLA-DQB1 Confer Risk of Psoriasis Vulgaris in Japanese. <i>Journal of Investigative Dermatology</i> , 2018 , 138, 542-548	4.3	29	
146	TBX4 is involved in the super-enhancer-driven transcriptional programs underlying features specific to lung fibroblasts. <i>American Journal of Physiology - Lung Cellular and Molecular Physiology</i> , 2018 , 314, L177-L191	5.8	26	
145	CXCL13 produced by macrophages due to Fli1 deficiency may contribute to the development of tissue fibrosis, vasculopathy and immune activation in systemic sclerosis. <i>Experimental Dermatology</i> , 2018 , 27, 1030-1037	4	27	
144	Rapid alteration of serum interleukin-6 levels may predict the reactivity of i.v. cyclophosphamide pulse therapy in systemic sclerosis-associated interstitial lung disease. <i>Journal of Dermatology</i> , 2018 , 45, 1221-1224	1.6	5	
143	Gastroesophageal Reflux Disease-Related Disorders of Systemic Sclerosis Based on the Analysis of 66 Patients. <i>Digestion</i> , 2018 , 98, 201-208	3.6	5	
142	Fli1-haploinsufficient dermal fibroblasts promote skin-localized transdifferentiation of Th2-like regulatory T cells. <i>Arthritis Research and Therapy</i> , 2018 , 20, 23	5.7	13	
141	An orally-active adiponectin receptor agonist mitigates cutaneous fibrosis, inflammation and microvascular pathology in a murine model of systemic sclerosis. <i>Scientific Reports</i> , 2018 , 8, 11843	4.9	24	
140	Diagnostic criteria, severity classification and guidelines of eosinophilic fasciitis. <i>Journal of Dermatology</i> , 2018 , 45, 881-890	1.6	21	

139	Decreased IL-10-producing regulatory B cells in patients with advanced mycosis fungoides. European Journal of Dermatology, 2018 , 28, 314-319	0.8	2
138	Prediction of therapeutic response before and during i.v. cyclophosphamide pulse therapy for interstitial lung disease in systemic sclerosis: A longitudinal observational study. <i>Journal of Dermatology</i> , 2018 , 45, 1425-1433	1.6	15
137	Progranulin overproduction due to constitutively activated c-Abl/PKC-IFli1 pathway contributes to the resistance of dermal fibroblasts to the anti-fibrotic effect of tumor necrosis factor-In localized scleroderma. <i>Journal of Dermatological Science</i> , 2018 , 92, 207-214	4.3	4
136	Aberrant CD137 ligand expression induced by GATA6 overexpression promotes tumor progression in cutaneous T-cell lymphoma. <i>Blood</i> , 2018 , 132, 1922-1935	2.2	23
135	Linear immunoglobulin A bullous dermatosis limited to oral mucosa associated with ulcerative colitis. <i>Journal of Dermatology</i> , 2018 , 45, e281-e282	1.6	0
134	Epithelial Fli1 deficiency drives systemic autoimmunity and fibrosis: Possible roles in scleroderma. Journal of Experimental Medicine, 2017 , 214, 1129-1151	16.6	58
133	CXCL17 Attenuates Imiquimod-Induced Psoriasis-like Skin Inflammation by Recruiting Myeloid-Derived Suppressor Cells and Regulatory T Cells. <i>Journal of Immunology</i> , 2017 , 198, 3897-3908	5.3	27
132	Serum H-ficolin levels: Clinical association with interstitial lung disease in patients with systemic sclerosis. <i>Journal of Dermatology</i> , 2017 , 44, 1168-1171	1.6	6
131	Fli1 Deficiency Induces CXCL6 Expression in Dermal Fibroblasts and Endothelial Cells, Contributing to the Development of Fibrosis and Vasculopathy in Systemic Sclerosis. <i>Journal of Rheumatology</i> , 2017 , 44, 1198-1205	4.1	15
130	Case of mycosis fungoides with gastric and central nervous system involvement. <i>Journal of Dermatology</i> , 2017 , 44, e166-e167	1.6	
129	Critical contribution of the interleukin-6/signal transducer and activator of transcription 3 axis to vasculopathy associated with systemic sclerosis. <i>Journal of Dermatology</i> , 2017 , 44, 967-971	1.6	12
128	Systemic Sclerosis Dermal Fibroblasts Suppress Th1 Cytokine Production via Galectin-9 Overproduction due to Fli1 Deficiency. <i>Journal of Investigative Dermatology</i> , 2017 , 137, 1850-1859	4.3	26
127	Contribution of Soluble Forms of Programmed Death 1 and Programmed Death Ligand 2 to Disease Severity and Progression in Systemic Sclerosis. <i>Arthritis and Rheumatology</i> , 2017 , 69, 1879-1890	9.5	28
126	The impact of transcription factor Fli1 deficiency on the regulation of angiogenesis. <i>Experimental Dermatology</i> , 2017 , 26, 912-918	4	16
125	Twelve-week, multicenter, placebo-controlled, randomized, double-blind, parallel-group, comparative phase II/III study of benzoyl peroxide gel in patients with acne vulgaris: A secondary publication. <i>Journal of Dermatology</i> , 2017 , 44, 774-782	1.6	6
124	Increased Interleukin-19 Expression in Cutaneous T-cell Lymphoma and Atopic Dermatitis. <i>Acta Dermato-Venereologica</i> , 2017 , 97, 1172-1177	2.2	15
123	Placental Growth Factor and Vascular Endothelial Growth Factor Together Regulate Tumour Progression via Increased Vasculature in Cutaneous T-cell Lymphoma. <i>Acta Dermato-Venereologica</i> , 2017 , 97, 586-592	2.2	12
122	A possible implication of reduced levels of LIF, LIFR, and gp130 in vasculopathy related to systemic sclerosis. <i>Archives of Dermatological Research</i> , 2017 , 309, 833-842	3.3	6

121	Glycyrrhizin Ameliorates Fibrosis, Vasculopathy, and Inflammation in Animal Models of Systemic Sclerosis. <i>Journal of Investigative Dermatology</i> , 2017 , 137, 631-640	4.3	21	
120	Plasma plasmin-2-plasmin inhibitor complex levels may predict the effect of cyclophosphamide for systemic sclerosis-related interstitial lung disease. <i>Modern Rheumatology</i> , 2017 , 27, 618-622	3.3	4	
119	Circulating galectin-1 concentrations in systemic sclerosis: potential contribution to digital vasculopathy. <i>International Journal of Rheumatic Diseases</i> , 2016 , 19, 622-7	2.3	12	
118	Thymic Stromal Chemokine TSLP Acts through Th2 Cytokine Production to Induce Cutaneous T-cell Lymphoma. <i>Cancer Research</i> , 2016 , 76, 6241-6252	10.1	57	
117	Hematopoietic stem cell transplantation for cutaneous T-cell lymphoma: Summary of 11 cases from two facilities in Japan and Brazil. <i>Journal of Dermatology</i> , 2016 , 43, 638-42	1.6	11	
116	Association of anti-RNA polymerase III antibody and silicone breast implants in patients with systemic sclerosis. <i>Journal of Dermatology</i> , 2016 , 43, 808-10	1.6	8	
115	A potential contribution of altered cathepsin L expression to the development of dermal fibrosis and vasculopathy in systemic sclerosis. <i>Experimental Dermatology</i> , 2016 , 25, 287-92	4	17	
114	Tamibarotene Ameliorates Bleomycin-Induced Dermal Fibrosis by Modulating Phenotypes of Fibroblasts, Endothelial Cells, and Immune Cells. <i>Journal of Investigative Dermatology</i> , 2016 , 136, 387-3	9 8 ·3	18	
113	Nucleosome in patients with systemic sclerosis: possible association with immunological abnormalities via abnormal activation of T and B cells. <i>Annals of the Rheumatic Diseases</i> , 2016 , 75, 1858	-65 ⁴	11	
112	Effect of ambrisentan on peripheral circulation in patients with systemic sclerosis. <i>Modern Rheumatology</i> , 2016 , 26, 454-7	3.3	1	
111	Human Leukocyte Antigen and Systemic Sclerosis in Japanese: The Sign of the Four Independent Protective Alleles, DRB1*13:02, DRB1*14:06, DQB1*03:01, and DPB1*02:01. <i>PLoS ONE</i> , 2016 , 11, e0154	1235	17	
110	Autoimmunity in Systemic Sclerosis: Overview 2016 , 21-37			
109	A case of penile basal cell carcinoma reconstructed by scrotal myofasciocutaneous flap. <i>Dermatologic Therapy</i> , 2016 , 29, 349-352	2.2	1	
108	Serum levels of interleukin-18-binding protein isoform a: Clinical association with inflammation and pulmonary hypertension in systemic sclerosis. <i>Journal of Dermatology</i> , 2016 , 43, 912-8	1.6	9	
107	Decreased interleukin-21 expression in skin and blood in advanced mycosis fungoides. <i>Journal of Dermatology</i> , 2016 , 43, 819-22	1.6	4	
106	Increased expression of chemerin in endothelial cells due to Fli1 deficiency may contribute to the development of digital ulcers in systemic sclerosis. <i>Rheumatology</i> , 2015 , 54, 1308-16	3.9	34	
105	Amelioration of tissue fibrosis by toll-like receptor 4 knockout in murine models of systemic sclerosis. <i>Arthritis and Rheumatology</i> , 2015 , 67, 254-65	9.5	49	
104	Elevated serum galectin-9 levels in patients with atopic dermatitis. <i>Journal of Dermatology</i> , 2015 , 42, 723-6	1.6	11	

103	Regulatory B cells in human inflammatory and autoimmune diseases: from mouse models to clinical research. <i>International Immunology</i> , 2015 , 27, 495-504	4.9	68
102	CCR4 is expressed on infiltrating cells in lesional skin of early mycosis fungoides and atopic dermatitis. <i>Journal of Dermatology</i> , 2015 , 42, 613-5	1.6	33
101	Endothelin receptor blockade ameliorates vascular fragility in endothelial cell-specific Fli-1-knockout mice by increasing Fli-1 DNA binding ability. <i>Arthritis and Rheumatology</i> , 2015 , 67, 1335-4	14 ^{9.5}	25
100	Vasculopathy in scleroderma. <i>Seminars in Immunopathology</i> , 2015 , 37, 489-500	12	105
99	Progranulin Overproduction Due to Fli-1 Deficiency Contributes to the Resistance of Dermal Fibroblasts to Tumor Necrosis Factor in Systemic Sclerosis. <i>Arthritis and Rheumatology</i> , 2015 , 67, 3245-5	5 3 ·5	24
98	Multifaceted contribution of the TLR4-activated IRF5 transcription factor in systemic sclerosis. Proceedings of the National Academy of Sciences of the United States of America, 2015, 112, 15136-41	11.5	39
97	Fibrosis, vascular activation, and immune abnormalities resembling systemic sclerosis in bleomycin-treated Fli-1-haploinsufficient mice. <i>Arthritis and Rheumatology</i> , 2015 , 67, 517-26	9.5	67
96	Abnormal B lymphocyte activation and function in systemic sclerosis. <i>Annals of Dermatology</i> , 2015 , 27, 1-9	0.4	28
95	Deficiency of both L-selectin and ICAM-1 exacerbates imiquimod-induced psoriasis-like skin inflammation through increased infiltration of antigen presenting cells. <i>Clinical Immunology</i> , 2015 , 157, 43-55	9	13
94	A possible contribution of endothelial CCN1 downregulation due to Fli1 deficiency to the development of digital ulcers in systemic sclerosis. <i>Experimental Dermatology</i> , 2015 , 24, 127-32	4	37
93	Lymphatic dysfunction attenuates tumor immunity through impaired antigen presentation. <i>Oncotarget</i> , 2015 , 6, 18081-93	3.3	30
92	TLR4, rather than TLR2, regulates wound healing through TGF-land CCL5 expression. <i>Journal of Dermatological Science</i> , 2014 , 73, 117-24	4.3	64
91	Skin barrier dysfunction and low antimicrobial peptide expression in cutaneous T-cell lymphoma. <i>Clinical Cancer Research</i> , 2014 , 20, 4339-48	12.9	22
90	Successful experience of rituximab therapy for systemic sclerosis-associated interstitial lung disease with concomitant systemic lupus erythematosus. <i>Journal of Dermatology</i> , 2014 , 41, 418-20	1.6	19
89	An association study of 36 psoriasis susceptibility loci for psoriasis vulgaris and atopic dermatitis in a Japanese population. <i>Journal of Dermatological Science</i> , 2014 , 76, 156-7	4.3	10
88	Effects of the immunosuppressant rapamycin on the expression of human 2 (I) collagen and matrix metalloproteinase 1 genes in scleroderma dermal fibroblasts. <i>Journal of Dermatological Science</i> , 2014 , 74, 251-9	4.3	22
87	Genetic polymorphism in the TRAF3IP2 gene is associated with psoriasis vulgaris in a Japanese population. <i>Journal of Dermatological Science</i> , 2014 , 73, 264-5	4.3	8
86	Serum autotaxin levels correlate with pruritus in patients with atopic dermatitis. <i>Journal of Investigative Dermatology</i> , 2014 , 134, 1745-1747	4.3	17

85	The role of IL-32 in cutaneous T-cell lymphoma. Journal of Investigative Dermatology, 2014, 134, 1428-1	4455	40
84	Serum levels of angiopoietin-2, but not angiopoietin-1, are elevated in patients with erythrodermic cutaneous T-cell lymphoma. <i>Acta Dermato-Venereologica</i> , 2014 , 94, 9-13	2.2	14
83	Clinical correlation of brachial artery flow-mediated dilation in patients with systemic sclerosis. <i>Modern Rheumatology</i> , 2014 , 24, 106-11	3.3	20
82	Simultaneous downregulation of KLF5 and Fli1 is a key feature underlying systemic sclerosis. <i>Nature Communications</i> , 2014 , 5, 5797	17.4	98
81	Serum levels of matrix metalloproteinase-13 in patients with eosinophilic fasciitis. <i>Journal of Dermatology</i> , 2014 , 41, 746-8	1.6	5
80	Novel IL36RN gene mutation revealed by analysis of 8 Japanese patients with generalized pustular psoriasis. <i>Journal of Dermatological Science</i> , 2014 , 76, 267-9	4.3	15
79	Cilostazol improves lymphatic function by inducing proliferation and stabilization of lymphatic endothelial cells. <i>Journal of Dermatological Science</i> , 2014 , 74, 150-8	4.3	15
78	Animal models of scleroderma: current state and recent development. <i>Current Rheumatology Reports</i> , 2013 , 15, 382	4.9	23
77	Increased production of soluble inducible costimulator in patients with diffuse cutaneous systemic sclerosis. <i>Archives of Dermatological Research</i> , 2013 , 305, 17-23	3.3	13
76	ICAM-1 deficiency exacerbates sarcoid-like granulomatosis induced by Propionibacterium acnes through impaired IL-10 production by regulatory T cells. <i>American Journal of Pathology</i> , 2013 , 183, 173	1- 1 739	9
75	Regulatory B cells suppress imiquimod-induced, psoriasis-like skin inflammation. <i>Journal of Leukocyte Biology</i> , 2013 , 94, 563-73	6.5	71
74	CXCR4 negatively regulates keratinocyte proliferation in IL-23-mediated psoriasiform dermatitis. <i>Journal of Investigative Dermatology</i> , 2013 , 133, 2530-2537	4.3	15
74 73		4·3 3·9	15 34
	Journal of Investigative Dermatology, 2013, 133, 2530-2537 Decreased cathepsin V expression due to Fli1 deficiency contributes to the development of dermal		
73	Decreased cathepsin V expression due to Fli1 deficiency contributes to the development of dermal fibrosis and proliferative vasculopathy in systemic sclerosis. <i>Rheumatology</i> , 2013 , 52, 790-9 Exogenous application of hydrogen sulfide donor attenuates inflammatory reactions through the L-selectin-involved pathway in the cutaneous reverse passive Arthus reaction. <i>Journal of Leukocyte</i>	3.9	34
73 72	Decreased cathepsin V expression due to Fli1 deficiency contributes to the development of dermal fibrosis and proliferative vasculopathy in systemic sclerosis. <i>Rheumatology</i> , 2013 , 52, 790-9 Exogenous application of hydrogen sulfide donor attenuates inflammatory reactions through the L-selectin-involved pathway in the cutaneous reverse passive Arthus reaction. <i>Journal of Leukocyte Biology</i> , 2013 , 93, 573-84 Delayed wound healing due to increased interleukin-10 expression in mice with lymphatic	3.9 6.5	34
73 72 71	Decreased cathepsin V expression due to Fli1 deficiency contributes to the development of dermal fibrosis and proliferative vasculopathy in systemic sclerosis. <i>Rheumatology</i> , 2013 , 52, 790-9 Exogenous application of hydrogen sulfide donor attenuates inflammatory reactions through the L-selectin-involved pathway in the cutaneous reverse passive Arthus reaction. <i>Journal of Leukocyte Biology</i> , 2013 , 93, 573-84 Delayed wound healing due to increased interleukin-10 expression in mice with lymphatic dysfunction. <i>Journal of Leukocyte Biology</i> , 2013 , 94, 137-45 Clinical significance of monitoring serum adiponectin levels during intravenous pulse cyclophosphamide therapy in interstitial lung disease associated with systemic sclerosis. <i>Modern</i>	3.9 6.5 6.5	34 6 19

67	Clinical significance of monitoring serum adiponectin levels during intravenous pulse cyclophosphamide therapy in interstitial lung disease associated with systemic sclerosis. <i>Modern Rheumatology</i> , 2013 , 23, 323-9	3.3	11
66	A possible contribution of altered cathepsin B expression to the development of skin sclerosis and vasculopathy in systemic sclerosis. <i>PLoS ONE</i> , 2012 , 7, e32272	3.7	31
65	Proteasome inhibitor bortezomib ameliorates intestinal injury in mice. PLoS ONE, 2012, 7, e34587	3.7	10
64	Angiogenin levels are increased in lesional skin and sera in patients with erythrodermic cutaneous T cell lymphoma. <i>Archives of Dermatological Research</i> , 2012 , 304, 401-6	3.3	14
63	Increased serum levels of soluble CD163 in patients with scleroderma. <i>Clinical Rheumatology</i> , 2012 , 31, 1059-64	3.9	28
62	Serum IL-31 levels are increased in patients with cutaneous T-cell lymphoma. <i>Acta Dermato-Venereologica</i> , 2012 , 92, 282-3	2.2	48
61	Low herpesvirus entry mediator (HVEM) expression on dermal fibroblasts contributes to a Th2-dominant microenvironment in advanced cutaneous T-cell lymphoma. <i>Journal of Investigative Dermatology</i> , 2012 , 132, 1280-9	4.3	21
60	Lymphatic dysfunction impairs antigen-specific immunization, but augments tissue swelling following contact with allergens. <i>Journal of Investigative Dermatology</i> , 2012 , 132, 667-76	4.3	16
59	Clinical significance of serum growth differentiation factor-15 levels in systemic sclerosis: association with disease severity. <i>Modern Rheumatology</i> , 2012 , 22, 668-675	3.3	26
58	P2Y6 receptor signaling pathway mediates inflammatory responses induced by monosodium urate crystals. <i>Journal of Immunology</i> , 2012 , 188, 436-44	5.3	62
57	IL-10-producing regulatory B10 cells inhibit intestinal injury in a mouse model. <i>American Journal of Pathology</i> , 2011 , 178, 735-43	5.8	138
56	Serum CCL23 levels are increased in patients with systemic sclerosis. <i>Archives of Dermatological Research</i> , 2011 , 303, 29-34	3.3	15
55	Serum IL-33 levels are raised in patients with systemic sclerosis: association with extent of skin sclerosis and severity of pulmonary fibrosis. <i>Clinical Rheumatology</i> , 2011 , 30, 825-30	3.9	91
54	Increasing levels of serum antioxidant status, total antioxidant power, in systemic sclerosis. <i>Clinical Rheumatology</i> , 2011 , 30, 921-5	3.9	23
53	A portable dermatoscope for easy, rapid examination of periungual nailfold capillary changes in patients with systemic sclerosis. <i>Rheumatology International</i> , 2011 , 31, 1601-6	3.6	17
52	The specific free radical scavenger edaravone suppresses fibrosis in the bleomycin-induced and tight skin mouse models of systemic sclerosis. <i>Arthritis and Rheumatism</i> , 2011 , 63, 3086-97		30
51	Immunization with DNA topoisomerase I and Freund's complete adjuvant induces skin and lung fibrosis and autoimmunity via interleukin-6 signaling. <i>Arthritis and Rheumatism</i> , 2011 , 63, 3575-85		60
50	Significant attenuation of macrovascular involvement by bosentan in a patient with diffuse cutaneous systemic sclerosis with multiple digital ulcers and gangrene. <i>Modern Rheumatology</i> , 2011 21 548-552	3.3	13

(2008-2011)

49	The development of Th1-mediated sarcoidosis improves the clinical course of Th2-mediated atopic dermatitis. <i>Modern Rheumatology</i> , 2011 , 21, 406-409	3.3	4
48	Disseminated cutaneous and visceral Kaposill sarcoma in a patient with rheumatoid arthritis receiving corticosteroids and tacrolimus. <i>Modern Rheumatology</i> , 2011 , 21, 309-312	3.3	3
47	Elevated serum interleukin-27 levels in patients with systemic sclerosis: association with T cell, B cell and fibroblast activation. <i>Annals of the Rheumatic Diseases</i> , 2011 , 70, 194-200	2.4	46
46	Significant attenuation of macrovascular involvement by bosentan in a patient with diffuse cutaneous systemic sclerosis with multiple digital ulcers and gangrene. <i>Modern Rheumatology</i> , 2011 , 21, 548-52	3.3	7
45	Kunihiko Tamaki, M.D., Ph.D., Emeritus Professor, University of Tokyo, 19462010. <i>Journal of Dermatology</i> , 2010 , 37, 777-779	1.6	1
44	The proteasome inhibitor bortezomib inhibits T cell-dependent inflammatory responses. <i>Journal of Leukocyte Biology</i> , 2010 , 88, 117-22	6.5	23
43	Regulatory B cells (B10 cells) have a suppressive role in murine lupus: CD19 and B10 cell deficiency exacerbates systemic autoimmunity. <i>Journal of Immunology</i> , 2010 , 184, 4801-9	5.3	242
42	Cell adhesion molecules regulate fibrotic process via Th1/Th2/Th17 cell balance in a bleomycin-induced scleroderma model. <i>Journal of Immunology</i> , 2010 , 185, 2502-15	5.3	103
41	Mobilization of endothelial progenitor cells by intravenous cyclophosphamide in patients with systemic sclerosis. <i>Rheumatology</i> , 2010 , 49, 2375-80	3.9	26
40	Bortezomib: a promising treatment for inflammatory diseases. <i>International Journal of Clinical Rheumatology</i> , 2010 , 5, 513-514	1.5	
39	Studies on 1-(2-phenethyl)-4-(N-propionylanilino)piperidine (fentanyl) and related compounds VII. Quantification of alpha-methylfentanyl metabolites excreted in rat urine. <i>Forensic Science International</i> , 2010 , 195, 68-72	2.6	16
38	Treatment with rapamycin prevents fibrosis in tight-skin and bleomycin-induced mouse models of systemic sclerosis. <i>Arthritis and Rheumatism</i> , 2010 , 62, 2476-87		100
37	Increased serum pentraxin 3 in patients with systemic sclerosis. <i>Journal of Rheumatology</i> , 2009 , 36, 976	-8431	37
36	Clinical significance of serum HMGB-1 and sRAGE levels in systemic sclerosis: association with disease severity. <i>Journal of Clinical Immunology</i> , 2009 , 29, 180-9	5.7	83
35	CD19, a response regulator of B lymphocytes, regulates wound healing through hyaluronan-induced TLR4 signaling. <i>American Journal of Pathology</i> , 2009 , 175, 649-60	5.8	65
34	Clinical association of serum interleukin-17 levels in systemic sclerosis: is systemic sclerosis a Th17 disease?. <i>Journal of Dermatological Science</i> , 2008 , 50, 240-2	4.3	100
33	CD19 regulates skin and lung fibrosis via Toll-like receptor signaling in a model of bleomycin-induced scleroderma. <i>American Journal of Pathology</i> , 2008 , 172, 1650-63	5.8	165
32	Increased Serum Soluble OX40 in Patients with Systemic Sclerosis. <i>Journal of Rheumatology</i> , 2008 , 35, 2359-2362	4.1	44

31	Clinical significance of serum hyaluronan levels in systemic sclerosis: association with disease severity. <i>Journal of Rheumatology</i> , 2008 , 35, 1825-9	4.1	23
30	BAFF antagonist attenuates the development of skin fibrosis in tight-skin mice. <i>Journal of Investigative Dermatology</i> , 2007 , 127, 2772-80	4.3	58
29	Intercellular adhesion molecule-1 deficiency attenuates the development of skin fibrosis in tight-skin mice. <i>Journal of Immunology</i> , 2007 , 179, 698-707	5.3	32
28	Phase-dependent roles of E-selectin during chronic contact hypersensitivity responses. <i>American Journal of Pathology</i> , 2007 , 170, 1649-58	5.8	12
27	CD19 expression in B cells is important for suppression of contact hypersensitivity. <i>American Journal of Pathology</i> , 2007 , 171, 560-70	5.8	92
26	Elevated serum BAFF levels in patients with systemic sclerosis: enhanced BAFF signaling in systemic sclerosis B lymphocytes. <i>Arthritis and Rheumatism</i> , 2006 , 54, 192-201		202
25	E- and P-selectins synergistically inhibit bleomycin-induced pulmonary fibrosis. <i>American Journal of Pathology</i> , 2006 , 169, 740-9	5.8	13
24	Longitudinal analysis of serum cytokine concentrations in systemic sclerosis: association of interleukin 12 elevation with spontaneous regression of skin sclerosis. <i>Journal of Rheumatology</i> , 2006 , 33, 275-84	4.1	101
23	P-selectin glycoprotein ligand-1 is required for the development of cutaneous vasculitis induced by immune complex deposition. <i>Journal of Leukocyte Biology</i> , 2004 , 76, 374-82	6.5	14
22	Altered blood B lymphocyte homeostasis in systemic sclerosis: expanded naive B cells and diminished but activated memory B cells. <i>Arthritis and Rheumatism</i> , 2004 , 50, 1918-27		223
22			223 69
	diminished but activated memory B cells. <i>Arthritis and Rheumatism</i> , 2004 , 50, 1918-27 Association of a functional CD19 polymorphism with susceptibility to systemic sclerosis. <i>Arthritis</i>	4.3	
21	diminished but activated memory B cells. <i>Arthritis and Rheumatism</i> , 2004 , 50, 1918-27 Association of a functional CD19 polymorphism with susceptibility to systemic sclerosis. <i>Arthritis and Rheumatism</i> , 2004 , 50, 4002-7 Altered B lymphocyte function induces systemic autoimmunity in systemic sclerosis. <i>Molecular</i>	4.3	69
21	diminished but activated memory B cells. <i>Arthritis and Rheumatism</i> , 2004 , 50, 1918-27 Association of a functional CD19 polymorphism with susceptibility to systemic sclerosis. <i>Arthritis and Rheumatism</i> , 2004 , 50, 4002-7 Altered B lymphocyte function induces systemic autoimmunity in systemic sclerosis. <i>Molecular Immunology</i> , 2004 , 41, 1123-33 Relative contributions of selectins and intercellular adhesion molecule-1 to tissue injury induced by		69 99
21 20 19	Association of a functional CD19 polymorphism with susceptibility to systemic sclerosis. <i>Arthritis and Rheumatism</i> , 2004 , 50, 4002-7 Altered B lymphocyte function induces systemic autoimmunity in systemic sclerosis. <i>Molecular Immunology</i> , 2004 , 41, 1123-33 Relative contributions of selectins and intercellular adhesion molecule-1 to tissue injury induced by immune complex deposition. <i>American Journal of Pathology</i> , 2003 , 162, 1463-73 L-selectin or ICAM-1 deficiency reduces an immediate-type hypersensitivity response by preventing mast cell recruitment in repeated elicitation of contact hypersensitivity. <i>Journal of Immunology</i> ,	5.8	69 99 32
21 20 19	Association of a functional CD19 polymorphism with susceptibility to systemic sclerosis. <i>Arthritis and Rheumatism</i> , 2004 , 50, 4002-7 Altered B lymphocyte function induces systemic autoimmunity in systemic sclerosis. <i>Molecular Immunology</i> , 2004 , 41, 1123-33 Relative contributions of selectins and intercellular adhesion molecule-1 to tissue injury induced by immune complex deposition. <i>American Journal of Pathology</i> , 2003 , 162, 1463-73 L-selectin or ICAM-1 deficiency reduces an immediate-type hypersensitivity response by preventing mast cell recruitment in repeated elicitation of contact hypersensitivity. <i>Journal of Immunology</i> , 2003 , 170, 4325-34 The cutaneous reverse Arthus reaction requires intercellular adhesion molecule 1 and L-selectin	5.8 5·3	69 99 32 50
21 20 19 18	Association of a functional CD19 polymorphism with susceptibility to systemic sclerosis. <i>Arthritis and Rheumatism</i> , 2004 , 50, 4002-7 Altered B lymphocyte function induces systemic autoimmunity in systemic sclerosis. <i>Molecular Immunology</i> , 2004 , 41, 1123-33 Relative contributions of selectins and intercellular adhesion molecule-1 to tissue injury induced by immune complex deposition. <i>American Journal of Pathology</i> , 2003 , 162, 1463-73 L-selectin or ICAM-1 deficiency reduces an immediate-type hypersensitivity response by preventing mast cell recruitment in repeated elicitation of contact hypersensitivity. <i>Journal of Immunology</i> , 2003 , 170, 4325-34 The cutaneous reverse Arthus reaction requires intercellular adhesion molecule 1 and L-selectin expression. <i>Journal of Immunology</i> , 2002 , 168, 2970-8 Intercellular adhesion molecule-1 and L-selectin regulate bleomycin-induced lung fibrosis. <i>American</i>	5.8 5.3 5.3	69 99 32 50 41

LIST OF PUBLICATIONS

13	Serum levels of interleukin-6 and interleukin-10 correlate with total skin thickness score in patients with systemic sclerosis. <i>Journal of Dermatological Science</i> , 2001 , 27, 140-6	4.3	216
12	Quantitative genetic variation in CD19 expression correlates with autoimmunity. <i>Journal of Immunology</i> , 2000 , 165, 6635-43	5.3	255
11	Delayed wound healing in the absence of intercellular adhesion molecule-1 or L-selectin expression. <i>American Journal of Pathology</i> , 2000 , 157, 237-47	5.8	132
10	Abnormalities of adhesion molecules and chemokines in scleroderma. <i>Current Opinion in Rheumatology</i> , 1999 , 11, 503-507	5.3	29
9	Elevated soluble CD23 levels in the sera from patients with localized scleroderma. <i>Archives of Dermatological Research</i> , 1996 , 288, 74-8	3.3	17
8	Soluble CD4 and CD8 in serum from patients with localized scleroderma. <i>Archives of Dermatological Research</i> , 1996 , 288, 358-62	3.3	30
7	Elevated soluble CD23 levels in the sera from patients with localized scleroderma 1996 , 288, 74		2
6	Soluble CD4 and CD8 in serum from patients with localized scleroderma 1996 , 288, 358		1
5	Autoantibodies to pyruvate dehydrogenase complex in patients with systemic sclerosis. Possible role of anti-E1 alpha antibody as a serologic indicator for development of primary biliary cirrhosis. <i>Arthritis and Rheumatism</i> , 1995 , 38, 985-9		20
4	Antihistone antibodies in systemic sclerosis. Association with pulmonary fibrosis. <i>Arthritis and Rheumatism</i> , 1994 , 37, 391-4		104
3	Diagnostic significance of nailfold bleeding in scleroderma spectrum disorders. <i>Journal of the American Academy of Dermatology</i> , 1993 , 28, 198-203	4.5	23
2	Antihistone antibodies in patients with localized scleroderma. Arthritis and Rheumatism, 1993, 36, 1137-	-41	136
1	Parabiotic Experiments on the Development of Neurohypophysis (With Bufo vulgaris formosus). Okajimas Folia Anatomica Japonica, 1959, 32, 313-317 3	0.3	