

Di Che

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2495059/publications.pdf>

Version: 2024-02-01

39
papers

506
citations

1039406

9
h-index

752256

20
g-index

42
all docs

42
docs citations

42
times ranked

751
citing authors

#	ARTICLE	IF	CITATIONS
1	Large retrospective cohort study of the association between maternal 25-hydroxyvitamin D status and birth weight of neonate. <i>Journal of Maternal-Fetal and Neonatal Medicine</i> , 2024, 35, 7231-7237.	0.7	4
2	Correlation between vitamin D levels and bone metabolism in children with cow's milk allergy. <i>Journal of International Medical Research</i> , 2022, 50, 030006052110660.	0.4	0
3	Association study of miR-149, miR-196a2, and miR-499a polymorphisms with coronary artery aneurysm of Kawasaki disease in southern Chinese population. <i>Journal of Gene Medicine</i> , 2022, 24, e3405.	1.4	3
4	Association between the rs3802201 polymorphism of the lncRNA MIR2052HG gene and the risk of recurrent miscarriage in a Southern Chinese population. <i>Journal of Clinical Laboratory Analysis</i> , 2022, 36, e24167.	0.9	2
5	The rs7404339 AA Genotype in CDH5 Contributes to Increased Risks of Kawasaki Disease and Coronary Artery Lesions in a Southern Chinese Child Population. <i>Frontiers in Cardiovascular Medicine</i> , 2022, 9, 760982.	1.1	0
6	Upregulation of PEDF Predicts a Poor Prognosis and Promotes Esophageal Squamous Cell Carcinoma Progression by Modulating the MAPK/ERK Signaling Pathway. <i>Frontiers in Oncology</i> , 2021, 11, 625612.	1.3	4
7	Homozygous of MRP4 Gene rs1751034 C Allele Is Related to Increased Risk of Intravenous Immunoglobulin Resistance in Kawasaki Disease. <i>Frontiers in Genetics</i> , 2021, 12, 510350.	1.1	4
8	The lncRNA ANRIL Gene rs2151280 GG Genotype is Associated with Increased Susceptibility to Recurrent Miscarriage in a Southern Chinese Population. <i>Journal of Inflammation Research</i> , 2021, Volume 14, 2865-2872.	1.6	3
9	FND1 Polymorphism (rs3003174 C > T) Increased the Incidence of Coronary Artery Aneurysm in Patients with Kawasaki Disease in a Southern Chinese Population. <i>Journal of Inflammation Research</i> , 2021, Volume 14, 2633-2640.	1.6	5
10	The lncRNA CCAT2 Rs6983267 G Variant Contributes to Increased Sepsis Susceptibility in a Southern Chinese Population. <i>Infection and Drug Resistance</i> , 2021, Volume 14, 2969-2976.	1.1	2
11	Protective Effect of TNFRSF11A rs7239667 G > C Gene Polymorphism on Coronary Outcome of Kawasaki Disease in Southern Chinese Population. <i>Frontiers in Genetics</i> , 2021, 12, 691282.	1.1	0
12	Heightened Local Th17 Cell Inflammation Is Associated with Severe Community-Acquired Pneumonia in Children under the Age of 1 Year. <i>Mediators of Inflammation</i> , 2021, 2021, 1-13.	1.4	4
13	The SERPINA4 rs2070777 AA Genotype is Associated with an Increased Risk of Recurrent Miscarriage in a Southern Chinese Population. <i>International Journal of Women's Health</i> , 2021, Volume 13, 111-117.	1.1	2
14	Effect of Pandemic-Related Confinement on Vitamin D Status Among Children Aged 0-6 Years in Guangzhou, China: A Cross-Sectional Study. <i>Risk Management and Healthcare Policy</i> , 2020, Volume 13, 2669-2675.	1.2	22
15	lncRNA SOX2OT rs9839776 Polymorphism Reduces Sepsis Susceptibility in Southern Chinese Children. <i>Journal of Inflammation Research</i> , 2020, Volume 13, 1095-1101.	1.6	5
16	The rs1051931 G>A Polymorphism in the PLA2G7 Gene Confers Resistance to Immunoglobulin Therapy in Kawasaki Disease in a Southern Chinese Population. <i>Frontiers in Pediatrics</i> , 2020, 8, 338.	0.9	3
17	MiR-124-3p helps to protect against acute respiratory distress syndrome by targeting p65. <i>Bioscience Reports</i> , 2020, 40, .	1.1	24
18	Estrogen-Related Hormones Induce Apoptosis by Stabilizing Schlafen-12 Protein Turnover. <i>Molecular Cell</i> , 2019, 75, 1103-1116.e9.	4.5	55

#	ARTICLE	IF	CITATIONS
19	Mucosal Profiling of Pediatric-Onset Colitis and IBD Reveals Common Pathogenics and Therapeutic Pathways. <i>Cell</i> , 2019, 179, 1160-1176.e24.	13.5	163
20	LncRNA HULC Polymorphism Is Associated With Recurrent Spontaneous Abortion Susceptibility in the Southern Chinese Population. <i>Frontiers in Genetics</i> , 2019, 10, 918.	1.1	10
21	Association between the TOX3 rs3803662 C>T polymorphism and recurrent miscarriage in a southern Chinese population. <i>Journal of Clinical Laboratory Analysis</i> , 2019, 33, e22992.	0.9	2
22	Association between the rs2288947 polymorphism of the lncRNA <i>TINCR</i> gene and the risk of recurrent miscarriage in a Southern Chinese population. <i>Journal of Clinical Laboratory Analysis</i> , 2019, 33, e22919.	0.9	7
23	The IL-1B Gene Polymorphisms rs16944 and rs1143627 Contribute to an Increased Risk of Coronary Artery Lesions in Southern Chinese Children with Kawasaki Disease. <i>Journal of Immunology Research</i> , 2019, 2019, 1-7.	0.9	24
24	The <i>miRNA-608</i> rs4919510 G>C polymorphism confers reduce coronary injury of Kawasaki disease in a Southern Chinese population. <i>Bioscience Reports</i> , 2019, 39, .	1.1	3
25	The lncRNA MALAT1 rs619586 G Variant Confers Decreased Susceptibility to Recurrent Miscarriage. <i>Frontiers in Physiology</i> , 2019, 10, 385.	1.3	24
26	The lncRNA <i>CCAT2</i> rs6983267 G allele is associated with decreased susceptibility to recurrent miscarriage. <i>Journal of Cellular Physiology</i> , 2019, 234, 20577-20583.	2.0	16
27	Serum exosomal microRNA <i>letâ€“7iâ€“3p</i> as candidate diagnostic biomarker for Kawasaki disease patients with coronary artery aneurysm. <i>IUBMB Life</i> , 2019, 71, 891-900.	1.5	19
28	An Angiotensinogen Gene Polymorphism (rs5050) Is Associated with the Risk of Coronary Artery Aneurysm in Southern Chinese Children with Kawasaki Disease. <i>Disease Markers</i> , 2019, 2019, 1-7.	0.6	9
29	The lncRNA SOX2OT rs9839776 C>T Polymorphism Indicates Recurrent Miscarriage Susceptibility in a Southern Chinese Population. <i>Mediators of Inflammation</i> , 2019, 2019, 1-6.	1.4	9
30	A <i>PEAR1</i> polymorphism (<i>rs12041331</i>) is associated with risk of coronary artery aneurysm in Kawasaki disease. <i>Annals of Human Genetics</i> , 2019, 83, 54-62.	0.3	12
31	P2RY12:rs7637803 TT variant genotype increases coronary artery aneurysm risk in Kawasaki disease in a southern Chinese population. <i>Journal of Gene Medicine</i> , 2019, 21, e3066.	1.4	10
32	Lack of association between <i>miR-218</i> rs11134527 A>G and Kawasaki disease susceptibility. <i>Bioscience Reports</i> , 2018, 38, .	1.1	5
33	ABCC4 Variants Modify Susceptibility to Kawasaki Disease in a Southern Chinese Population. <i>Disease Markers</i> , 2018, 2018, 1-7.	0.6	9
34	TBXA2R rs4523 G allele is associated with decreased susceptibility to Kawasaki disease. <i>Cytokine</i> , 2018, 111, 216-221.	1.4	9
35	Immature platelets and antiplatelet therapy response to aspirin in Kawasaki disease. <i>Drug Design, Development and Therapy</i> , 2018, Volume 12, 1353-1362.	2.0	12
36	The rs1625579 T>G polymorphism in the <i>miRNA-137</i> gene confers a risk of early-onset Kawasaki disease in a southern Chinese population. <i>Infection and Drug Resistance</i> , 2018, Volume 11, 1055-1060.	1.1	11

#	ARTICLE	IF	CITATIONS
37	The association between the <i>miR-146a</i> rs2910164 C>G polymorphism and Kawasaki disease in a southern Chinese population. Bioscience Reports, 2018, 38, .	1.1	9
38	The EIF2AK4/rs4594236 AG/GG Genotype Is a Hazard Factor of Immunoglobulin Therapy Resistance in Southern Chinese Kawasaki Disease Patients. Frontiers in Genetics, 0, 13, .	1.1	1
39	The rs8506 TT Genotype in lincRNA-NR_024015 Contributes to the Risk of Sepsis in a Southern Chinese Child Population. Frontiers in Public Health, 0, 10, .	1.3	0