

Tatsuya Umecky

List of Publications by Citations

Source: <https://exaly.com/author-pdf/2500440/tatsuya-umecky-publications-by-citations.pdf>

Version: 2024-04-24

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

53
papers

1,229
citations

21
h-index

33
g-index

54
ext. papers

1,302
ext. citations

2.9
avg, IF

4.24
L-index

#	Paper	IF	Citations
53	Existing condition and migration property of ions in lithium electrolytes with ionic liquid solvent. <i>Journal of Physical Chemistry B</i> , 2007 , 111, 11794-802	3.4	109
52	Self-diffusion coefficients of 1-butyl-3-methylimidazolium hexafluorophosphate with pulsed-field gradient spin-echo NMR technique. <i>Fluid Phase Equilibria</i> , 2005 , 228-229, 329-333	2.5	79
51	Solution structures of 1-butyl-3-methylimidazolium hexafluorophosphate ionic liquid saturated with CO ₂ : Experimental evidence of specific anion-CO ₂ interaction. <i>Journal of Physical Chemistry B</i> , 2005 , 109, 13847-50	3.4	78
50	Low Melting and Electrochemically Stable Ionic Liquids Based on Asymmetric Fluorosulfonyl(trifluoromethylsulfonyl)amide. <i>Chemistry Letters</i> , 2008 , 37, 1020-1021	1.7	59
49	Direct measurements of ionic mobility of ionic liquids using the electric field applying pulsed gradient spin-echo NMR. <i>Journal of Physical Chemistry B</i> , 2009 , 113, 8466-8	3.4	49
48	Ionization condition of lithium ionic liquid electrolytes under the solvation effect of liquid and solid solvents. <i>Journal of Physical Chemistry B</i> , 2008 , 112, 3357-64	3.4	49
47	Electrical Conductivities, Viscosities, and Densities of N-Methoxymethyl- and N-Butyl-N-methylpyrrolidinium Ionic Liquids with the Bis(fluorosulfonyl)amide Anion. <i>Journal of Chemical & Engineering Data</i> , 2012 , 57, 751-755	2.8	47
46	Infinite Dilution Binary Diffusion Coefficients of Several α -Amino Acids in Water over a Temperature Range from (293.2 to 333.2) K with the Taylor Dispersion Technique. <i>Journal of Chemical & Engineering Data</i> , 2006 , 51, 1705-1710	2.8	47
45	CO ₂ solubility in and physical properties for ionic liquid mixtures of 1-butyl-3-methylimidazolium acetate and 1-butyl-3-methylimidazolium bis(trifluoromethanesulfonyl)amide. <i>Journal of Molecular Liquids</i> , 2016 , 217, 112-119	6	46
44	CO ₂ absorption properties, densities, viscosities, and electrical conductivities of ethylimidazolium and 1-ethyl-3-methylimidazolium ionic liquids. <i>Fluid Phase Equilibria</i> , 2014 , 362, 300-306	2.5	41
43	Interactions of perfluoroalkyltrifluoroborate anions with Li ⁺ ion and imidazolium cation: effects of perfluoroalkyl chain on motion of ions in ionic liquids. <i>Journal of Physical Chemistry B</i> , 2010 , 114, 11390-3	3.4	40
42	Microscopic interactions of the imidazolium-based ionic liquid with molecular liquids depending on their electron-donicity. <i>Physical Chemistry Chemical Physics</i> , 2014 , 16, 23627-38	3.6	38
41	Alkoxy chains in ionic liquid anions; effect of introducing ether oxygen into perfluoroalkylborate on physical and thermal properties. <i>Chemical Communications</i> , 2010 , 46, 1730-2	5.8	37
40	Water-induced Acceleration of Transport Properties in Hydrophobic 1-Butyl-3-methylimidazolium Hexafluorophosphate Ionic Liquid. <i>Chemistry Letters</i> , 2005 , 34, 324-325	1.7	35
39	Electrical Conductivities, Viscosities, and Densities of N-Acetoxyethyl-N,N-dimethyl-N-ethylammonium and N,N-Dimethyl-N-ethyl-N-methoxyethoxyethylammonium Bis(trifluoromethanesulfonyl)amide and Their Nonfunctionalized Analogues. <i>Journal of Chemical & Engineering Data</i> , 2013 , 58, 270-276	2.8	31
38	Effects of alkyl chain on transport properties in 1-alkyl-3-methylimidazolium hexafluorophosphates. <i>Journal of Molecular Liquids</i> , 2005 , 119, 77-81	6	29
37	Pressure-volume-temperature-composition relations for carbon dioxide+pyrrolidinium-based ionic liquid binary systems. <i>Fluid Phase Equilibria</i> , 2013 , 360, 253-259	2.5	27

36	Binary Diffusion Coefficients of Aqueous Phenylalanine, Tyrosine Isomers, and Aminobutyric Acids at Infinitesimal Concentration and Temperatures from (293.2 to 333.2) K. <i>Journal of Chemical & Engineering Data</i> , 2013 , 58, 1909-1917	2.8	25
35	Experimental Determination of Reorientational Correlation Time of CO ₂ over a Wide Range of Density and Temperature. <i>Journal of Physical Chemistry B</i> , 2003 , 107, 12003-12008	3.4	25
34	Possible Proton Conduction Mechanism in Pseudo-Protic Ionic Liquids: A Concept of Specific Proton Conduction. <i>Journal of Physical Chemistry B</i> , 2019 , 123, 6244-6252	3.4	24
33	CO ₂ Solubilities in Ammonium Bis(trifluoromethanesulfonyl)amide Ionic Liquids: Effects of Ester and Ether Groups. <i>Journal of Chemical & Engineering Data</i> , 2014 , 59, 1435-1440	2.8	24
32	Effects of Tetrafluoroborate and Bis(trifluoromethylsulfonyl)amide Anions on the Microscopic Structures of 1-Methyl-3-octylimidazolium-Based Ionic Liquids and Benzene Mixtures: A Multiple Approach by ATR-IR, NMR, and Femtosecond Raman-Induced Kerr Effect Spectroscopy. <i>Journal of Physical Chemistry B</i> , 2016 , 120, 513-26	3.4	21
31	CO ₂ solubility and physical properties of N-(2-hydroxyethyl)pyridinium bis(trifluoromethanesulfonyl)amide. <i>Fluid Phase Equilibria</i> , 2013 , 357, 64-70	2.5	20
30	Effect of partial pressure on CO ₂ solubility in ionic liquid mixtures of 1-butyl-3-methylimidazolium acetate and 1-butyl-3-methylimidazolium bis(trifluoromethanesulfonyl)amide. <i>Fluid Phase Equilibria</i> , 2016 , 420, 74-82	2.5	17
29	Effects of dissolved water on Li ⁺ solvation in 1-ethyl-3-methylimidazolium bis(trifluoromethanesulfonyl)amide ionic liquid studied by NMR. <i>Journal of Physical Chemistry B</i> , 2013 , 117, 16219-26	3.4	17
28	High-pressure NMR studies on solvation structure in supercritical carbon dioxide. <i>Fluid Phase Equilibria</i> , 2002 , 194-197, 859-868	2.5	16
27	⁹ Be NMR Relaxation Measurements of Bis(acetylacetonato)beryllium(II) in Liquid and Supercritical Carbon Dioxide: A Clear Evidence of Near-Critical Solvation Effect on Rotational Correlation Time. <i>Journal of Physical Chemistry B</i> , 2002 , 106, 11114-11119	3.4	16
26	Effects of hydroxyl groups on binary diffusion coefficients of amino acids in dilute aqueous solutions. <i>Fluid Phase Equilibria</i> , 2008 , 264, 18-22	2.5	15
25	CO ₂ Absorption Properties and Mechanisms for 1-Ethyl-3-methylimidazolium Ether-Functionalized Carboxylates. <i>Industrial & Engineering Chemistry Research</i> , 2016 , 55, 12949-12961	3.9	15
24	Development of High-Pressure Electric Conductivity Cell and its Application: Pressure Effect of Carbon Dioxide on Electric Conductivity of Ionic Liquid. <i>Electrochemistry</i> , 2004 , 72, 703-705	1.2	13
23	Binary Diffusion Coefficients of Aqueous Straight-Chain Amino Acids at Infinitesimal Concentration and Temperatures from (298.2 to 333.2) K. <i>Journal of Chemical & Engineering Data</i> , 2013 , 58, 2848-2853	2.8	12
22	High-Pressure ¹⁹ F NMR Measurements of a Series of Fluorinated Benzenes in Supercritical Carbon Dioxide. <i>Journal of Solution Chemistry</i> , 2004 , 33, 863-874	1.8	12
21	SANS, ATR-IR, and 1D- and 2D-NMR studies of mixing states of imidazolium-based ionic liquid and aryl solvents. <i>Physical Chemistry Chemical Physics</i> , 2013 , 15, 20565-76	3.6	11
20	Determination of anisotropic solvation structure of octafluorotoluene in supercritical carbon dioxide by means of solvent-induced ¹⁹ F NMR chemical shift. <i>Chemical Physics Letters</i> , 2001 , 338, 95-100	2.5	11
19	Solvent-Dependent Properties and Higher-Order Structures of Aryl Alcohol + Surfactant Molecular Gels. <i>Langmuir</i> , 2016 , 32, 4352-60	4	10

18	Effect of CO ₂ dissolution on electrical conductivity and self-diffusion coefficients of 1-butyl-3-methylimidazolium hexafluorophosphate ionic liquid. <i>Fluid Phase Equilibria</i> , 2013 , 357, 76-79	2.5	10
17	CO ₂ absorption features of 1-ethyl-3-methylimidazolium ionic liquids with 2,4-pentanedionate and its fluorine derivatives. <i>Journal of CO₂ Utilization</i> , 2019 , 31, 75-84	7.6	9
16	A Study of the Solvation Structure of L-Leucine in Alcohol-Water Binary Solvents through Molecular Dynamics Simulations and FTIR and NMR Spectroscopy. <i>ChemPhysChem</i> , 2015 , 16, 3190-9	3.2	9
15	Solvation structure and dynamics of Li ⁺ in Lewis-basic ionic liquid of 1-octyl-4-aza-1-azoniabicyclo[2.2.2]octane bis(trifluoromethanesulfonyl)amide. <i>Journal of Molecular Liquids</i> , 2015 , 209, 557-562	6	8
14	A Novel High-Pressure NMR Cell Consisting of Double Tube Structure for the Convenient On-Line Measurements. <i>Chemistry Letters</i> , 2002 , 31, 118-119	1.7	8
13	Structures of Naphthol⊗OT Self-assembly Organogels and Their Applications to Dispersing Media of Rare-earth Complexes. <i>Chemistry Letters</i> , 2014 , 43, 1861-1863	1.7	6
12	Solvation Structure of 1,3-Butanediol in Aqueous Binary Solvents with Acetonitrile, 1,4-Dioxane, and Dimethyl Sulfoxide Studied by IR, NMR, and Molecular Dynamics Simulation. <i>Journal of Physical Chemistry B</i> , 2017 , 121, 4864-4872	3.4	5
11	Complex formation of nickel(ii) with dimethyl sulfoxide, methanol, and acetonitrile in a TFSA-based ionic liquid of [Cmim][TFSA]. <i>Physical Chemistry Chemical Physics</i> , 2017 , 19, 31335-31344	3.6	5
10	SANS, infrared, and ⁷ Li and ²³ Na NMR studies on phase separation of alkali halide-acetonitrile-water mixtures by cooling. <i>Journal of Physical Chemistry B</i> , 2013 , 117, 2438-48	3.4	5
9	Fluorination effects on rotational correlation times of tris(β-diketonato)aluminum(III) in CO ₂ by ²⁷ Al NMR relaxation measurements. <i>Journal of Physical Chemistry B</i> , 2011 , 115, 10622-30	3.4	5
8	Proton Conduction Properties of Sulfonicacid Type Polymer Gel Electrolytes. <i>Journal of Physical Chemistry C</i> , 2009 , 113, 3021-3028	3.8	3
7	¹⁹ F NMR chemical shifts of CF ₄ in CO ₂ over a wide pressure range at different temperatures. <i>Magnetic Resonance in Chemistry</i> , 2003 , 41, 75-76	2.1	3
6	Role of water in complexation of 1,4,7,10,13,16-hexaoxacyclooctadecane (18-crown-6) with Li and K in hydrophobic 1-ethyl-3-methylimidazolium bis(trifluoromethanesulfonyl)amide ionic liquid. <i>Journal of Inclusion Phenomena and Macrocyclic Chemistry</i> , 2014 , 80, 401-407	1.7	2
5	NMR Studies on Solution Structures of Methanol and Ethanol Saturated with CO ₂ . <i>Journal of Solution Chemistry</i> , 2014 , 43, 1539-1549	1.8	2
4	New Ionic Liquids Containing Fluorosulfonyl(trifluoromethylsulfonyl)amide and 5-Phosphoniaspiro[4.4]nonan. <i>ECS Transactions</i> , 2010 , 33, 35-40	1	2
3	Ion Mobility of 1-Ethyl-3-methylimidazolium Tetrafluoroborate and 1-Ethyl-3-methylimidazolium Bis(trifluorosulfonyl)amide Ionic Liquids. <i>ECS Transactions</i> , 2009 , 25, 23-29	1	2
2	In-Situ Observation of Functional Solvents at High Pressures. <i>Review of High Pressure Science and Technology/Koatsuryoku No Kagaku To Gijutsu</i> , 2018 , 28, 88-94	0	
1	Development of CO ₂ Separation Technology Using Ionic Liquids. <i>Zairyo/Journal of the Society of Materials Science, Japan</i> , 2018 , 67, 514-520	0.1	

