

Halil Genç

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2504353/publications.pdf>

Version: 2024-02-01

30
papers

875
citations

566801

15
h-index

552369

26
g-index

30
all docs

30
docs citations

30
times ranked

1626
citing authors

#	ARTICLE	IF	CITATIONS
1	Plasma visfatin levels in patients with newly diagnosed and untreated type 2 diabetes mellitus and impaired glucose tolerance. <i>Diabetes Research and Clinical Practice</i> , 2007, 76, 24-29.	1.1	178
2	Association of plasma visfatin with hepatic and systemic inflammation in nonalcoholic fatty liver disease. <i>Annals of Hepatology</i> , 2013, 12, 380-387.	0.6	83
3	Plasma Apelin and ADMA Levels in Patients with Essential Hypertension. <i>Clinical and Experimental Hypertension</i> , 2010, 32, 179-183.	0.5	82
4	Plasma fetuin-A is associated with endothelial dysfunction and subclinical atherosclerosis in subjects with nonalcoholic fatty liver disease. <i>Clinical Endocrinology</i> , 2013, 78, 712-717.	1.2	73
5	The relationship of serum uric acid with non-alcoholic fatty liver disease. <i>Clinical Biochemistry</i> , 2014, 47, 383-388.	0.8	62
6	Low- and high-density lipoprotein subclasses in subjects with nonalcoholic fatty liver disease. <i>Journal of Clinical Lipidology</i> , 2015, 9, 576-582.	0.6	56
7	Mean platelet volume and its relationship with carotid atherosclerosis in subjects with non-alcoholic fatty liver disease. <i>Upsala Journal of Medical Sciences</i> , 2010, 115, 253-259.	0.4	46
8	LDL-cholesterol lowering increases plasma apelin in isolated hypercholesterolemia. <i>Atherosclerosis</i> , 2009, 204, 222-228.	0.4	45
9	Elevated asymmetric dimethylarginine in plasma: An early marker for endothelial dysfunction in non-alcoholic fatty liver disease?. <i>Diabetes Research and Clinical Practice</i> , 2012, 96, 47-52.	1.1	39
10	Insulin Resistance but Not Visceral Adiposity Index Is Associated with Liver Fibrosis in Nondiabetic Subjects with Nonalcoholic Fatty Liver Disease. <i>Metabolic Syndrome and Related Disorders</i> , 2015, 13, 319-325.	0.5	31
11	Chitotriosidase activity predicts endothelial dysfunction in type-2 diabetes mellitus. <i>Endocrine</i> , 2010, 37, 455-459.	1.1	30
12	Choroidal Thickness Changes in the Acute Attack Period in Patients with Familial Mediterranean Fever. <i>Ophthalmologica</i> , 2016, 235, 72-77.	1.0	28
13	Neutrophil-to-lymphocyte ratio is not a predictor of liver histology in patients with nonalcoholic fatty liver disease. <i>European Journal of Gastroenterology and Hepatology</i> , 2015, 27, 1144-1148.	0.8	23
14	The relationship of circulating fetuin-a with liver histology and biomarkers of systemic inflammation in nondiabetic subjects with nonalcoholic fatty liver disease. <i>Saudi Journal of Gastroenterology</i> , 2015, 21, 139.	0.5	21
15	Circulating vaspin and its relationship with insulin sensitivity, adiponectin, and liver histology in subjects with non-alcoholic steatohepatitis. <i>Scandinavian Journal of Gastroenterology</i> , 2011, 46, 1355-1361.	0.6	16
16	Soluble CD40 ligand, soluble P-selectin and von Willebrand factor levels in subjects with prediabetes: The impact of metabolic syndrome. <i>Clinical Biochemistry</i> , 2012, 45, 92-95.	0.8	13
17	Circulating levels of interleukin-18 in patients with non-alcoholic fatty liver disease. <i>Scandinavian Journal of Clinical and Laboratory Investigation</i> , 2010, 70, 399-403.	0.6	11
18	Efficacy of a Modified Sequential Therapy Including Bismuth Subcitrate as First-Line Therapy to Eradicate <i>Helicobacter pylori</i> in a Turkish Population. <i>Helicobacter</i> , 2012, 17, 486-490.	1.6	9

#	ARTICLE	IF	CITATIONS
19	Non-alcoholic fatty liver disease per se is not associated with carotid atherosclerosis. <i>Gulhane Medical Journal</i> , 2013, 55, 84.	0.1	6
20	The relationship between blood urea nitrogen levels and metabolic, biochemical, and histopathologic findings of nondiabetic, nonhypertensive patients with nonalcoholic fatty liver disease. <i>Turkish Journal of Medical Sciences</i> , 2016, 46, 985-991.	0.4	5
21	The presence of <i>MED1</i> gene mutations in patients with primary osteoarthritis who require surgery. <i>Korean Journal of Internal Medicine</i> , 2013, 28, 594.	0.7	4
22	Soluble CD40 Ligand Levels in Otherwise Healthy Subjects With Impaired Fasting Glucose. <i>Mediators of Inflammation</i> , 2006, 2006, 1-5.	1.4	3
23	Comment and reply on: Serum chemerin and vaspin in non-alcoholic fatty liver disease. <i>Scandinavian Journal of Gastroenterology</i> , 2010, 45, 1000-1002.	0.6	3
24	C reactive protein levels in non-alcoholic fatty liver disease. <i>Journal of Hepatology</i> , 2012, 56, 507-508.	1.8	3
25	Carotid atherosclerosis or nonalcoholic fatty liver disease: Which comes first?. <i>Atherosclerosis</i> , 2013, 231, 187-188.	0.4	3
26	Platelet Aggregation is not Enhanced in Patients with Prediabetes. <i>Upsala Journal of Medical Sciences</i> , 2007, 112, 338-346.	0.4	2
27	Microalbuminuria and nonalcoholic fatty liver disease. <i>Metabolism: Clinical and Experimental</i> , 2010, 59, e5.	1.5	0
28	Non-invasive prediction of fibrosis in non-alcoholic fatty liver disease. <i>Journal of Hepatology</i> , 2011, 55, 498-499.	1.8	0
29	İNFLAMMATUAR BAĞIŞIRSAK HASTALIKLARININ KOROİDAL VE MAKULAR KALINLIK İZERİNE ETKİSİ. <i>Sleyman Demirel Üniversitesi Tıp Fakültesi Dergisi</i> , 0, , .	0.0	0
30	Transforming growth factor β^2 (TGF- β^2) levels in otherwise healthy subjects with impaired glucose tolerance. <i>Endokrynologia Polska</i> , 2010, 61, 691-4.	0.3	0