

# Halil Genç

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2504353/publications.pdf>

Version: 2024-02-01

30  
papers

875  
citations

567281  
15  
h-index

552781  
26  
g-index

30  
all docs

30  
docs citations

30  
times ranked

1626  
citing authors

#	ARTICLE	IF	CITATIONS
1	The relationship between blood urea nitrogen levels and metabolic, biochemical, and histopathologic findings of nondiabetic, nonhypertensive patients with nonalcoholic fatty liver disease. Turkish Journal of Medical Sciences, 2016, 46, 985-991.	0.9	5
2	Choroidal Thickness Changes in the Acute Attack Period in Patients with Familial Mediterranean Fever. Ophthalmologica, 2016, 235, 72-77.	1.9	28
3	Neutrophil-to-lymphocyte ratio is not a predictor of liver histology in patients with nonalcoholic fatty liver disease. European Journal of Gastroenterology and Hepatology, 2015, 27, 1144-1148.	1.6	23
4	Insulin Resistance but Not Visceral Adiposity Index Is Associated with Liver Fibrosis in Nondiabetic Subjects with Nonalcoholic Fatty Liver Disease. Metabolic Syndrome and Related Disorders, 2015, 13, 319-325.	1.3	31
5	Low- and high-density lipoprotein subclasses in subjects with nonalcoholic fatty liver disease. Journal of Clinical Lipidology, 2015, 9, 576-582.	1.5	56
6	The relationship of circulating fetuin-a with liver histology and biomarkers of systemic inflammation in nondiabetic subjects with nonalcoholic fatty liver disease. Saudi Journal of Gastroenterology, 2015, 21, 139.	1.1	21
7	The relationship of serum uric acid with non-alcoholic fatty liver disease. Clinical Biochemistry, 2014, 47, 383-388.	1.9	62
8	Carotid atherosclerosis or nonalcoholic fatty liver disease: Which comes first?. Atherosclerosis, 2013, 231, 187-188.	0.8	3
9	Plasma fetuin-A is associated with endothelial dysfunction and subclinical atherosclerosis in subjects with nonalcoholic fatty liver disease. Clinical Endocrinology, 2013, 78, 712-717.	2.4	73
10	Non-alcoholic fatty liver disease per se is not associated with carotid atherosclerosis. Gulhane Medical Journal, 2013, 55, 84.	0.2	6
11	Association of plasma visfatin with hepatic and systemic inflammation in nonalcoholic fatty liver disease. Annals of Hepatology, 2013, 12, 380-387.	1.5	83
12	The presence of MEFV gene mutations in patients with primary osteoarthritis who require surgery. Korean Journal of Internal Medicine, 2013, 28, 594.	1.7	4
13	Efficacy of a Modified Sequential Therapy Including Bismuth Subcitrate as First-Line Therapy to Eradicate <i>Helicobacter pylori</i> in a Turkish Population. Helicobacter, 2012, 17, 486-490.	3.5	9
14	Soluble CD40 ligand, soluble P-selectin and von Willebrand factor levels in subjects with prediabetes: The impact of metabolic syndrome. Clinical Biochemistry, 2012, 45, 92-95.	1.9	13
15	C reactive protein levels in non-alcoholic fatty liver disease. Journal of Hepatology, 2012, 56, 507-508.	3.7	3
16	Elevated asymmetric dimethylarginine in plasma: An early marker for endothelial dysfunction in non-alcoholic fatty liver disease?. Diabetes Research and Clinical Practice, 2012, 96, 47-52.	2.8	39
17	Circulating vaspin and its relationship with insulin sensitivity, adiponectin, and liver histology in subjects with non-alcoholic steatohepatitis. Scandinavian Journal of Gastroenterology, 2011, 46, 1355-1361.	1.5	16
18	Non-invasive prediction of fibrosis in non-alcoholic fatty liver disease. Journal of Hepatology, 2011, 55, 498-499.	3.7	0

#	ARTICLE	IF	CITATIONS
19	Chitotriosidase activity predicts endothelial dysfunction in type-2 diabetes mellitus. <i>Endocrine</i> , 2010, 37, 455-459.	2.3	30
20	Microalbuminuria and nonalcoholic fatty liver disease. <i>Metabolism: Clinical and Experimental</i> , 2010, 59, e5.	3.4	0
21	Plasma Apelin and ADMA Levels in Patients with Essential Hypertension. <i>Clinical and Experimental Hypertension</i> , 2010, 32, 179-183.	1.3	82
22	Mean platelet volume and its relationship with carotid atherosclerosis in subjects with non-alcoholic fatty liver disease. <i>Uppsala Journal of Medical Sciences</i> , 2010, 115, 253-259.	0.9	46
23	Circulating levels of interleukin-18 in patients with non-alcoholic fatty liver disease. <i>Scandinavian Journal of Clinical and Laboratory Investigation</i> , 2010, 70, 399-403.	1.2	11
24	Comment and reply on: Serum chemerin and vaspin in non-alcoholic fatty liver disease. <i>Scandinavian Journal of Gastroenterology</i> , 2010, 45, 1000-1002.	1.5	3
25	Transforming growth factor $\beta^2$ (TGF- $\beta^2$ ) levels in otherwise healthy subjects with impaired glucose tolerance. <i>Endokrynologia Polska</i> , 2010, 61, 691-4.	1.0	0
26	LDL-cholesterol lowering increases plasma apelin in isolated hypercholesterolemia. <i>Atherosclerosis</i> , 2009, 204, 222-228.	0.8	45
27	Platelet Aggregation is not Enhanced in Patients with Prediabetes. <i>Uppsala Journal of Medical Sciences</i> , 2007, 112, 338-346.	0.9	2
28	Plasma visfatin levels in patients with newly diagnosed and untreated type 2 diabetes mellitus and impaired glucose tolerance. <i>Diabetes Research and Clinical Practice</i> , 2007, 76, 24-29.	2.8	178
29	Soluble CD40 Ligand Levels in Otherwise Healthy Subjects With Impaired Fasting Glucose. <i>Mediators of Inflammation</i> , 2006, 2006, 1-5.	3.0	3
30	İNFLAMMATUAR BAĞIRSAK HASTALIKLARININ KOROİDAL VE MAKULAR KALINLIK İLERİLEME ETKİSİ. Sileyman Demirel Üniversitesi Tıp Fakültesi Dergisi, 0, , .	0.2	0