

# Halil Gen

## List of Publications by Year in Descending Order

**Source:** <https://exaly.com/author-pdf/2504353/halil-genc-publications-by-year.pdf>

**Version:** 2024-04-24

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

29  
papers

690  
citations

14  
h-index

26  
g-index

30  
ext. papers

765  
ext. citations

3.3  
avg, IF

2.92  
L-index

#	Paper	IF	Citations
29	Choroidal Thickness Changes in the Acute Attack Period in Patients with Familial Mediterranean Fever. <i>Ophthalmologica</i> , <b>2016</b> , 235, 72-7	3.7	17
28	The relationship between blood urea nitrogen levels and metabolic, biochemical, and histopathologic findings of nondiabetic, nonhypertensive patients with nonalcoholic fatty liver disease. <i>Turkish Journal of Medical Sciences</i> , <b>2016</b> , 46, 985-91	2.7	4
27	Insulin Resistance but Not Visceral Adiposity Index Is Associated with Liver Fibrosis in Nondiabetic Subjects with Nonalcoholic Fatty Liver Disease. <i>Metabolic Syndrome and Related Disorders</i> , <b>2015</b> , 13, 319-26	2.6	20
26	Low- and high-density lipoprotein subclasses in subjects with nonalcoholic fatty liver disease. <i>Journal of Clinical Lipidology</i> , <b>2015</b> , 9, 576-82	4.9	42
25	Neutrophil-to-lymphocyte ratio is not a predictor of liver histology in patients with nonalcoholic fatty liver disease. <i>European Journal of Gastroenterology and Hepatology</i> , <b>2015</b> , 27, 1144-8	2.2	16
24	The relationship of circulating fetuin-a with liver histology and biomarkers of systemic inflammation in nondiabetic subjects with nonalcoholic fatty liver disease. <i>Saudi Journal of Gastroenterology</i> , <b>2015</b> , 21, 139-45	3	16
23	The relationship of serum uric acid with non-alcoholic fatty liver disease. <i>Clinical Biochemistry</i> , <b>2014</b> , 47, 383-8	3.5	53
22	Carotid atherosclerosis or nonalcoholic fatty liver disease: which comes first?. <i>Atherosclerosis</i> , <b>2013</b> , 231, 187-8	3.1	3
21	Plasma fetuin-A is associated with endothelial dysfunction and subclinical atherosclerosis in subjects with nonalcoholic fatty liver disease. <i>Clinical Endocrinology</i> , <b>2013</b> , 78, 712-7	3.4	59
20	Non-alcoholic fatty liver disease per se is not associated with carotid atherosclerosis. <i>Gulhane Medical Journal</i> , <b>2013</b> , 55, 84		3
19	Association of plasma visfatin with hepatic and systemic inflammation in nonalcoholic fatty liver disease. <i>Annals of Hepatology</i> , <b>2013</b> , 12, 380-387	3.1	64
18	The presence of MEFV gene mutations in patients with primary osteoarthritis who require surgery. <i>Korean Journal of Internal Medicine</i> , <b>2013</b> , 28, 594-8	2.5	2
17	Efficacy of a modified sequential therapy including bismuth subcitrate as first-line therapy to eradicate <i>Helicobacter pylori</i> in a Turkish population. <i>Helicobacter</i> , <b>2012</b> , 17, 486-90	4.9	7
16	Soluble CD40 ligand, soluble P-selectin and von Willebrand factor levels in subjects with prediabetes: the impact of metabolic syndrome. <i>Clinical Biochemistry</i> , <b>2012</b> , 45, 92-5	3.5	11
15	C reactive protein levels in non-alcoholic fatty liver disease. <i>Journal of Hepatology</i> , <b>2012</b> , 56, 507-8; author reply 508-10	13.4	2
14	Elevated asymmetric dimethylarginine in plasma: an early marker for endothelial dysfunction in non-alcoholic fatty liver disease?. <i>Diabetes Research and Clinical Practice</i> , <b>2012</b> , 96, 47-52	7.4	26
13	Circulating vaspin and its relationship with insulin sensitivity, adiponectin, and liver histology in subjects with non-alcoholic steatohepatitis. <i>Scandinavian Journal of Gastroenterology</i> , <b>2011</b> , 46, 1355-61	2.4	12

12	Non-invasive prediction of fibrosis in non-alcoholic fatty liver disease. <i>Journal of Hepatology</i> , <b>2011</b> , 55, 498-9; author reply 499-500	13.4	
11	Plasma apelin and ADMA Levels in patients with essential hypertension. <i>Clinical and Experimental Hypertension</i> , <b>2010</b> , 32, 179-83	2.2	72
10	Mean platelet volume and its relationship with carotid atherosclerosis in subjects with non-alcoholic fatty liver disease. <i>Upsala Journal of Medical Sciences</i> , <b>2010</b> , 115, 253-9	2.8	34
9	Circulating levels of interleukin-18 in patients with non-alcoholic fatty liver disease. <i>Scandinavian Journal of Clinical and Laboratory Investigation</i> , <b>2010</b> , 70, 399-403	2	7
8	Comment and reply on: Serum chemerin and vaspin in non-alcoholic fatty liver disease. <i>Scandinavian Journal of Gastroenterology</i> , <b>2010</b> , 45, 1000-1; author reply 1001-2	2.4	3
7	Chitotriosidase activity predicts endothelial dysfunction in type-2 diabetes mellitus. <i>Endocrine</i> , <b>2010</b> , 37, 455-9	4	25
6	Microalbuminuria and nonalcoholic fatty liver disease. <i>Metabolism: Clinical and Experimental</i> , <b>2010</b> , 59, E5; author reply E6	12.7	
5	Transforming growth factor [TGF- $\beta$ ] levels in otherwise healthy subjects with impaired glucose tolerance. <i>Endokrynologia Polska</i> , <b>2010</b> , 61, 691-4	1.1	
4	LDL-cholesterol lowering increases plasma apelin in isolated hypercholesterolemia. <i>Atherosclerosis</i> , <b>2009</b> , 204, 222-8	3.1	42
3	Platelet aggregation is not enhanced in patients with prediabetes. <i>Upsala Journal of Medical Sciences</i> , <b>2007</b> , 112, 338-46	2.8	1
2	Plasma visfatin levels in patients with newly diagnosed and untreated type 2 diabetes mellitus and impaired glucose tolerance. <i>Diabetes Research and Clinical Practice</i> , <b>2007</b> , 76, 24-9	7.4	149
1	Soluble CD40 ligand levels in otherwise healthy subjects with impaired fasting glucose. <i>Mediators of Inflammation</i> , <b>2006</b> , 2006, 32508	4.3	0