

Jing Li

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2506499/publications.pdf>

Version: 2024-02-01

22
papers

420
citations

933447

10
h-index

752698

20
g-index

22
all docs

22
docs citations

22
times ranked

473
citing authors

#	ARTICLE	IF	CITATIONS
1	Clinicopathological and Prognostic Characteristics in Spinal Chondroblastomas: A Pooled Analysis of Individual Patient Data From a Single Institute and 27 Studies. <i>Global Spine Journal</i> , 2023, 13, 713-723.	2.3	7
2	Letter: A Retrospective Analysis in 1347 Patients Undergoing Cement Augmentation for Osteoporotic Vertebral Compression Fracture: Is the Sandwich Vertebra at a Higher Risk of Further Fracture?. <i>Neurosurgery</i> , 2021, 88, E562-E563.	1.1	1
3	Prognostic Significance of Tumor-Associated Macrophages in Chondroblastoma and Their Association with Response to Adjuvant Radiotherapy. <i>Journal of Inflammation Research</i> , 2021, Volume 14, 1991-2005.	3.5	6
4	Clinicopathological and Prognostic Characteristics in Dedifferentiated/Poorly Differentiated Chordomas: A Pooled Analysis of Individual Patient Data From 58 Studies and Comparison With Conventional Chordomas. <i>Frontiers in Oncology</i> , 2021, 11, 686565.	2.8	8
5	Coexpression of HHLA2 and PD-L1 on Tumor Cells Independently Predicts the Survival of Spinal Chordoma Patients. <i>Frontiers in Immunology</i> , 2021, 12, 797407.	4.8	9
6	Letter: Tumor Growth Rate as a New Predictor of Progression-Free Survival After Chordoma Surgery. <i>Neurosurgery</i> , 2021, Publish Ahead of Print, e19.	1.1	1
7	Development and Validation of a 6-miRNA Prognostic Signature in Spinal Chordoma. <i>Frontiers in Oncology</i> , 2020, 10, 556902.	2.8	9
8	A four-factor immune risk score signature predicts the clinical outcome of patients with spinal chordoma. <i>Clinical and Translational Medicine</i> , 2020, 10, 224-237.	4.0	22
9	Clinical Impact of the Immune Microenvironment in Spinal Chordoma: Immunoscore as an Independent Favorable Prognostic Factor. <i>Neurosurgery</i> , 2019, 84, E318-E333.	1.1	33
10	The Relationship Between Tumor-Stroma Ratio, the Immune Microenvironment, and Survival in Patients With Spinal Chordoma. <i>Neurosurgery</i> , 2019, 85, E1095-E1110.	1.1	29
11	Clinicopathological and Prognostic Characteristics in Extra-Axial Chordomas: An Integrative Analysis of 86 Cases and Comparison With Axial Chordomas. <i>Neurosurgery</i> , 2019, 85, E527-E542.	1.1	13
12	Letter to the Editor. Brachyury as prognostic biomarker in chordoma. <i>Journal of Neurosurgery</i> , 2018, 129, 273-275.	1.6	4
13	Clinicopathologic implications of CD8+/Foxp3+ ratio and miR-574-3p/PD-L1 axis in spinal chordoma patients. <i>Cancer Immunology, Immunotherapy</i> , 2018, 67, 209-224.	4.2	40
14	Prognostic Factors in Skull Base Chordoma: A Systematic Literature Review and Meta-Analysis. <i>World Neurosurgery</i> , 2018, 109, 307-327.	1.3	47
15	Prognostic factors in spinal chordoma: An update of current systematic review and meta-analysis. <i>Journal of Surgical Oncology</i> , 2017, 115, 497-500.	1.7	7
16	Letter: Factors Predicting Recurrence after Resection of Clival Chordoma Using Variable Surgical Approaches and Radiation Modalities. <i>Neurosurgery</i> , 2017, 81, E28-E31.	1.1	6
17	Prognostic Biomarkers in Spinal Chordoma: A Systematic Review. <i>Journal of Neuropathology and Experimental Neurology</i> , 2016, 75, 1184-1187.	1.7	9
18	Upregulated human telomerase reverse transcriptase (hTERT) expression is associated with spinal chordoma growth, invasion and poor prognosis. <i>American Journal of Translational Research (discontinued)</i> , 2016, 8, 516-29.	0.0	17

#	ARTICLE	IF	CITATIONS
19	Expression of programmed death-1 ligand (PD-L1) in tumor-infiltrating lymphocytes is associated with favorable spinal chordoma prognosis. <i>American Journal of Translational Research</i> (discontinued), 2016, 8, 3274-87.	0.0	35
20	Prognostic factors in spinal chordoma: A systematic review. <i>Clinical Neurology and Neurosurgery</i> , 2015, 139, 110-118.	1.4	32
21	Reduced expression of miRNA-1237-3p associated with poor survival of spinal chordoma patients. <i>European Spine Journal</i> , 2015, 24, 1738-1746.	2.2	49
22	Identification of miR-140-3p as a marker associated with poor prognosis in spinal chordoma. <i>International Journal of Clinical and Experimental Pathology</i> , 2014, 7, 4877-85.	0.5	36