

Ying Zhang

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2512377/publications.pdf>

Version: 2024-02-01

10
papers

230
citations

1163117

8
h-index

1372567

10
g-index

10
all docs

10
docs citations

10
times ranked

333
citing authors

#	ARTICLE	IF	CITATIONS
1	Effects of Low-Dose Spironolactone Combined with Metformin or Either Drug Alone on Insulin Resistance in Patients with Polycystic Ovary Syndrome: A Pilot Study. <i>International Journal of Endocrinology</i> , 2022, 2022, 1-8.	1.5	4
2	Effect of metformin and exenatide on pregnancy rate and pregnancy outcomes in overweight or obese infertility PCOS women: long-term follow-up of an RCT. <i>Archives of Gynecology and Obstetrics</i> , 2022, 306, 1711-1721.	1.7	10
3	Circulating zinc- α 2-glycoprotein is reduced in women with polycystic ovary syndrome, but can be increased by exenatide or metformin treatment. <i>Endocrine Journal</i> , 2019, 66, 555-562.	1.6	17
4	Comparison of Oral Antidiabetic Drugs as Add-On Treatments in Patients with Type 2 Diabetes Uncontrolled on Metformin: A Network Meta-Analysis. <i>Diabetes Therapy</i> , 2018, 9, 1945-1958.	2.5	20
5	Serum levels of nesfatin-1 are increased in gestational diabetes mellitus. <i>Gynecological Endocrinology</i> , 2017, 33, 621-624.	1.7	18
6	Efficacy and safety of linagliptin/metformin single-pill combination as initial therapy in drug-naïve Asian patients with type 2 diabetes. <i>Diabetes Research and Clinical Practice</i> , 2017, 124, 48-56.	2.8	13
7	Efficacy of exenatide on weight loss, metabolic parameters and pregnancy in overweight/obese polycystic ovary syndrome. <i>Clinical Endocrinology</i> , 2017, 87, 767-774.	2.4	75
8	Short term monotherapy with exenatide is superior to metformin in weight loss, improving insulin resistance and inflammation in Chinese overweight/obese PCOS women. <i>Obesity Medicine</i> , 2017, 7, 15-20.	0.9	18
9	Changes in serum adipocyte fatty acid-binding protein in women with gestational diabetes mellitus and normal pregnant women during mid- and late pregnancy. <i>Journal of Diabetes Investigation</i> , 2016, 7, 797-804.	2.4	45
10	Clinical study of treatment switching from premixed insulin to basal insulin combined with oral hypoglycemic drugs in patients with type 2 diabetes. <i>Diabetology and Metabolic Syndrome</i> , 2014, 6, 37.	2.7	10