

Vadim A Kuznetsov

List of Publications by Year in descending order

Source: [//exaly.com/author-pdf/2512879/publications.pdf](https://exaly.com/author-pdf/2512879/publications.pdf)

Version: 2024-02-01

73
papers

410
citations

1045038

9
h-index

840818

18
g-index

82
all docs

82
docs citations

82
times ranked

546
citing authors

#	ARTICLE	IF	CITATIONS
1	The Multidimensional Scale of Perceived Social Support (MSPSS): Reliability and Validity of Russian Version. <i>Clinical Gerontologist</i> , 2020, 43, 331-339.	2.3	22
2	INTERRELATION OF TRANSCUTANEOUS CORONARY INTERVENTIONS FOR ACUTE FORMS OF CORONARY HEART DISEASE AND MORTALITY PARAMETERS IN TYUMEN REGION INHABITANTS. <i>Russian Journal of Cardiology</i> , 2014, , 42-46.	1.4	12
3	A LIFE-EXHAUSTION: INTERRELATION WITH THE PREVALENCE OF ISCHEMIC HEART DISEASE. <i>Russian Journal of Cardiology</i> , 2014, , 68-72.	1.4	12
4	SELECTED PARAMETERS OF CHRONIC SOCIAL STRESS IN OPEN POPULATION " ASSOCIATION WITH THE PREVALENCE OF ISCHEMIC HEART DISEASE. <i>Cardiovascular Therapy and Prevention (Russian Federation)</i> , 2014, 13, 28-31.	1.4	11
5	RELIABILITY AND VALIDITY OF RUSSIAN VERSION OF DS14 SCORE FOR ISCHEMIC HEART DISEASE PATIENTS. <i>Russian Journal of Cardiology</i> , 2016, , 50-54.	1.4	10
6	Behavioral risk factors for cardiovascular diseases in men having different work patterns: Results of a cross-sectional epidemiological study. <i>Profilakticheskaya Meditsina</i> , 2016, 19, 49.	0.7	10
7	Associations between high levels of hostility and coronary heart disease in an open urban population among "64-year-old men. <i>Terapevticheskii Arkhiv</i> , 2017, 89, 28-31.	0.8	10
8	Influence of Total Ischemic Time on Clinical Outcomes in Patients with ST-Segment Elevation Myocardial Infarction. <i>Kardiologiya</i> , 2021, 61, 40-46.	1.1	8
9	Impact of hyperglycemia on the results of percutaneous coronary interventions in patients with acute ST-segment elevation myocardial infarction. <i>Terapevticheskii Arkhiv</i> , 2017, 89, 25-29.	0.8	8
10	CLINICAL CHARACTERISTICS OF PATIENTS AND RESULTS OF CATHETER ABLATION IN ATRIAL FIBRILLATION IN RUSSIA: SUBANALYSIS OF THE EUROPEAN REGISTRY 2012-2016. <i>Russian Journal of Cardiology</i> , 2018, , 7-15.	1.4	8
11	Asymmetric septal hypertrophy in patients with coronary artery disease. <i>European Journal of Echocardiography</i> , 2010, 11, 698-702.	2.0	6
12	Social support for patients with coronary artery disease after percutaneous coronary intervention. <i>Journal of Psychosomatic Research</i> , 2019, 119, 74-78.	2.9	6
13	Echocardiographic markers of dyssynchrony as predictors of super-response to cardiac resynchronisation therapy " a pilot study. <i>Cardiovascular Ultrasound</i> , 2018, 16, 24.	1.7	5
14	SOCIO-ECONOMIC RISK FACTORS FOR CARDIOVASCULAR DEATH: DATA FROM 12-YEAR PROSPECTIVE EPIDEMIOLOGIC STUDY. <i>Russian Journal of Cardiology</i> , 2014, , 7-11.	1.4	5
15	Physical activity and social gradient in an open urban population: Gender differences. <i>Profilakticheskaya Meditsina</i> , 2017, 20, 31.	0.7	5
16	ATTITUDE TO PREVENTION OF HEART DISEASES AND STRESS IN THE FAMILY IN THE OPEN CITY POPULATION: PREVALENCE AND INTERRELATIONS. <i>Sibirskij Å¾urnal KliniÄeskoj I ÄksperimentalÉnoj Mediciny</i> , 2019, 33, 148-153.	0.4	5
17	ASSOCIATION BETWEEN CORONARY HEART DISEASE PREVALENCE AND ATTITUDES TOWARDS HEALTH AND PREVENTION IN THE MALE POPULATION OF TUMEN CITY. <i>Cardiovascular Therapy and Prevention (Russian)</i> Tj ETQq14l 0.7843 14 rgBT	0.7843	14
18	Attitude to health and disease prevention of as a parameter of public health. <i>Cardiovascular Therapy and Prevention (Russian Federation)</i> , 2019, 18, 156-160.	1.4	5

#	ARTICLE	IF	CITATIONS
19	Percutaneous coronary interventions for ST elevation myocardial infarction in different age groups. The Siberian Scientific Medical Journal, 2021, 41, 56-65.	0.3	5
20	Comparison of Squatting Stress Echocardiography and Dobutamine Stress Echocardiography for the Diagnosis of Coronary Artery Disease. Echocardiography, 2012, 29, 695-699.	0.9	4
21	Type D personality in Russian patients with cardiovascular disease: validity of the Russian DS14 (DS14-RU). BMC Cardiovascular Disorders, 2019, 19, 78.	1.7	4
22	LONG TERM RESULTS OF PERCUTANEOUS CORONARY INTERVENTIONS COMPARING WITH CONSERVATIVE MANAGEMENT IN TREATMENT OF STABLE ISCHEMIC HEART DISEASE PATIENTS UNDER REAL CIRCUMSTANCES. Russian Journal of Cardiology, 2016, , 7-11.	1.4	4
23	Associations of the prevalence of cardiovascular diseases with a high level of depression in the open population of an average urbanized Siberian town. Terapevticheskii Arkhiv, 2014, 86, 27.	0.8	4
24	Predictors of High-Grade Ventricular Arrhythmias in Patients Referred for the Coronary Angiography. Kardiologiya, 2014, 8_2014, 44-48.	1.1	4
25	Does the lack of left ventricular reverse remodeling always mean non - response to cardiac resynchronization therapy?. Terapevticheskii Arkhiv, 2019, 91, 10-15.	0.8	4
26	The relationship of the prolonged PR interval with the long-term survival in patients with heart failure undergoing cardiac resynchronization therapy. Russian Journal of Cardiology, 2020, 25, 33-38.	1.4	3
27	Association of Atrial Fibrillation with Coronary Lesion in Ischemic Heart Disease Patients. Kardiologiya, 2019, 59, 5-12.	1.1	3
28	Clinical Profile and Treatment of Left Main Coronary Artery Disease in a Real-World Practice. Kardiologiya, 2014, 1_2014, 55-60.	1.1	3
29	Levels of depression and life exhaustion in the open population of the middle urbanized Siberian city: gender differences. Terapevticheskii Arkhiv, 2019, 91, 48-52.	0.8	3
30	Antiarrhythmic drug therapy after atrial fibrillation ablation: data of the ESC-EHRA registry. Russian Journal of Cardiology, 2020, 25, 3874.	1.4	3
31	The risk score for in-hospital mortality in patients with ST-segment elevation myocardial infarction. Kardiologiya, 2021, 61, 11-19.	1.1	3
32	Type D personality in patients with coronary heart disease underwent coronary stenting: a prospective study. Kardiologiya, 2020, 59, 18-24.	1.1	3
33	Gender in cardiac resynchronisation therapy. Journal of Cardiovascular and Thoracic Research, 2018, 10, 197-202.	0.9	2
34	CLINICAL CHARACTERISTICS OF ISCHEMIC HEART DISEASE PATIENTS WITH TYPE D PERSONALITY, UNDERWENT PERCUTANEOUS CORONARY INTERVENTIONS. Cardiovascular Therapy and Prevention (Russian) Tj ETQq0 0 0 rgBT1/0 Overlock 10 Tf 50 1	1.0	2
35	Right ventricular dilatation in patients with coronary heart disease without myocardial infarction: According to the data of the Coronary Angiography Surgery Registry. Terapevticheskii Arkhiv, 2015, 87, 34.	0.8	2
36	Clusters and components of the metabolic syndrome are associations with the prevalence of depression in open urban male population. Cardiovascular Therapy and Prevention (Russian) Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 57 Td	1.0	2

#	ARTICLE	IF	CITATIONS
37	The prevalence of hyperglycemia, as a component of the metabolic syndrome, in men of an open population of Siberian city (epidemiological study). Russian Journal of Cardiology, 2019, , 92-96.	1.4	2
38	Clinical manifestations and risk factors of coronary artery disease in patients with diabetes mellitus in western Siberia. International Journal of Circumpolar Health, 2010, 69, 278-284.	1.3	1
39	Five-year survival and biomarkers of sympatho-adrenal, neurohumoral, immune activation, fibrosis in patients with early and late superresponse to cardiac resynchronization therapy. Journal of Arrhythmology, 2021, 28, 18-27.	0.2	1
40	Overweight in an urban Siberian population: 12-year trends. Cardiovascular Therapy and Prevention (Russian Federation), 2012, 11, 58-61.	1.4	1
41	Populational Characteristics of Metabolic Syndrome in Open Male Population. Kardiologiya, 2014, 6_2014, 58-61.	1.1	1
42	Anxiety, coronary heart disease and metabolic syndrome in male population: prevalence and interrelations. Arterial Hypertension (Russian Federation), 2015, 21, 138-144.	0.4	1
43	Population characteristics of metabolic syndrome components in 25-64-year-old males of an average urbanized Siberian town. Terapevticheskii Arkhiv, 2016, 88, 79.	0.8	1
44	Percutaneous Coronary Intervention in Patients With Acute Myocardial Infarction With ST-Segment Elevation After Prehospital Thrombolysis. Kardiologiya, 2016, 2_2016, 5-10.	1.1	1
45	Primary percutaneous interventions into the coronary arteries in patients with diabetes mellitus. Terapevticheskii Arkhiv, 2016, 88, 35-39.	0.8	1
46	Prediction of superresponse to cardiac resynchronisation therapy in patients with congestive heart failure. Medical Visualization, 2018, , 49-59.	0.4	1
47	ASSOCIATIONS OF SOMATIC RISK FACTORS OF CORONARY HEART DISEASES AND LEVELS OF HOSTILITY OF MEN OF OPEN CITY POPULATION. Sibirskij Å¾urnal KliniÅeskoj I ÅksperimentalÅnoj Mediciny, 2018, 33, 83-89.	0.4	1
48	ULTRASOUND CONTRAST-ENHANCED STRESS ECHOCARDIOGRAPHY WITH MYOCARDIAL PERFUSION FOR FUNCTIONAL ASSESSMENT OF INTERMEDIATE STENOSIS IN THE LEFT ANTERIOR DESCENDING CORONARY ARTERY IN PATIENT WITH PREMATURE VENTRICULAR COMPLEXE. Sibirskij Å¾urnal KliniÅeskoj I ÅksperimentalÅnoj Mediciny, 2019, 34, 137-144.	0.4	1
49	Association of atrial fibrillation with coronary bed lesions (according to the coronary angiography) Tj ETQq1 1 0.784314 rgBT/Overlo	1.4	1
50	Percutaneous Coronary Interventions in Patients With ST-Elevation Myocardial Infarction: 10-Years Follow-up. Kardiologiya, 2020, 60, 69-75.	1.1	1
51	The incidence of frailty and its relationship with long-term survival in patients with heart failure and implanted cardiac resynchronization therapy devices. Russian Journal of Cardiology, 2020, 25, 3685.	1.4	1
52	Association between anxiety and mortality in patients with congestive heart failure after implantation of cardiac electronic devices. Journal of Psychosomatic Research, 2022, 152, 110686.	2.9	1
53	Worldwide geographic differences of patients selected for cardiac resynchronization therapy. European Heart Journal, 2013, 34, P3203-P3203.	2.4	0
54	Decreased sympathoadrenal reactivity during active orthostatic test as a marker of severe chronic heart failure. European Heart Journal, 2013, 34, P5723-P5723.	2.4	0

#	ARTICLE	IF	CITATIONS
55	Attitudes towards the effectiveness of cardiovascular disease prevention and treatment among Tyumen men with and without coronary artery disease. <i>European Heart Journal</i> , 2013, 34, P5146-P5146.	2.4	0
56	P724 Association of biochemical markers with long-term outcomes in patients after coronary angioplasty and stenting. <i>Cardiovascular Research</i> , 2014, 103, S132.3-S132.	3.7	0
57	P200 Biochemical markers in patients with coronary artery disease and diabetes mellitus: prospective observational study. <i>Cardiovascular Research</i> , 2014, 103, S35.4-S36.	3.7	0
58	The new Russian language version of the DS14-RU questionnaire: reliability and validity assessment. <i>Al'manah Kliničeskoj Mediciny</i> , 2021, 49, 113-124.	0.3	0
59	Prediction of 5-year survival in patients with heart failure and implanted cardiac resynchronization therapy devices. <i>Russian Journal of Cardiology</i> , 2021, 26, 4409.	1.4	0
60	Algorithm for assessing the total 10 years risk of death from cardiovascular diseases in women 25-64 years old in Tyumen (Tyumen risk scale). <i>Eurasian Heart Journal</i> , 2021, , 14-21.	0.8	0
61	A Study of the taste sensitivity threshold to the salt as one of arterial hypertension risk factors in schoolchildren of Tyumen. <i>Cardiovascular Therapy and Prevention (Russian Federation)</i> , 2014, , 4-7.	1.4	0
62	The Results of the Method Percutaneous Intracoronary Intervention in Patients of Both Male and Female. <i>Kardiologiya</i> , 2016, 7_2016, 72-77.	1.1	0
63	SOCIAL SUPPORT AS A FACTOR OF CHRONIC SOCIAL STRESS IN MEN OF OPEN CITY POPULATION (BASED ON) Tj ETQq1 1 0,784314 0.4	0.4	0
64	Associations of the spread of low physical activity and working capacity in an open population with and without coronary heart disease. <i>Profilaktičeskaya Meditsina</i> , 2019, 22, 48.	0.7	0
65	Associations between the prevalence of metabolic syndrome and workplace stress in able-bodied males. <i>Profilaktičeskaya Meditsina</i> , 2019, 22, 62.	0.7	0
66	Association of Strategy of Direct Stenting of Infarct-Related Artery with Reduction of Mortality in Woman with ST-Elevation Myocardial Infarction. <i>Kardiologiya</i> , 2019, 59, 5-11.	1.1	0
67	ULTRASOUND CONTRAST-ENHANCED STRESS ECHOCARDIOGRAPHY WITH MYOCARDIAL PERFUSION FOR FUNCTIONAL ASSESSMENT OF INTERMEDIATE STENOSIS IN THE LEFT ANTERIOR DESCENDING CORONARY ARTERY IN PATIENT WITH PREMATURE VENTRICULAR COMPLEXE. <i>Sibirskij Ā¼urnal Kliničeskoj I Ā¼ksperimental'noj Mediciny</i> , 2019, 34, 137-144.	0.4	0
68	FEATURES OF THE ATTITUDE TOWARDS SMOKING IN CONNECTION WITH THE NATURE OF WORK AND MARITAL STATUS IN OPEN URBAN POPULATION: GENDER DIFFERENCES. <i>Sibirskij Ā¼urnal Kliničeskoj I Ā¼ksperimental'noj Mediciny</i> , 2019, 34, 162-169.	0.4	0
69	Ā¼hanges in quality of life in patients with congestive heart failure after cardiac resynchronization therapy. <i>Kardiologiya</i> , 2019, 59, 36-43.	1.1	0
70	Psychosocial risk factors among young people of medium urban Siberian city: a gender aspect (according to a cross-sectional epidemiological study). <i>Cardiovascular Therapy and Prevention (Russian Federation)</i> , 2019, 18, 13-18.	1.4	0
71	Electroanatomic bipolar mapping for detection of arrhythmogenic substrate in catheter ablation of atrial fibrillation. <i>Journal of Arrhythmology</i> , 2020, 26, 32-38.	0.2	0
72	Attitude to health and disease prevention of as a parameter of public health. <i>Cardiovascular Therapy and Prevention (Russian Federation)</i> , 2019, 18, 156-161.	1.4	0

#	ARTICLE	IF	CITATIONS
73	Clinical and demographic characteristics of patients with congestive heart failure and implanted devices for cardiac resynchronization therapy. The Siberian Scientific Medical Journal, 2021, 41, 100-108.	0.3	0