

Satoshi Taba

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2514792/publications.pdf>

Version: 2024-02-01

16

papers

108

citations

1478505

6

h-index

1372567

10

g-index

16

all docs

16

docs citations

16

times ranked

74

citing authors

#	ARTICLE	IF	CITATIONS
1	Nematicidal activity of Okinawa Island plants on the root-knot nematode <i>Meloidogyne incognita</i> (Kofoid and White) Chitwood. <i>Plant and Soil</i> , 2008, 303, 207-216.	3.7	24
2	Alternaria leaf spot of basil caused by <i>Alternaria alternata</i> in Japan. <i>Journal of General Plant Pathology</i> , 2009, 75, 160-162.	1.0	18
3	Identification and use of a wild plant with antimicrobial activity against <i>Ralstonia solanacearum</i> , the cause of bacterial wilt of potato. <i>Weed Biology and Management</i> , 2004, 4, 187-194.	1.4	17
4	Fruit rot of Strawberry pear (pitaya) caused by <i>Bipolaris cactivora</i> . <i>Journal of General Plant Pathology</i> , 2007, 73, 374-376.	1.0	15
5	Black band of Jewâ€™s marrow caused by <i>Lasiodiplodia theobromae</i> . <i>Journal of General Plant Pathology</i> , 2008, 74, 91-93.	1.0	7
6	First report of anthracnose of jaboticaba caused by <i>Colletotrichum tropicale</i> in Japan. <i>Journal of General Plant Pathology</i> , 2020, 86, 65-69.	1.0	6
7	Nematicidal activity of <i>Bidens pilosa</i> var. <i>radiata</i> boiled extracts on plant-parasitic nematodes and stability of the activity. <i>Journal of Weed Science and Technology</i> , 2011, 57, 1-6.	0.1	6
8	<i>Bidens pilosa</i> Extract Effects on Pine Wilt: Causal Agents and Their Natural Enemies. <i>Forest Science</i> , 2020, 66, 284-290.	1.0	5
9	First report of sclerotinia rot of mango caused by <i>Sclerotinia sclerotiorum</i> in Japan. <i>Journal of General Plant Pathology</i> , 2018, 84, 70-72.	1.0	3
10	Efficacy of several control methods on the southern root-knot nematode, <i>Meloidogyne incognita</i>, using <i>Bidens pilosa</i> L. var. <i>radiata</i> Scherff.. <i>Nihon Senchu Gakkai Shi = Japanese Journal of Nematology</i> , 2014, 44, 29-35.	0.3	3
11	Efficacy of a Simple Formulation Composed of Nematode-Trapping Fungi and <i>Bidens pilosa</i> var. <i>radiata</i> Scherff Aqueous Extracts (BPE) for Controlling the Southern Root-Knot Nematode. <i>Microbes and Environments</i> , 2018, 33, 4-9.	1.6	2
12	First report of leaf spot (<i>Corynespora cassiicola</i>) in Assam indigo in Japan. <i>Journal of General Plant Pathology</i> , 2021, 87, 192-195.	1.0	1
13	Ecology of pitaya stem rot caused by <i>Gilbertella persicaria</i> and its biological control by phyllosphere bacteria. <i>Plant Pathology</i> , 0, , .	2.4	1
14	Suppressive effect of polyvinyl alcohol (PVA) film containing <i>Bidens pilosa</i> L. var. <i>radiata</i> aqueous extract on the southern-root knot nematode (<i>Meloidogyne incognita</i>). <i>Nihon Senchu Gakkai Shi = Japanese Journal of Nematology</i> , 2017, 47, 1-8.	0.3	0
15	Development of an environmentally-friendly nematode control agent applying <i>Bidens pilosa</i> L. var. <i>radiata</i> extract. <i>Nihon Senchu Gakkai Shi = Japanese Journal of Nematology</i> , 2020, 50, 39-44.	0.3	0
16	First report of anthracnose caused by <i>Colletotrichum cigarro</i> in pitcher plant in Japan. <i>Journal of General Plant Pathology</i> , 0, , 1.	1.0	0