

Lorenzo Nardo

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2522204/publications.pdf>

Version: 2024-02-01

86
papers

4,613
citations

117625

34
h-index

106344

65
g-index

88
all docs

88
docs citations

88
times ranked

6202
citing authors

#	ARTICLE	IF	CITATIONS
1	Total-Body PET Multiparametric Imaging of Cancer Using a Voxelwise Strategy of Compartmental Modeling. <i>Journal of Nuclear Medicine</i> , 2022, 63, 1274-1281.	5.0	39
2	Potential and Most Relevant Applications of Total Body PET/CT Imaging. <i>Clinical Nuclear Medicine</i> , 2022, 47, 43-55.	1.3	15
3	Total-body PET/CT – First Clinical Experiences and Future Perspectives. <i>Seminars in Nuclear Medicine</i> , 2022, 52, 330-339.	4.6	14
4	Marrow uptake on FDG PET/CT is associated with progression from smoldering to symptomatic multiple myeloma. <i>Skeletal Radiology</i> , 2021, 50, 79-85.	2.0	4
5	Total Body PET: Exploring New Horizons. <i>PET Clinics</i> , 2021, 16, xvii-xviii.	3.0	4
6	Oncologic Applications of Long Axial Field-of-View PET/Computed Tomography. <i>PET Clinics</i> , 2021, 16, 65-73.	3.0	9
7	Clinical Implementation of Total-Body PET/CT at University of California, Davis. <i>PET Clinics</i> , 2021, 16, 1-7.	3.0	19
8	Phase 1 Trial of MLN0128 (Sapanisertib) and CB-839 HCl (Telaglenastat) in Patients With Advanced NSCLC (NCI 10327): Rationale and Study Design. <i>Clinical Lung Cancer</i> , 2021, 22, 67-70.	2.6	33
9	Meaningful words in rectal MRI synoptic reports: How “epolypoid” may be prognostic. <i>Clinical Imaging</i> , 2021, 80, 371-376.	1.5	3
10	Two-bed SPECT/CT versus planar bone scintigraphy: prospective comparison of reproducibility and diagnostic performance. <i>Nuclear Medicine Communications</i> , 2021, 42, 360-368.	1.1	4
11	Venous thromboembolism detected by FDG-PET/CT in cancer patients: a common, yet life-threatening observation. <i>American Journal of Nuclear Medicine and Molecular Imaging</i> , 2021, 11, 99-106.	1.0	4
12	Potential Roles of Total-Body PET/Computed Tomography in Pediatric Imaging. <i>PET Clinics</i> , 2020, 15, 271-279.	3.0	20
13	Qualitative evaluation of MRI features of lipoma and atypical lipomatous tumor: results from a multicenter study. <i>Skeletal Radiology</i> , 2020, 49, 1005-1014.	2.0	24
14	Development and Validation of a Multitask Deep Learning Model for Severity Grading of Hip Osteoarthritis Features on Radiographs. <i>Radiology</i> , 2020, 295, 136-145.	7.3	57
15	Beneficial Effects of Melatonin on Apolipoprotein-E Knockout Mice by Morphological and 18F-FDG PET/CT Assessments. <i>International Journal of Molecular Sciences</i> , 2020, 21, 2920.	4.1	3
16	Radiomics-based prediction of microsatellite instability in colorectal cancer at initial computed tomography evaluation. <i>Abdominal Radiology</i> , 2019, 44, 3755-3763.	2.1	74
17	First Human Imaging Studies with the EXPLORER Total-Body PET Scanner*. <i>Journal of Nuclear Medicine</i> , 2019, 60, 299-303.	5.0	453
18	The Influence of Percentage Weight-Bearing on Foot Radiographs. <i>Foot and Ankle Specialist</i> , 2019, 12, 363-369.	1.0	19

#	ARTICLE	IF	CITATIONS
19	A Deep Learning Model to Predict a Diagnosis of Alzheimer Disease by Using ¹⁸ F-FDG PET of the Brain. <i>Radiology</i> , 2019, 290, 456-464.	7.3	413
20	MRI and biomechanics multidimensional data analysis reveals $R^2 < 1$ as an early predictor of cartilage lesion progression in knee osteoarthritis. <i>Journal of Magnetic Resonance Imaging</i> , 2018, 47, 78-90.	3.4	40
21	Quantitative and Visual Assessments toward Potential Sub-mSv or Ultrafast FDG PET Using High-Sensitivity TOF PET in PET/MRI. <i>Molecular Imaging and Biology</i> , 2018, 20, 492-500.	2.6	12
22	Trabecular bone microstructure is impaired in the proximal femur of human immunodeficiency virus-infected men with normal bone mineral density. <i>Quantitative Imaging in Medicine and Surgery</i> , 2018, 8, 5-13.	2.0	12
23	Vertebral and femoral bone mineral density and bone strength in prostate cancer patients assessed in phantomless PET/CT examinations. <i>Bone</i> , 2017, 101, 62-69.	2.9	28
24	Cyclops lesions detected by MRI are frequent findings after ACL surgical reconstruction but do not impact clinical outcome over 2 years. <i>European Radiology</i> , 2017, 27, 3499-3508.	4.5	25
25	Sporadic Inclusion Body Myositis: MRI Findings and Correlation With Clinical and Functional Parameters. <i>American Journal of Roentgenology</i> , 2017, 209, 1340-1347.	2.2	41
26	Axial or Helical? Considerations for wide collimation CT scanners capable of volumetric imaging in both modes. <i>Medical Physics</i> , 2017, 44, 5718-5725.	3.0	17
27	Cross-sectional and Longitudinal Associations of Diffuse Idiopathic Skeletal Hyperostosis and Thoracic Kyphosis in Older Men and Women. <i>Arthritis Care and Research</i> , 2017, 69, 1245-1252.	3.4	16
28	Tumor immune profiling predicts response to anti-PD-1 therapy in human melanoma. <i>Journal of Clinical Investigation</i> , 2016, 126, 3447-3452.	8.2	439
29	Degeneration in ACL Injured Knees with and without Reconstruction in Relation to Muscle Size and Fat Content—Data from the Osteoarthritis Initiative. <i>PLoS ONE</i> , 2016, 11, e0166865.	2.5	20
30	Impact of Whole-Body Radiation Dose on Response and Toxicity in Patients With Neuroblastoma After Therapy With ¹³¹ I-Metaiodobenzylguanidine (MIBG). <i>Pediatric Blood and Cancer</i> , 2016, 63, 436-442.	1.5	18
31	The Role of PET/CT in the Assessment of Primary Bone Tumors. <i>Current Radiology Reports</i> , 2016, 4, 1.	1.4	2
32	Associations between patellofemoral joint cartilage T1 and T2 and knee flexion moment and impulse during gait in individuals with and without patellofemoral joint osteoarthritis. <i>Osteoarthritis and Cartilage</i> , 2016, 24, 1554-1564.	1.3	26
33	Quantitative assessment of morphology, T1, and T2 of shoulder cartilage using MRI. <i>European Radiology</i> , 2016, 26, 4656-4663.	4.5	10
34	Longitudinal assessment of MRI in hip osteoarthritis using SHOMRI and correlation with clinical progression. <i>Seminars in Arthritis and Rheumatism</i> , 2016, 45, 648-655.	3.4	26
35	Zonal differences in meniscus MR relaxation times in response to in vivo static loading in knee osteoarthritis. <i>Journal of Orthopaedic Research</i> , 2016, 34, 249-261.	2.3	19
36	T1 and T2 relaxation times are associated with progression of hip osteoarthritis. <i>Osteoarthritis and Cartilage</i> , 2016, 24, 1399-1407.	1.3	46

#	ARTICLE	IF	CITATIONS
37	MRI findings associated with development of incident knee pain over 48 months: data from the osteoarthritis initiative. <i>Skeletal Radiology</i> , 2016, 45, 653-660.	2.0	13
38	MR T1 and T2 of meniscus after acute anterior cruciate ligament injuries. <i>Osteoarthritis and Cartilage</i> , 2016, 24, 631-639.	1.3	30
39	Pitfalls in 18 F-FDG PET imaging in gynecological malignancies. <i>Quarterly Journal of Nuclear Medicine and Molecular Imaging</i> , 2016, 60, 124-38.	0.7	1
40	In vitro assessment of knee MRI in the presence of metal implants comparing MAVRIC-SL and conventional fast spin echo sequences at 1.5 and 3 T field strength. <i>Journal of Magnetic Resonance Imaging</i> , 2015, 41, 1291-1299.	3.4	29
41	Scoring hip osteoarthritis with MRI (SHOMRI): A whole joint osteoarthritis evaluation system. <i>Journal of Magnetic Resonance Imaging</i> , 2015, 41, 1549-1557.	3.4	98
42	Association of Physical Activity Measured by Accelerometer, Knee Joint Abnormalities, and Cartilage T2 Measurements Obtained From 3T Magnetic Resonance Imaging: Data From the Osteoarthritis Initiative. <i>Arthritis Care and Research</i> , 2015, 67, 1272-1280.	3.4	36
43	A reference database of cartilage 3T MRI T2 values in knees without diagnostic evidence of cartilage degeneration: data from the osteoarthritis initiative. <i>Osteoarthritis and Cartilage</i> , 2015, 23, 897-905.	1.3	44
44	Femoroacetabular Impingement: Prevalent and Often Asymptomatic in Older Men: The Osteoporotic Fractures in Men Study. <i>Clinical Orthopaedics and Related Research</i> , 2015, 473, 2578-2586.	1.5	32
45	Relationship of unilateral total hip arthroplasty (THA) to contralateral and ipsilateral knee joint degeneration – a longitudinal 3T MRI study from the Osteoarthritis Initiative (OAI). <i>Osteoarthritis and Cartilage</i> , 2015, 23, 1144-1153.	1.3	26
46	Spatial variations in magnetic resonance-based diffusion of articular cartilage in knee osteoarthritis. <i>Magnetic Resonance Imaging</i> , 2015, 33, 1051-1058.	1.8	10
47	Cartilage T1 and T2 Relaxation Times in Patients With Mild to Moderate Radiographic Hip Osteoarthritis. <i>Arthritis and Rheumatology</i> , 2015, 67, 1548-1556.	5.6	34
48	Anatomic correlates of reduced hip extension during walking in individuals with mild to moderate radiographic hip osteoarthritis. <i>Journal of Orthopaedic Research</i> , 2015, 33, 527-534.	2.3	39
49	Association of cartilage degeneration with four year weight gain – 3T MRI data from the Osteoarthritis Initiative. <i>Osteoarthritis and Cartilage</i> , 2015, 23, 525-531.	1.3	42
50	Magnetic resonance rotator cuff fat fraction and its relationship with tendon tear severity and subject characteristics. <i>Journal of Shoulder and Elbow Surgery</i> , 2015, 24, 1442-1451.	2.6	69
51	Are There Sex Differences in Knee Cartilage Composition and Walking Mechanics in Healthy and Osteoarthritis Populations?. <i>Clinical Orthopaedics and Related Research</i> , 2015, 473, 2548-2558.	1.5	29
52	Metal artefact suppression at 3T MRI: comparison of MAVRIC-SL with conventional fast spin echo sequences in patients with Hip joint arthroplasty. <i>European Radiology</i> , 2015, 25, 2403-2411.	4.5	24
53	Femoral condyle insufficiency fractures: associated clinical and morphological findings and impact on outcome. <i>Skeletal Radiology</i> , 2015, 44, 1785-1794.	2.0	27
54	A comparison of melatonin and α -lipoic acid in the induction of antioxidant defences in L6 rat skeletal muscle cells. <i>Age</i> , 2015, 37, 9824.	3.0	18

#	ARTICLE	IF	CITATIONS
55	Metal artifact suppression at the hip: diagnostic performance at 3.0T versus 1.5 Tesla. <i>Skeletal Radiology</i> , 2015, 44, 1609-1616.	2.0	13
56	Diffuse Idiopathic Skeletal Hyperostosis Association With Thoracic Spine Kyphosis. <i>Spine</i> , 2014, 39, E1418-E1424.	2.0	17
57	Thymus and aging: morphological, radiological, and functional overview. <i>Age</i> , 2014, 36, 313-351.	3.0	146
58	Physical Activity and Spatial Differences in Medial Knee T1rho and T2 Relaxation Times in Knee Osteoarthritis. <i>Journal of Orthopaedic and Sports Physical Therapy</i> , 2014, 44, 964-972.	3.5	23
59	Quantitative assessment of fat infiltration in the rotator cuff muscles using water-fat MRI. <i>Journal of Magnetic Resonance Imaging</i> , 2014, 39, 1178-1185.	3.4	88
60	Lesser Tuberosity Avulsions in Adolescents. <i>HSS Journal</i> , 2014, 10, 201-207.	1.7	5
61	PET/MRI Radiotracer Beyond 18F-FDG. <i>PET Clinics</i> , 2014, 9, 345-349.	3.0	5
62	Differences in the Association of Hip Cartilage Lesions and Cam-Type Femoroacetabular Impingement With Movement Patterns: A Preliminary Study. <i>PM and R</i> , 2014, 6, 681-689.	1.6	56
63	Longitudinal evaluation of T1rho and T2 spatial distribution in osteoarthritic and healthy medial knee cartilage. <i>Osteoarthritis and Cartilage</i> , 2014, 22, 51-62.	1.3	45
64	Cartilage Lesion Score: Comparison of a Quantitative Assessment Score with Established Semiquantitative MR Scoring Systems. <i>Radiology</i> , 2014, 271, 479-487.	7.3	43
65	Quadriceps intramuscular fat fraction rather than muscle size is associated with knee osteoarthritis. <i>Osteoarthritis and Cartilage</i> , 2014, 22, 226-234.	1.3	108
66	Magnetic resonance imaging of ankle tendon pathology: benefits of additional axial short-tau inversion recovery imaging to reduce magic angle effects. <i>Skeletal Radiology</i> , 2013, 42, 499-510.	2.0	10
67	Bone marrow changes related to disuse. <i>European Radiology</i> , 2013, 23, 3422-3431.	4.5	11
68	T1rho and T2 relaxation times predict progression of knee osteoarthritis. <i>Osteoarthritis and Cartilage</i> , 2013, 21, 69-76.	1.3	119
69	Association of cartilage defects, and other MRI findings with pain and function in individuals with mild to moderate radiographic hip osteoarthritis and controls. <i>Osteoarthritis and Cartilage</i> , 2013, 21, 1685-1692.	1.3	64
70	Correlation of magnetic resonance imaging-based knee cartilage T2 measurements and focal knee lesions with body mass index: Thirty-six-month followup data from a longitudinal, observational multicenter study. <i>Arthritis Care and Research</i> , 2013, 65, 23-33.	3.4	47
71	Trabecular bone structure and spatial differences in articular cartilage MR relaxation times in individuals with posterior horn medial meniscal tears. <i>Osteoarthritis and Cartilage</i> , 2013, 21, 86-93.	1.3	24
72	Focal knee lesions in knee pairs of asymptomatic and symptomatic subjects with OA risk factors: Data from the Osteoarthritis Initiative. <i>European Journal of Radiology</i> , 2013, 82, e367-e373.	2.6	10

#	ARTICLE	IF	CITATIONS
73	Cartilage morphology and T1 ρ and T2 quantification in ACL-reconstructed knees: a 2-year follow-up. <i>Osteoarthritis and Cartilage</i> , 2013, 21, 1058-1067.	1.3	119
74	Association of Frequent Knee Bending Activity With Focal Knee Lesions Detected With 3T Magnetic Resonance Imaging: Data From the Osteoarthritis Initiative. <i>Arthritis Care and Research</i> , 2013, 65, 1441-1448.	3.4	24
75	Association of Metabolic Risk Factors With Cartilage Degradation Assessed by T2 Relaxation Time at the Knee: Data From the Osteoarthritis Initiative. <i>Arthritis Care and Research</i> , 2013, 65, 1942-1950.	3.4	64
76	T ₂ relaxation time measurements are limited in monitoring progression, once advanced cartilage defects at the knee occur: Longitudinal data from the osteoarthritis initiative. <i>Journal of Magnetic Resonance Imaging</i> , 2013, 38, 1415-1424.	3.4	64
77	Do Cartilage Repair Procedures Prevent Degenerative Meniscus Changes?: Longitudinal T _{1ρ} and Morphological Evaluation With 3.0-T MRI. <i>American Journal of Sports Medicine</i> , 2012, 40, 2700-2708.	4.2	17
78	Lumbosacral Transitional Vertebrae: Association with Low Back Pain. <i>Radiology</i> , 2012, 265, 497-503.	7.3	121
79	Characterization of the regional distribution of skeletal muscle adipose tissue in type 2 diabetes using chemical shift-based water/fat separation. <i>Journal of Magnetic Resonance Imaging</i> , 2012, 35, 899-907.	3.4	103
80	Obesity increases the prevalence and severity of focal knee abnormalities diagnosed using 3T MRI in middle-aged subjects—data from the Osteoarthritis Initiative. <i>Skeletal Radiology</i> , 2012, 41, 633-641.	2.0	78
81	Comparison of clinical semi-quantitative assessment of muscle fat infiltration with quantitative assessment using chemical shift-based water/fat separation in MR studies of the calf of post-menopausal women. <i>European Radiology</i> , 2012, 22, 1592-1600.	4.5	58
82	Baseline mean and heterogeneity of MR cartilage T2 are associated with morphologic degeneration of cartilage, meniscus, and bone marrow over 3years — data from the Osteoarthritis Initiative. <i>Osteoarthritis and Cartilage</i> , 2012, 20, 727-735.	1.3	125
83	Does vertebral bone marrow fat content correlate with abdominal adipose tissue, lumbar spine bone mineral density, and blood biomarkers in women with type 2 diabetes mellitus?. <i>Journal of Magnetic Resonance Imaging</i> , 2012, 35, 117-124.	3.4	196
84	Association of magnetic resonance imaging-based knee cartilage T2 measurements and focal knee lesions with knee pain: Data from the Osteoarthritis Initiative. <i>Arthritis Care and Research</i> , 2012, 64, 248-255.	3.4	96
85	Texture analysis of cartilage T2 maps: individuals with risk factors for OA have higher and more heterogeneous knee cartilage MR T2 compared to normal controls - data from the osteoarthritis initiative. <i>Arthritis Research and Therapy</i> , 2011, 13, R153.	3.5	105
86	Radiology Quiz Case 2. <i>JAMA Otolaryngology</i> , 2011, 137, 629.	1.2	0