

Liselotte E Jensen

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2522928/publications.pdf>

Version: 2024-02-01

30
papers

1,483
citations

279487

23
h-index

476904

29
g-index

30
all docs

30
docs citations

30
times ranked

2328
citing authors

#	ARTICLE	IF	CITATIONS
1	Regulation of serum amyloid A protein expression during the acute-phase response. <i>Biochemical Journal</i> , 1998, 334, 489-503.	1.7	353
2	IL-1 Signaling Cascade in Liver Cells and the Involvement of a Soluble Form of the IL-1 Receptor Accessory Protein. <i>Journal of Immunology</i> , 2000, 164, 5277-5286.	0.4	104
3	The Double-Stranded RNA Analogue Polyinosinic-Polycytidylic Acid Induces Keratinocyte Pyroptosis and Release of IL-36 ¹ . <i>Journal of Investigative Dermatology</i> , 2012, 132, 1346-1353.	0.3	89
4	Mouse model of imiquimod-induced psoriatic itch. <i>Pain</i> , 2016, 157, 2536-2543.	2.0	83
5	Serum amyloid A transcription in Atlantic salmon (<i>Salmo salar</i> L.) hepatocytes is enhanced by stimulation with macrophage factors, recombinant human IL-1 ² , IL-6 and TNF ¹ or bacterial lipopolysaccharide. <i>Developmental and Comparative Immunology</i> , 2000, 24, 553-563.	1.0	68
6	<i>Staphylococcus aureus</i> Stimulates Neutrophil Targeting Chemokine Expression in Keratinocytes through an Autocrine IL-1 ¹ Signaling Loop. <i>Journal of Investigative Dermatology</i> , 2010, 130, 1866-1876.	0.3	64
7	Unprocessed Interleukin-36 ¹ Regulates Psoriasis-Like Skin Inflammation in Cooperation With Interleukin-1. <i>Journal of Investigative Dermatology</i> , 2015, 135, 2992-3000.	0.3	63
8	Pellino3, a Novel Member of the Pellino Protein Family, Promotes Activation of c-Jun and Elk-1 and May Act as a Scaffolding Protein. <i>Journal of Immunology</i> , 2003, 171, 1500-1506.	0.4	62
9	IL-1R1 Signaling Facilitates <i>Munro</i> 's Microabscess Formation in Psoriasiform Imiquimod-Induced Skin Inflammation. <i>Journal of Investigative Dermatology</i> , 2013, 133, 1541-1549.	0.3	61
10	Pellino2 activates the mitogen activated protein kinase pathway. <i>FEBS Letters</i> , 2003, 545, 199-202.	1.3	45
11	Pellino Proteins Contain a Cryptic FHA Domain that Mediates Interaction with Phosphorylated IRAK1. <i>Structure</i> , 2008, 16, 1806-1816.	1.6	44
12	Interleukin-1 regulates keratinocyte expression of T cell targeting chemokines through interleukin-1 receptor associated kinase-1 (IRAK1) dependent and independent pathways. <i>Cellular Signalling</i> , 2009, 21, 685-694.	1.7	39
13	Loss of function polymorphisms in NAT1 protect against spina bifida. <i>Human Genetics</i> , 2006, 120, 52-57.	1.8	35
14	Maternal genotype for the monocyte chemoattractant protein 1 A(-2518)G promoter polymorphism is associated with the risk of spina bifida in offspring. <i>American Journal of Medical Genetics, Part A</i> , 2006, 140A, 1114-1118.	0.7	35
15	Interleukin-1 ¹ released from HSV-1-infected keratinocytes acts as a functional alarmin in the skin. <i>Nature Communications</i> , 2014, 5, 5230.	5.8	35
16	Expression of alternatively spliced interleukin-1 receptor accessory protein mRNAs is differentially regulated during inflammation and apoptosis. <i>Cellular Signalling</i> , 2003, 15, 793-802.	1.7	33
17	Targeting the IL-1 family members in skin inflammation. <i>Current Opinion in Investigational Drugs</i> , 2010, 11, 1211-20.	2.3	33
18	Imiquimod Treatment Causes Systemic Disease in Mice Resembling Generalized Pustular Psoriasis in an IL-1 and IL-36 Dependent Manner. <i>Mediators of Inflammation</i> , 2016, 2016, 1-10.	1.4	29

#	ARTICLE	IF	CITATIONS
19	Ubiquitin activated tumor necrosis factor receptor associated factor-6 (TRAF6) is recycled via deubiquitination. <i>FEBS Letters</i> , 2003, 553, 190-194.	1.3	28
20	The NAT1 C1095A polymorphism, maternal multivitamin use and smoking, and the risk of spina bifida. <i>Birth Defects Research Part A: Clinical and Molecular Teratology</i> , 2005, 73, 512-516.	1.6	28
21	The human T locus and spina bifida risk. <i>Human Genetics</i> , 2004, 115, 475-482.	1.8	27
22	Interleukin-36 cytokines may overcome microbial immune evasion strategies that inhibit interleukin-1 family signaling. <i>Science Signaling</i> , 2017, 10, .	1.6	26
23	Interleukin-36 β provides protection against HSV-1 infection, but does not modulate initiation of adaptive immune responses. <i>Scientific Reports</i> , 2017, 7, 5799.	1.6	24
24	The 3' UTR Untranslated Region of the Membrane-Bound IL-1R Accessory Protein mRNA Confers Tissue-Specific Destabilization. <i>Journal of Immunology</i> , 2004, 173, 6248-6258.	0.4	23
25	IL-36 promotes anti-viral immunity by boosting sensitivity to IFN- β in IRF1 dependent and independent manners. <i>Nature Communications</i> , 2019, 10, 4700.	5.8	23
26	ELAM-1/E-selectin promoter contains an inducible AP-1/CREB site and is not NF- κ B-specific. <i>BioTechniques</i> , 2003, 35, 54-58.	0.8	13
27	A common ABCC2 promoter polymorphism is not a determinant of the risk of spina bifida. <i>Birth Defects Research Part A: Clinical and Molecular Teratology</i> , 2004, 70, 396-399.	1.6	9
28	Interleukin-36 (IL-36) system in the 1-fluoro-2,4-dinitrobenzene (DNFB) mouse model of allergic contact dermatitis. <i>Allergy: European Journal of Allergy and Clinical Immunology</i> , 2020, 75, 2078-2081.	2.7	6
29	Transient Induction of Fever in the Imiquimod C57BL/6 Mouse Model of Psoriasis-Like Disease Involves IL-1 and IL-6 but Not IL-36. <i>Journal of Investigative Dermatology</i> , 2022, 142, 247-251.e2.	0.3	1
30	Interleukin-1 facilitates recruitment of leukocytes into the skin through a complex assortment of chemokines. <i>FASEB Journal</i> , 2008, 22, 1070.19.	0.2	0