

# Liselotte E Jensen

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2522928/publications.pdf>

Version: 2024-02-01

30  
papers

1,483  
citations

279487

23  
h-index

476904

29  
g-index

30  
all docs

30  
docs citations

30  
times ranked

2328  
citing authors

#	ARTICLE	IF	CITATIONS
1	Transient Induction of Fever in the Imiquimod C57BL/6 Mouse Model of Psoriasis-Like Disease Involves IL-1 and IL-6 but Not IL-36. <i>Journal of Investigative Dermatology</i> , 2022, 142, 247-251.e2.	0.3	1
2	Interleukin-36 (IL-36) system in the 1-fluoro-2,4-dinitrobenzene (DNFB) mouse model of allergic contact dermatitis. <i>Allergy: European Journal of Allergy and Clinical Immunology</i> , 2020, 75, 2078-2081.	2.7	6
3	IL-36 promotes anti-viral immunity by boosting sensitivity to IFN- $\beta$ in IRF1 dependent and independent manners. <i>Nature Communications</i> , 2019, 10, 4700.	5.8	23
4	Interleukin-36 $\beta$ provides protection against HSV-1 infection, but does not modulate initiation of adaptive immune responses. <i>Scientific Reports</i> , 2017, 7, 5799.	1.6	24
5	Interleukin-36 cytokines may overcome microbial immune evasion strategies that inhibit interleukin-1 family signaling. <i>Science Signaling</i> , 2017, 10, .	1.6	26
6	Imiquimod Treatment Causes Systemic Disease in Mice Resembling Generalized Pustular Psoriasis in an IL-1 and IL-36 Dependent Manner. <i>Mediators of Inflammation</i> , 2016, 2016, 1-10.	1.4	29
7	Mouse model of imiquimod-induced psoriatic itch. <i>Pain</i> , 2016, 157, 2536-2543.	2.0	83
8	Unprocessed Interleukin-36 $\beta$ Regulates Psoriasis-Like Skin Inflammation in Cooperation With Interleukin-1. <i>Journal of Investigative Dermatology</i> , 2015, 135, 2992-3000.	0.3	63
9	Interleukin-1 $\beta$ released from HSV-1-infected keratinocytes acts as a functional alarmin in the skin. <i>Nature Communications</i> , 2014, 5, 5230.	5.8	35
10	IL-1R1 Signaling Facilitates Munro's Microabscess Formation in Psoriasiform Imiquimod-Induced Skin Inflammation. <i>Journal of Investigative Dermatology</i> , 2013, 133, 1541-1549.	0.3	61
11	The Double-Stranded RNA Analogue Polyinosinic-Polycytidylic Acid Induces Keratinocyte Pyroptosis and Release of IL-36 $\beta$ . <i>Journal of Investigative Dermatology</i> , 2012, 132, 1346-1353.	0.3	89
12	Staphylococcus aureus Stimulates Neutrophil Targeting Chemokine Expression in Keratinocytes through an Autocrine IL-1 $\beta$ Signaling Loop. <i>Journal of Investigative Dermatology</i> , 2010, 130, 1866-1876.	0.3	64
13	Targeting the IL-1 family members in skin inflammation. <i>Current Opinion in Investigational Drugs</i> , 2010, 11, 1211-20.	2.3	33
14	Interleukin-1 regulates keratinocyte expression of T cell targeting chemokines through interleukin-1 receptor associated kinase-1 (IRAK1) dependent and independent pathways. <i>Cellular Signalling</i> , 2009, 21, 685-694.	1.7	39
15	Pellino Proteins Contain a Cryptic FHA Domain that Mediates Interaction with Phosphorylated IRAK1. <i>Structure</i> , 2008, 16, 1806-1816.	1.6	44
16	Interleukin-1 facilitates recruitment of leukocytes into the skin through a complex assortment of chemokines. <i>FASEB Journal</i> , 2008, 22, 1070.19.	0.2	0
17	Loss of function polymorphisms in NAT1 protect against spina bifida. <i>Human Genetics</i> , 2006, 120, 52-57.	1.8	35
18	Maternal genotype for the monocyte chemoattractant protein 1 A(-2518)G promoter polymorphism is associated with the risk of spina bifida in offspring. <i>American Journal of Medical Genetics, Part A</i> , 2006, 140A, 1114-1118.	0.7	35

#	ARTICLE	IF	CITATIONS
19	The NAT1 C1095A polymorphism, maternal multivitamin use and smoking, and the risk of spina bifida. Birth Defects Research Part A: Clinical and Molecular Teratology, 2005, 73, 512-516.	1.6	28
20	The 3' UTR Untranslated Region of the Membrane-Bound IL-1R Accessory Protein mRNA Confers Tissue-Specific Destabilization. Journal of Immunology, 2004, 173, 6248-6258.	0.4	23
21	The human T locus and spina bifida risk. Human Genetics, 2004, 115, 475-482.	1.8	27
22	A common ABCC2 promoter polymorphism is not a determinant of the risk of spina bifida. Birth Defects Research Part A: Clinical and Molecular Teratology, 2004, 70, 396-399.	1.6	9
23	Expression of alternatively spliced interleukin-1 receptor accessory protein mRNAs is differentially regulated during inflammation and apoptosis. Cellular Signalling, 2003, 15, 793-802.	1.7	33
24	Pellino2 activates the mitogen activated protein kinase pathway. FEBS Letters, 2003, 545, 199-202.	1.3	45
25	Ubiquitin activated tumor necrosis factor receptor associated factor-6 (TRAF6) is recycled via deubiquitination. FEBS Letters, 2003, 553, 190-194.	1.3	28
26	Pellino3, a Novel Member of the Pellino Protein Family, Promotes Activation of c-Jun and Elk-1 and May Act as a Scaffolding Protein. Journal of Immunology, 2003, 171, 1500-1506.	0.4	62
27	ELAM-1/E-selectin promoter contains an inducible AP-1/CREB site and is not NF- $\kappa$ B-specific. BioTechniques, 2003, 35, 54-58.	0.8	13
28	IL-1 Signaling Cascade in Liver Cells and the Involvement of a Soluble Form of the IL-1 Receptor Accessory Protein. Journal of Immunology, 2000, 164, 5277-5286.	0.4	104
29	Serum amyloid A transcription in Atlantic salmon ( <i>Salmo salar</i> L.) hepatocytes is enhanced by stimulation with macrophage factors, recombinant human IL-1 $\beta$ , IL-6 and TNF $\alpha$ or bacterial lipopolysaccharide. Developmental and Comparative Immunology, 2000, 24, 553-563.	1.0	68
30	Regulation of serum amyloid A protein expression during the acute-phase response. Biochemical Journal, 1998, 334, 489-503.	1.7	353