Eva Grill

List of Publications by Citations

Source: https://exaly.com/author-pdf/2525749/eva-grill-publications-by-citations.pdf

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

183
papers3,051
citations30
h-index43
g-index214
ext. papers3,718
ext. citations3.8
avg, IF5.23
L-index

#	Paper	IF	Citations
183	Rationale and principles of early rehabilitation care after an acute injury or illness. <i>Disability and Rehabilitation</i> , 2005 , 27, 353-9	2.4	107
182	ICF Core Sets development for the acute hospital and early post-acute rehabilitation facilities. <i>Disability and Rehabilitation</i> , 2005 , 27, 361-6	2.4	101
181	Incidence and risk factors of respiratory syncytial virus-related hospitalizations in premature infants in Germany. <i>European Journal of Pediatrics</i> , 2003 , 162, 230-6	4.1	90
180	ICF Core Set for patients with neurological conditions in early post-acute rehabilitation facilities. <i>Disability and Rehabilitation</i> , 2005 , 27, 389-95	2.4	72
179	Health services utilization of patients with vertigo in primary care: a retrospective cohort study. Journal of Neurology, 2014 , 261, 1492-8	5.5	68
178	Rehabilitation outcome of anoxic-ischaemic encephalopathy survivors with prolonged disorders of consciousness. <i>Resuscitation</i> , 2013 , 84, 1409-15	4	67
177	Burden of disability attributable to vertigo and dizziness in the aged: results from the KORA-Age study. <i>European Journal of Public Health</i> , 2014 , 24, 802-7	2.1	66
176	Economic evaluation of possible prevention of RSV-related hospitalizations in premature infants in Germany. <i>European Journal of Pediatrics</i> , 2003 , 162, 237-44	4.1	56
175	ICF Core Set for patients with neurological conditions in the acute hospital. <i>Disability and Rehabilitation</i> , 2005 , 27, 367-73	2.4	55
174	Down on heights? One in three has visual height intolerance. <i>Journal of Neurology</i> , 2013 , 260, 597-604	5.5	53
173	How much does it hurt to be lonely? Mental and physical differences between older men and women in the KORA-Age Study. <i>International Journal of Geriatric Psychiatry</i> , 2014 , 29, 245-52	3.9	52
172	ICF Core Set for geriatric patients in early post-acute rehabilitation facilities. <i>Disability and Rehabilitation</i> , 2005 , 27, 411-7	2.4	50
171	Association between sleep disturbances and falls among the elderly: results from the German Cooperative Health Research in the Region of Augsburg-Age study. <i>Sleep Medicine</i> , 2013 , 14, 1356-63	4.6	46
170	Gender differences in the association between grip strength and mortality in older adults: results from the KORA-age study. <i>BMC Geriatrics</i> , 2016 , 16, 201	4.1	46
169	Patient goals in post-acute geriatric rehabilitation-goal attainment is an indicator for improved functioning. <i>Journal of Rehabilitation Medicine</i> , 2011 , 43, 156-61	3.4	43
168	Systematic review of measures and their concepts used in published studies focusing on rehabilitation in the acute hospital and in early post-acute rehabilitation facilities. <i>Disability and Rehabilitation</i> , 2005 , 27, 419-29	2.4	43
167	Scales could be developed based on simple clinical ratings of International Classification of Functioning, Disability and Health Core Set categories. <i>Journal of Clinical Epidemiology</i> , 2009 , 62, 891-8	5.7	42

(2005-2015)

166	Persistent vegetative state and minimally conscious state: a systematic review and meta-analysis of diagnostic procedures. <i>Deutsches A&#x0308;rzteblatt International</i> , 2015 , 112, 235-42	2.5	40
165	Health outcome after major trauma: what are we measuring?. PLoS ONE, 2014 , 9, e103082	3.7	38
164	Description of spatio-temporal gait parameters in elderly people and their association with history of falls: results of the population-based cross-sectional KORA-Age study. <i>BMC Geriatrics</i> , 2015 , 15, 32	4.1	37
163	Distribution and determinants of functioning and disability in aged adultsresults from the German KORA-Age study. <i>BMC Public Health</i> , 2013 , 13, 137	4.1	36
162	Health care utilization, prognosis and outcomes of vestibular disease in primary care settings: systematic review. <i>Journal of Neurology</i> , 2016 , 263 Suppl 1, S36-44	5.5	36
161	Identification of relevant ICF categories by patients with neurological conditions in early post-acute rehabilitation facilities. <i>Disability and Rehabilitation</i> , 2005 , 27, 459-65	2.4	35
160	Assessing observer agreement when describing and classifying functioning with the International Classification of Functioning, Disability and Health. <i>Acta Dermato-Venereologica</i> , 2007 , 39, 71-6	2.2	34
159	ICF Core Set for patients with cardiopulmonary conditions in early post-acute rehabilitation facilities. <i>Disability and Rehabilitation</i> , 2005 , 27, 397-404	2.4	32
158	Identification of relevant ICF categories by patients in the acute hospital. <i>Disability and Rehabilitation</i> , 2005 , 27, 447-58	2.4	32
157	Time to evaluate COVID-19 contact-tracing apps. <i>Nature Medicine</i> , 2021 , 27, 361-362	50.5	32
157 156	Time to evaluate COVID-19 contact-tracing apps. <i>Nature Medicine</i> , 2021 , 27, 361-362 Understanding human functioning using graphical models. <i>BMC Medical Research Methodology</i> , 2010 , 10, 14	50.5 4·7	31
	Understanding human functioning using graphical models. BMC Medical Research Methodology,		
156	Understanding human functioning using graphical models. <i>BMC Medical Research Methodology</i> , 2010 , 10, 14 Visual height intolerance and acrophobia: clinical characteristics and comorbidity patterns.	4.7	31
156 155	Understanding human functioning using graphical models. <i>BMC Medical Research Methodology</i> , 2010 , 10, 14 Visual height intolerance and acrophobia: clinical characteristics and comorbidity patterns. <i>European Archives of Psychiatry and Clinical Neuroscience</i> , 2015 , 265, 375-85 Contractures with special reference in elderly: definition and risk factors - a systematic review with	4.7	31
156 155 154	Understanding human functioning using graphical models. <i>BMC Medical Research Methodology</i> , 2010 , 10, 14 Visual height intolerance and acrophobia: clinical characteristics and comorbidity patterns. <i>European Archives of Psychiatry and Clinical Neuroscience</i> , 2015 , 265, 375-85 Contractures with special reference in elderly: definition and risk factors - a systematic review with practical implications. <i>Disability and Rehabilitation</i> , 2014 , 36, 529-38 International Classification of Functioning, Disability and Health (ICF) Core Set for patients with vertigo, dizziness and balance disorders. <i>Journal of Vestibular Research: Equilibrium and Orientation</i> ,	4·7 5.1 2·4	31 30 30
156 155 154	Understanding human functioning using graphical models. <i>BMC Medical Research Methodology</i> , 2010 , 10, 14 Visual height intolerance and acrophobia: clinical characteristics and comorbidity patterns. <i>European Archives of Psychiatry and Clinical Neuroscience</i> , 2015 , 265, 375-85 Contractures with special reference in elderly: definition and risk factors - a systematic review with practical implications. <i>Disability and Rehabilitation</i> , 2014 , 36, 529-38 International Classification of Functioning, Disability and Health (ICF) Core Set for patients with vertigo, dizziness and balance disorders. <i>Journal of Vestibular Research: Equilibrium and Orientation</i> , 2012 , 22, 261-71 The ICF forms a useful framework for classifying individual patient goals in post-acute	4·7 5.1 2·4 2.5	31 30 30 30
156 155 154 153	Understanding human functioning using graphical models. <i>BMC Medical Research Methodology</i> , 2010 , 10, 14 Visual height intolerance and acrophobia: clinical characteristics and comorbidity patterns. <i>European Archives of Psychiatry and Clinical Neuroscience</i> , 2015 , 265, 375-85 Contractures with special reference in elderly: definition and risk factors - a systematic review with practical implications. <i>Disability and Rehabilitation</i> , 2014 , 36, 529-38 International Classification of Functioning, Disability and Health (ICF) Core Set for patients with vertigo, dizziness and balance disorders. <i>Journal of Vestibular Research: Equilibrium and Orientation</i> , 2012 , 22, 261-71 The ICF forms a useful framework for classifying individual patient goals in post-acute rehabilitation. <i>Journal of Rehabilitation Medicine</i> , 2011 , 43, 151-5	4·7 5.1 2·4 2·5	31 30 30 30 30

148	Impact of vertigo and dizziness on self-perceived participation and autonomy in older adults: results from the KORA-Age study. <i>Quality of Life Research</i> , 2014 , 23, 2301-8	3.7	27
147	On the relationship between gingival biotypes and supracrestal gingival height, crown form and papilla height. <i>Clinical Oral Implants Research</i> , 2014 , 25, 894-8	4.8	27
146	Graphical models illustrated complex associations between variables describing human functioning. <i>Journal of Clinical Epidemiology</i> , 2009 , 62, 922-33	5.7	27
145	Economic burden of vertigo: a systematic review. <i>Health Economics Review</i> , 2019 , 9, 37	2	27
144	Validation of International Classification of Functioning, Disability, and Health (ICF) Core Sets for early postacute rehabilitation facilities: comparisons with three other functional measures. American Journal of Physical Medicine and Rehabilitation, 2006, 85, 640-9	2.6	26
143	Rehabilitation outcome of unconscious traumatic brain injury patients. <i>Journal of Neurotrauma</i> , 2013 , 30, 1476-83	5.4	24
142	Prevalence and Predictors of Subclinical Micronutrient Deficiency in German Older Adults: Results from the Population-Based KORA-Age Study. <i>Nutrients</i> , 2017 , 9,	6.7	24
141	Neighborhood and healthy aging in a German city: distances to green space and senior service centers and their associations with physical constitution, disability, and health-related quality of life. <i>European Journal of Ageing</i> , 2015 , 12, 273-283	3.6	23
140	The ICF categories identified in nursing interventions administered to neurological patients with post-acute rehabilitation needs. <i>Disability and Rehabilitation</i> , 2005 , 27, 431-6	2.4	23
139	Country-specific and individual determinants of dizziness in Europe: results from the Survey of Health Ageing and Retirement in Europe (SHARE). <i>Public Health</i> , 2017 , 149, 1-10	4	22
138	Systematic Review and Meta-analysis of the Effectiveness of Implementation Strategies for Non-communicable Disease Guidelines in Primary Health Care. <i>Journal of General Internal Medicine</i> , 2018 , 33, 1142-1154	4	22
137	Sarcopenia is associated with disability status-results from the KORA-Age study. <i>Osteoporosis International</i> , 2017 , 28, 2069-2079	5.3	21
136	A New Questionnaire for Estimating the Severity of Visual Height Intolerance and Acrophobia by a Metric Interval Scale. <i>Frontiers in Neurology</i> , 2017 , 8, 211	4.1	21
135	Associations between multiple accelerometry-assessed physical activity parameters and selected health outcomes in elderly peopleresults from the KORA-age study. <i>PLoS ONE</i> , 2014 , 9, e111206	3.7	21
134	Validation of the comprehensive ICF Core Sets for patients receiving rehabilitation interventions in the acute care setting. <i>Journal of Rehabilitation Medicine</i> , 2011 , 43, 92-101	3.4	21
133	Goals of physiotherapy interventions can be described using the International Classification of Functioning, Disability and Health. <i>Physiotherapy</i> , 2008 , 94, 150-157	3	21
132	Neonatal hearing screening: modelling cost and effectiveness of hospital- and community-based screening. <i>BMC Health Services Research</i> , 2006 , 6, 14	2.9	21
131	Consequences of visual height intolerance for quality of life: a qualitative study. <i>Quality of Life Research</i> , 2014 , 23, 697-705	3.7	20

(2016-2015)

130	Subscales of the vestibular activities and participation questionnaire could be applied across cultures. <i>Journal of Clinical Epidemiology</i> , 2015 , 68, 211-9	5.7	20
129	Comparing the clinical effectiveness of different new-born hearing screening strategies. A decision analysis. <i>BMC Public Health</i> , 2005 , 5, 12	4.1	20
128	Functioning and disability in people living with spinal cord injury in high- and low-resourced countries: a comparative analysis of 14 countries. <i>International Journal of Public Health</i> , 2011 , 56, 341-56.	52 ⁴	19
127	Rationale and design of the prospective German registry of outcome in patients with severe disorders of consciousness after acute brain injury. <i>Archives of Physical Medicine and Rehabilitation</i> , 2013 , 94, 1870-6	2.8	18
126	The ICF as a way to specify goals and to assess the outcome of physiotherapeutic interventions in the acute hospital. <i>Journal of Rehabilitation Medicine</i> , 2011 , 43, 174-7	3.4	18
125	Identification of relevant ICF categories by health professionals in the acute hospital. <i>Disability and Rehabilitation</i> , 2005 , 27, 437-45	2.4	18
124	Determinants for utilization and transitions of long-term care in adults 65+ in Germany: results from the longitudinal KORA-Age study. <i>BMC Geriatrics</i> , 2018 , 18, 172	4.1	17
123	Validation of the comprehensive ICF Core Set for patients in geriatric post-acute rehabilitation facilities. <i>Journal of Rehabilitation Medicine</i> , 2011 , 43, 102-12	3.4	17
122	Criteria for validating comprehensive ICF Core Sets and developing brief ICF Core Set versions. Journal of Rehabilitation Medicine, 2011 , 43, 87-91	3.4	17
121	Goals of patients with rehabilitation needs in acute hospitals: goal achivement is an indicator for improved functioning. <i>Journal of Rehabilitation Medicine</i> , 2011 , 43, 145-50	3.4	17
120	Prevalence, Determinants, and Consequences of Vestibular Hypofunction. Results From the KORA-FF4 Survey. <i>Frontiers in Neurology</i> , 2018 , 9, 1076	4.1	17
119	Intervention goals determine physical therapistsQworkload in the acute care setting. <i>Physical Therapy</i> , 2010 , 90, 1468-78	3.3	16
118	Patients Qunctioning as predictor of nursing workload in acute hospital units providing rehabilitation care: a multi-centre cohort study. <i>BMC Health Services Research</i> , 2010 , 10, 295	2.9	16
117	Patients Qview on health-related aspects of functioning and disability of joint contractures: a qualitative interview study based on the International Classification of Functioning, Disability and Health (ICF). <i>Disability and Rehabilitation</i> , 2014 , 36, 2225-32	2.4	15
116	Brief ICF Core Set for patients in geriatric post-acute rehabilitation facilities. <i>Journal of Rehabilitation Medicine</i> , 2011 , 43, 139-44	3.4	15
115	Survey of motion sickness susceptibility in children and adolescents aged 3 months to 18 years. Journal of Neurology, 2019 , 266, 65-73	5.5	14
114	Impact of joint contracture on older persons in a geriatric setting: A cross-sectional study. <i>Zeitschrift Fur Gerontologie Und Geriatrie</i> , 2015 , 48, 625-32	2.7	13
113	Relevance of community structures and neighbourhood characteristics for participation of older adults: a qualitative study. <i>Quality of Life Research</i> , 2016 , 25, 143-52	3.7	13

112	Association between anemia and falls in community-dwelling older people: cross-sectional results from the KORA-Age study. <i>BMC Geriatrics</i> , 2014 , 14, 29	4.1	13
111	ICF Core Sets for early post-acute rehabilitation facilities. <i>Journal of Rehabilitation Medicine</i> , 2011 , 43, 131-8	3.4	13
110	Towards a European strategy to address the COVID-19 pandemic. Lancet, The, 2021, 398, 838-839	40	13
109	Examining Functioning and Contextual Factors in Individuals with Joint Contractures from the Health Professional Perspective Using the ICF: An International Internet-Based Qualitative Expert Survey. <i>Rehabilitation Nursing</i> , 2016 , 41, 170-8	1.3	12
108	Predictive Capability of an iPad-Based Medical Device (mex) for the Diagnosis of Vertigo and Dizziness. <i>Frontiers in Neurology</i> , 2018 , 9, 29	4.1	12
107	Effects of pharmaceutical counselling on antimicrobial use in surgical wards: intervention study with historical control group. <i>Pharmacoepidemiology and Drug Safety</i> , 2011 , 20, 739-46	2.6	12
106	Functioning and health in patients with cancer on home-parenteral nutrition: a qualitative study. Health and Quality of Life Outcomes, 2010 , 8, 41	3	12
105	Systematic review of measures and their concepts used in published studies focusing on the treatment of acute inflammatory arthritis. <i>Clinical Rheumatology</i> , 2006 , 25, 807-13	3.9	12
104	Worldwide survey on laboratory testing of vestibular function. <i>Neurology: Clinical Practice</i> , 2020 , 10, 379-387	1.7	12
103	Primary or secondary chronic functional dizziness: does it make a difference? A DizzyReg study in 356 patients. <i>Journal of Neurology</i> , 2020 , 267, 212-222	5.5	12
102	Common eye diseases in older adults of southern Germany: results from the KORA-Age study. <i>Age and Ageing</i> , 2017 , 46, 481-486	3	12
101	Susceptibility to Fear of Heights in Bilateral Vestibulopathy and Other Disorders of Vertigo and Balance. <i>Frontiers in Neurology</i> , 2018 , 9, 406	4.1	11
100	Directed acyclic graphs helped to identify confounding in the association of disability and electrocardiographic findings: results from the KORA-Age study. <i>Journal of Clinical Epidemiology</i> , 2014 , 67, 199-206	5.7	11
99	Quality of evidence of rehabilitation interventions in long-term care for people with severe disorders of consciousness after brain injury: A systematic review. <i>Journal of Rehabilitation Medicine</i> , 2015 , 47, 577-85	3.4	11
98	A high level of household physical activity compensates for lack of leisure time physical activity with regard to deficit accumulation: Results from the KORA-Age study. <i>Preventive Medicine</i> , 2016 , 86, 64-9	4.3	11
97	Prevalence of motion sickness in various vestibular disorders: a study on 749 patients. <i>Journal of Neurology</i> , 2018 , 265, 95-97	5.5	10
96	Impact of joint contractures on functioning and social participation in older individualsdevelopment of a standard set (JointConFunctionSet): study protocol. <i>BMC Geriatrics</i> , 2013 , 13, 18	4.1	10
95	Efficacy of a post-acute interval inpatient neurorehabilitation programme for severe brain injury. <i>Brain Injury</i> , 2014 , 28, 44-50	2.1	10

(2016-2013)

94	Using core sets of the international classification of functioning, disability and health (ICF) to measure disability in vestibular disorders: study protocol. <i>Journal of Vestibular Research: Equilibrium and Orientation</i> , 2013 , 23, 297-303	2.5	10
93	Assessment of functioning in the acute hospital: operationalisation and reliability testing of ICF categories relevant for physical therapists interventions. <i>Journal of Rehabilitation Medicine</i> , 2011 , 43, 162-73	3.4	10
92	Combined application of the International Classification of Functioning, Disability and Health and the NANDA-International Taxonomy II. <i>Journal of Advanced Nursing</i> , 2010 , 66, 1885-98	3.1	10
91	Identification of ICF categories relevant for nursing in the situation of acute and early post-acute rehabilitation. <i>BMC Nursing</i> , 2008 , 7, 3	3.2	10
90	Malnutrition and related risk factors in older adults from different health-care settings: an study. <i>Public Health Nutrition</i> , 2020 , 23, 446-456	3.3	10
89	DizzyReg: the prospective patient registry of the German Center for Vertigo and Balance Disorders. Journal of Neurology, 2017 , 264, 34-36	5.5	9
88	Vitamin D in Relation to Incident Sarcopenia and Changes in Muscle Parameters Among Older Adults: The KORA-Age Study. <i>Calcified Tissue International</i> , 2019 , 105, 173-182	3.9	9
87	Modern machine-learning can support diagnostic differentiation of central and peripheral acute vestibular disorders. <i>Journal of Neurology</i> , 2020 , 267, 143-152	5.5	9
86	Barriers and facilitators for the management of vertigo: a qualitative study with primary care providers. <i>Implementation Science</i> , 2018 , 13, 25	8.4	9
85	Men benefit more from midlife leisure-time physical activity than women regarding the development of late-life disabilityresults of the KORA-Age study. <i>Preventive Medicine</i> , 2014 , 62, 8-13	4.3	9
84	Interventions for the prevention and treatment of disability due to acquired joint contractures in older people: a systematic review. <i>Age and Ageing</i> , 2017 , 46, 373-382	3	9
83	Male sex and poverty predict abrupt health decline: Deficit accumulation patterns and trajectories in the KORA-Age cohort study. <i>Preventive Medicine</i> , 2017 , 102, 31-38	4.3	9
82	Atmospheric Pressure and Onset of Episodes of Menifle@ Disease - A Repeated Measures Study. <i>PLoS ONE</i> , 2016 , 11, e0152714	3.7	9
81	Visual height intolerance and acrophobia: distressing partners for life. <i>Journal of Neurology</i> , 2016 , 263, 1946-53	5.5	9
80	A comparative study of patient-reported functional outcomes in acute rheumatoid arthritis. <i>Journal of Rheumatology</i> , 2007 , 34, 64-9	4.1	9
79	Vertigo and dizziness cause considerable more health care resource use and costs: results from the KORA FF4 study. <i>Journal of Neurology</i> , 2019 , 266, 2120-2128	5.5	8
78	Development of an International Classification of Functioning, Disability and Health (ICF)-based standard set to describe the impact of joint contractures on participation of older individuals in geriatric care settings. <i>Archives of Gerontology and Geriatrics</i> , 2015 , 61, 61-6	4	8
77	Outcome measures in older persons with acquired joint contractures: a systematic review and content analysis using the ICF (International Classification of Functioning, Disability and Health) as a reference. <i>BMC Geriatrics</i> , 2016 , 16, 40	4.1	8

76	Usability of the Video Head Impulse Test: Lessons From the Population-Based Prospective KORA Study. <i>Frontiers in Neurology</i> , 2018 , 9, 659	4.1	8
75	Self-efficacy beliefs are associated with visual height intolerance: a cross-sectional survey. <i>PLoS ONE</i> , 2014 , 9, e116220	3.7	8
74	Brief ICF Core Sets for the acute hospital. <i>Journal of Rehabilitation Medicine</i> , 2011 , 43, 123-30	3.4	8
73	Early-stage impairments and limitations of functioning from the geriatric ICF core set as determinants of independent living in older patients after discharge from post-acute rehabilitation. <i>Acta Dermato-Venereologica</i> , 2007 , 39, 591-7	2.2	8
72	Quality of life in older individuals with joint contractures in geriatric care settings. <i>Quality of Life Research</i> , 2016 , 25, 2269-81	3.7	8
71	Who gets lost and why: A representative cross-sectional survey on sociodemographic and vestibular determinants of wayfinding strategies. <i>PLoS ONE</i> , 2019 , 14, e0204781	3.7	7
70	Being born in the aftermath of World War II increases the risk for health deficit accumulation in older age: results from the KORA-Age study. <i>European Journal of Epidemiology</i> , 2019 , 34, 675-687	12.1	7
69	Mediator Effect of Balance Problems on Association Between Grip Strength and Falls in Older Adults: Results From the KORA-Age Study. <i>Gerontology and Geriatric Medicine</i> , 2018 , 4, 2333721418760	123	7
68	Graphical modeling of binary data using the LASSO: a simulation study. <i>BMC Medical Research Methodology</i> , 2012 , 12, 16	4.7	7
67	Fear of heights and mild visual height intolerance independent of alcohol consumption. <i>Brain and Behavior</i> , 2013 , 3, 596-601	3.4	7
66	Graphical modeling can be used to illustrate associations between variables describing functioning in head and neck cancer patients. <i>Journal of Clinical Epidemiology</i> , 2011 , 64, 885-92	5.7	7
65	Effects of Whole Body Vibration Therapy and Classic Physiotherapy on Postural Stability in People With Back Pain: A Randomized Trial. <i>Clinical Spine Surgery</i> , 2019 , 32, E214-E220	1.8	7
64	Development of a complex intervention to improve participation of nursing home residents with joint contractures: a mixed-method study. <i>BMC Geriatrics</i> , 2018 , 18, 61	4.1	6
63	Perturbed Development of Striatal Dopamine Transporters in Fatty Versus Lean Zucker Rats: a Follow-up Small Animal PET Study. <i>Molecular Imaging and Biology</i> , 2015 , 17, 521-8	3.8	6
62	Psychosocial factors associated with the effects of physiotherapy in the acute hospital. <i>Disability and Rehabilitation</i> , 2011 , 33, 2311-21	2.4	6
61	ICF-Core-Sets im Akutkrankenhaus und in der Frßrehabilitation fßPatienten mit Erkrankungen des muskuloskelettalen Systems. <i>Physikalische Medizin Rehabilitationsmedizin Kurortmedizin</i> , 2009 , 19, 14-21	0.5	6
60	Operationalisierung von ICF-Kategorien fil physiotherapeutische Interventionen im Akutkrankenhaus. <i>Physioscience</i> , 2008 , 4, 7-16	0.3	6
59	Computerized clinical decision system and mobile application with expert support to optimize management of vertigo in primary care: study protocol for a pragmatic cluster-randomized controlled trial. <i>Journal of Neurology</i> , 2020 , 267, 45-50	5.5	6

58	Magnitude and predictors of khat use among patients with tuberculosis in Southwest Ethiopia: A longitudinal study. <i>PLoS ONE</i> , 2020 , 15, e0236154	3.7	6
57	The development of the ICF vestibular environmental scale. <i>Journal of Vestibular Research:</i> Equilibrium and Orientation, 2016 , 26, 297-302	2.5	6
56	The PaArticular Scales - A new outcome measure to quantify the impact of joint contractures on activities and participation in individuals in geriatric care: Development and Rasch analysis. <i>International Journal of Nursing Studies</i> , 2016 , 59, 107-17	5.8	6
55	Sociodemographic characteristics determine download and use of a Corona contact tracing app in Germany-Results of the COSMO surveys. <i>PLoS ONE</i> , 2021 , 16, e0256660	3.7	6
54	Multicenter data banking in management of dizzy patients: first results from the DizzyNet registry project. <i>Journal of Neurology</i> , 2018 , 265, 3-8	5.5	5
53	Exposure to anticholinergic and sedative medications using the Drug Burden Index and its association with vertigo, dizziness and balance problems in older people - Results from the KORA-FF4 Study. <i>Experimental Gerontology</i> , 2019 , 124, 110644	4.5	5
52	Multi-morbidity and disability, findings from the KORA-Age study. BMC Proceedings, 2013, 7, S10	2.3	5
51	Incremental cost-effectiveness analysis of a multidisciplinary renal education program for patients with chronic renal disease. <i>Disability and Rehabilitation</i> , 2010 , 32, 392-401	2.4	5
50	Identification of Relevant ICF Categories by Patients in Early Post-Acute Rehabilitation Facilities. <i>Physikalische Medizin Rehabilitationsmedizin Kurortmedizin</i> , 2005 , 15, 168-173	0.5	5
49	Developing and piloting a multifactorial intervention to address participation and quality of life in nursing home residents with joint contractures (JointConImprove): study protocol. <i>GMS German Medical Science</i> , 2015 , 13, Doc13	3.2	5
48	Implementation of a complex intervention to improve participation in older people with joint contractures living in nursing homes: a process evaluation of a cluster-randomised pilot trial. <i>BMC Geriatrics</i> , 2020 , 20, 270	4.1	5
47	Minimum data set to measure rehabilitation needs and health outcome after major trauma: application of an international framework. <i>European Journal of Physical and Rehabilitation Medicine</i> , 2016 , 52, 331-46	4.4	5
46	Validation of the comprehensive ICF Core Sets for patients in early post-acute rehabilitation facilities. <i>Journal of Rehabilitation Medicine</i> , 2011 , 43, 102-12	3.4	4
45	Effectiveness of a community-based 3-year advisory program after acquired brain injury. <i>European Journal of Neurology</i> , 2007 , 14, 1256-65	6	4
44	Lack of association of MRI determined subclinical cardiovascular disease with dizziness and vertigo in a cross-sectional population-based study. <i>PLoS ONE</i> , 2017 , 12, e0184858	3.7	4
43	Development and validation of a classification algorithm to diagnose and differentiate spontaneous episodic vertigo syndromes: results from the DizzyReg patient registry. <i>Journal of Neurology</i> , 2020 , 267, 160-167	5.5	4
42	RECAPDOC - a questionnaire for the documentation of rehabilitation care utilization in individuals with disorders of consciousness in long-term care in Germany: development and pretesting. <i>BMC Health Services Research</i> , 2018 , 18, 329	2.9	3
41	Improved participation of older people with joint contractures living in nursing homes: feasibility of study procedures in a cluster-randomised pilot trial. <i>Trials</i> , 2019 , 20, 411	2.8	3

40	The ICF and Florence Nightingale - bringing data to statistical proof. <i>Journal of Rehabilitation Medicine</i> , 2011 , 43, 1041-2	3.4	3
39	Separation from children as a specific risk factor to fathersQhealth and lifestyles. <i>International Journal of Public Health</i> , 2001 , 46, 272-8		3
38	Standardization of non-aggregated data: theory and practice. <i>Computer Methods and Programs in Biomedicine</i> , 2001 , 65, 207-27	6.9	3
37	Anticholinergic and sedative medications and the risk of vertigo or dizziness in the German primary care setting-A matched case-control study from the CONTENT registry. <i>Pharmacoepidemiology and Drug Safety</i> , 2018 , 27, 912-920	2.6	2
36	Chronic vestibular syndromes in the elderly: Presbyvestibulopathy - an isolated clinical entity?. <i>European Journal of Neurology</i> , 2022 ,	6	2
35	The times we are born into and our lifestyle choices determine our health trajectories in older age - Results from the KORA-Age study. <i>Preventive Medicine</i> , 2020 , 133, 106025	4.3	1
34	Patient trajectories and their impact on mobility, social participation and quality of life in patients with vertigo/dizziness/balance disorders and osteoarthritis (MobilE-TRA): study protocol of an observational, practice-based cohort study. <i>BMJ Open</i> , 2018 , 8, e022970	3	1
33	Pilot cluster randomized controlled trial of a complex intervention to improve management of vertigo in primary care (PRIMA-Vertigo): study protocol. <i>Current Medical Research and Opinion</i> , 2018 , 34, 1819-1828	2.5	1
32	Standardisierung der Ziele physiotherapeutischer Interventionen im Akutkrankenhaus mithilfe der ICF. <i>Physioscience</i> , 2008 , 4, 17-24	0.3	1
31	A Set of Eight Key Questions Helps to Classify Common Vestibular Disorders-Results From the DizzyReg Patient Registry. <i>Frontiers in Neurology</i> , 2021 , 12, 670944	4.1	1
30	Wealth and occupation determine health deficit accumulation onset in Europe - Results from the SHARE study. <i>Experimental Gerontology</i> , 2018 , 113, 74-79	4.5	1
29	Using Base-ml to Learn Classification of Common Vestibular Disorders on DizzyReg Registry Data. <i>Frontiers in Neurology</i> , 2021 , 12, 681140	4.1	1
28	Developing and Implementing Diagnostic Prediction Models for Vestibular Diseases in Primary Care. Studies in Health Technology and Informatics, 2016 , 228, 735-9	0.5	1
27	Vestibular paroxysmia: clinical characteristics and long-term course. Journal of Neurology,	5.5	1
26	Physiotherapeutischer Ressourcenaufwand im Akutkrankenhaus. <i>Physioscience</i> , 2008 , 4, 25-29	0.3	О
25	Netzwerke zur Versorgung Parkinsonerkrankter. <i>Nervenheilkunde</i> , 2021 , 40, 768-777	0.3	O
24	"WeQe in good hands there." - Acceptance, barriers and facilitators of a primary care-based health coaching programme for children and adolescents with mental health problems: a qualitative study (PrimA-QuO). BMC Family Practice, 2020, 21, 273	2.6	0
23	Development of a complex intervention to improve mobility and participation of older people with vertigo, dizziness and balance disorders in primary care: a mixed methods study. <i>BMC Family Practice</i> , 2021 , 22, 89	2.6	O

(2020-2019)

22	Administrators, health service providers, and consumers perspectives of functions of district health-care systems in Oromia region, Ethiopia: A qualitative study. <i>International Journal of Health Planning and Management</i> , 2019 , 34, 836-850	2.2	Ο
21	Economic evaluation of guideline implementation in primary care: a systematic review. <i>International Journal for Quality in Health Care</i> , 2020 , 32, 1-11	1.9	О
20	Living longer but less healthy: The female disadvantage in health expectancy. Results from the KORA-Age study. <i>Experimental Gerontology</i> , 2021 , 145, 111196	4.5	O
19	Substance use disorders and adherence to antituberculosis medications in Southwest Ethiopia: a prospective cohort study. <i>BMJ Open</i> , 2021 , 11, e043050	3	O
18	Mixed-methods evaluation of a structured primary care programme for children and adolescents with mental health problems (PrimA-QuO): a study protocol. <i>BMJ Open</i> , 2021 , 11, e052747	3	O
17	Gender-linked determinants for utilization of long-term care in community-dwelling adults 65+ in Germany: Results from the population-based KORA-Age study. <i>Experimental Gerontology</i> , 2021 , 153, 111500	4.5	O
16	Referral trajectories in patients with vertigo, dizziness and balance disorders and their impact on health-related quality of life and functioning: results from the longitudinal multicenter study MobilE-TRA <i>Journal of Neurology</i> , 2022 , 1	5.5	О
15	Associations of Disease Activity and Socio-demographic Factors with Activity and Participation of Patients with Rheumatoid Arthritis Results of a Cross-sectional Study. <i>Physikalische Medizin Rehabilitationsmedizin Kurortmedizin</i> , 2015 , 25, 141-147	0.5	
14	Epidemiologie - Ausbildung in Deutschland. <i>Public Health Forum</i> , 2012 , 20, 25-26	0.1	
13	P2-479 Determinants of functioning and disability in aged persons: results from the KORA-Age Study. <i>Journal of Epidemiology and Community Health</i> , 2011 , 65, A353-A353	5.1	
12	Epidemiology of Vestibular Impairments 2020 , 555-566		
11	Face and content validity of a prospective multidimensional performance instrument for service delivery in district health systems in low-income countries: a Delphi study. <i>International Health</i> , 2020 , 12, 184-191	2.4	
10	Identification of relevant performance indicators for district healthcare systems in Ethiopia: a systematic review and expert opinion. <i>International Journal for Quality in Health Care</i> , 2020 , 32, 161-17	72 ^{1.9}	
9	Effect of food insecurity on mental health of patients with tuberculosis in Southwest Ethiopia: a prospective cohort study. <i>BMJ Open</i> , 2021 , 11, e045434	3	
8	Beschreibung und Einordnung der Covid-19-Pandemie 2022 , 5-18		
7	Behavioural and patient-individual determinants of quality of life, functioning and physical activity in older adults (MobilE-TRA 2): study protocol of an observational cohort study in a tertiary care setting. <i>BMJ Open</i> , 2021 , 11, e051915	3	
6	Magnitude and predictors of khat use among patients with tuberculosis in Southwest Ethiopia: A longitudinal study 2020 , 15, e0236154		
5	Magnitude and predictors of khat use among patients with tuberculosis in Southwest Ethiopia: A longitudinal study 2020 , 15, e0236154		

- Magnitude and predictors of khat use among patients with tuberculosis in Southwest Ethiopia: A longitudinal study **2020**, 15, e0236154
- Magnitude and predictors of khat use among patients with tuberculosis in Southwest Ethiopia: A longitudinal study **2020**, 15, e0236154
- Magnitude and predictors of khat use among patients with tuberculosis in Southwest Ethiopia: A longitudinal study **2020**, 15, e0236154
- Magnitude and predictors of khat use among patients with tuberculosis in Southwest Ethiopia: A longitudinal study **2020**, 15, e0236154