

Tracey DiSipio

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/2531363/tracey-disipio-publications-by-year.pdf>

Version: 2024-04-20

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

23
papers

1,735
citations

17
h-index

23
g-index

23
ext. papers

2,074
ext. citations

4
avg, IF

4.69
L-index

#	Paper	IF	Citations
23	Risk of Second Primary Cancer in Survivors of In Situ Melanoma. <i>Journal of Investigative Dermatology</i> , 2019 , 139, 842-847	4.3	4
22	Cost-effectiveness of a pragmatic exercise intervention for women with breast cancer: results from a randomized controlled trial. <i>Psycho-Oncology</i> , 2017 , 26, 649-655	3.9	19
21	Systematic Review and Meta-Analysis of the Effects of Exercise for Those With Cancer-Related Lymphedema. <i>Archives of Physical Medicine and Rehabilitation</i> , 2016 , 97, 302-315.e13	2.8	57
20	A systematic review and meta-analysis of prescribing practices of antidepressants in cancer patients. <i>Psycho-Oncology</i> , 2016 , 25, 1002-16	3.9	37
19	Weight and weight change following breast cancer: evidence from a prospective, population-based, breast cancer cohort study. <i>BMC Cancer</i> , 2015 , 15, 28	4.8	44
18	Adverse breast cancer treatment effects: the economic case for making rehabilitative programs standard of care. <i>Supportive Care in Cancer</i> , 2015 , 23, 1807-17	3.9	29
17	The Lymphedema Evaluation in Gynecological cancer Study (LEGS): design of a prospective, longitudinal, cohort study. <i>Cancer Research Frontiers</i> , 2015 , 1, 104-118		7
16	Incidence of unilateral arm lymphoedema after breast cancer: a systematic review and meta-analysis. <i>Lancet Oncology</i> , 2013 , 14, 500-15	21.7	907
15	Exercise for health: a randomized, controlled trial evaluating the impact of a pragmatic, translational exercise intervention on the quality of life, function and treatment-related side effects following breast cancer. <i>Breast Cancer Research and Treatment</i> , 2013 , 137, 175-86	4.4	109
14	Bodyweight and other correlates of symptom-detected breast cancers in a population offered screening. <i>Cancer Causes and Control</i> , 2012 , 23, 89-102	2.8	6
13	Prevalence of breast cancer treatment sequelae over 6 years of follow-up: the Pulling Through Study. <i>Cancer</i> , 2012 , 118, 2217-25	6.4	96
12	Patterns, correlates, and prognostic significance of quality of life following breast cancer. <i>Psycho-Oncology</i> , 2011 , 20, 1084-91	3.9	17
11	Prevalence and prognostic significance of secondary lymphedema following breast cancer. <i>Lymphatic Research and Biology</i> , 2011 , 9, 135-41	2.3	56
10	Does quality of life among breast cancer survivors one year after diagnosis differ depending on urban and non-urban residence? A comparative study. <i>Health and Quality of Life Outcomes</i> , 2010 , 8, 3	3	30
9	Upper-body morbidity following breast cancer treatment is common, may persist longer-term and adversely influences quality of life. <i>Health and Quality of Life Outcomes</i> , 2010 , 8, 92	3	83
8	Prevalence and determinants of sunburn in Queensland. <i>Health Promotion Journal of Australia</i> , 2009 , 20, 102-6	1.7	5
7	The Queensland Cancer Risk Study: general population norms for the Functional Assessment of Cancer Therapy-General (FACT-G). <i>Psycho-Oncology</i> , 2009 , 18, 606-14	3.9	45

6	What determines the health-related quality of life among regional and rural breast cancer survivors?. <i>Australian and New Zealand Journal of Public Health</i> , 2009 , 33, 534-9	2.3	22
5	Health-related quality of life 18 months after breast cancer: comparison with the general population of Queensland, Australia. <i>Supportive Care in Cancer</i> , 2008 , 16, 1141-50	3.9	36
4	Solaria use in Queensland, Australia. <i>Australian and New Zealand Journal of Public Health</i> , 2006 , 30, 479-82	2.3	23
3	The Queensland Cancer Risk Study: behavioural risk factor results. <i>Australian and New Zealand Journal of Public Health</i> , 2006 , 30, 375-82	2.3	34
2	How do recovery advice and behavioural characteristics influence upper-body function and quality of life among women 6 months after breast cancer diagnosis?. <i>Supportive Care in Cancer</i> , 2006 , 14, 22-9	3.9	10
1	Perceptions of upper-body problems during recovery from breast cancer treatment. <i>Supportive Care in Cancer</i> , 2004 , 12, 106-13	3.9	59