

Jing Zhang

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/2535489/jing-zhang-publications-by-year.pdf>

Version: 2024-04-23

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

9
papers

12,926
citations

3
h-index

10
g-index

10
ext. papers

16,001
ext. citations

7.7
avg, IF

7.12
L-index

#	Paper	IF	Citations
9	Multi-Parametric Diffusion Tensor Imaging of The Optic Nerve for Detection of Dysthyroid Optic Neuropathy in Patients With Thyroid-Associated Ophthalmopathy.. <i>Frontiers in Endocrinology</i> , 2022 , 13, 851143	5.7	0
8	The diagnostic value of the IDEAL-T2WI sequence in dysthyroid optic neuropathy: a quantitative analysis of the optic nerve and cerebrospinal fluid in the optic nerve sheath. <i>European Radiology</i> , 2021 , 31, 7419-7428	8	1
7	Effect of early oxygen therapy and antiviral treatment on disease progression in patients with COVID-19: A retrospective study of medical charts in China. <i>PLoS Neglected Tropical Diseases</i> , 2021 , 15, e0009051	4.8	5
6	Baseline Volumetric T2 Relaxation Time Histogram Analysis: Can It Be Used to Predict the Response to Intravenous Methylprednisolone Therapy in Patients With Thyroid-Associated Ophthalmopathy?. <i>Frontiers in Endocrinology</i> , 2021 , 12, 614536	5.7	0
5	Prediction of treatment response to intravenous glucocorticoid in patients with thyroid-associated ophthalmopathy using T2 mapping and T2 IDEAL. <i>European Journal of Radiology</i> , 2021 , 142, 109839	4.7	1
4	Clinical Characteristics of 138 Hospitalized Patients With 2019 Novel Coronavirus-Infected Pneumonia in Wuhan, China. <i>JAMA - Journal of the American Medical Association</i> , 2020 , 323, 1061-1069	27.4	12910
3	Histogram analysis of T2 mapping for detecting early involvement of extraocular muscles in patients with thyroid-associated ophthalmopathy. <i>Scientific Reports</i> , 2020 , 10, 19445	4.9	2
2	Combined application of isotropic three-dimensional fast spin echo (3D-FSE-Cube) with 2-point Dixon fat/water separation (FLEX) and 3D-FSE-cube in MR dacryocystography. <i>British Journal of Radiology</i> , 2019 , 92, 20180157	3.4	3
1	Diagnostic performance of the three-dimensional fast spin echo-Cube sequence in comparison with a conventional imaging protocol in evaluation of the lachrymal drainage system. <i>European Radiology</i> , 2015 , 25, 635-43	8	4