

Rosa Elena Navarro-Hernández

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/254133/publications.pdf>

Version: 2024-02-01

37
papers

656
citations

516561

16
h-index

610775

24
g-index

40
all docs

40
docs citations

40
times ranked

1192
citing authors

#	ARTICLE	IF	CITATIONS
1	Macrophage migration inhibitory factor (MIF): Genetic evidence for participation in early onset and early stage rheumatoid arthritis. <i>Cytokine</i> , 2013, 61, 759-765.	1.4	59
2	Expression of ICAM1 and VCAM1 Serum Levels in Rheumatoid Arthritis Clinical Activity. Association with Genetic Polymorphisms. <i>Disease Markers</i> , 2009, 26, 119-126.	0.6	42
3	Tumor Necrosis Factor $\hat{\pm}$ -308 and -238 Polymorphisms in Rheumatoid Arthritis. Association With Messenger RNA Expression and sTNF- $\hat{\pm}$. <i>Journal of Investigative Medicine</i> , 2008, 56, 937-943.	0.7	34
4	The +1858C/T PTPN22 gene polymorphism confers genetic susceptibility to rheumatoid arthritis in Mexican population from the Western Mexico. <i>Immunology Letters</i> , 2012, 147, 41-46.	1.1	34
5	Serum levels of macrophage migration inhibitory factor are associated with rheumatoid arthritis course. <i>Rheumatology International</i> , 2012, 32, 2307-2311.	1.5	33
6	Serum Levels of Anticyclic Citrullinated Peptide Antibodies, Interleukin-6, Tumor Necrosis Factor- $\hat{\pm}$, and C-Reactive Protein Are Associated with Increased Carotid Intima-Media Thickness: A Cross-Sectional Analysis of a Cohort of Rheumatoid Arthritis Patients without Cardiovascular Risk Factors. <i>BioMed Research International</i> , 2015, 2015, 1-10.	0.9	30
7	Detection of autoantibodies to DSF70/LEDGFp75 in Mexican Hispanics using multiple complementary assay platforms. <i>Autoimmunity Highlights</i> , 2017, 8, 1.	3.9	28
8	Disease duration of rheumatoid arthritis is a predictor of vascular stiffness. <i>Medicine (United States)</i> , 2017, 96, e7862.	0.4	25
9	The Impact of <i>LEP</i> G-2548A and <i>LEPR</i> Gln223Arg Polymorphisms on Adiposity, Leptin, and Leptin-Receptor Serum Levels in a Mexican Mestizo Population. <i>BioMed Research International</i> , 2015, 2015, 1-10.	0.9	24
10	Subclinical parameters of arterial stiffness and arteriosclerosis correlate with QRISK3 in systemic lupus erythematosus. <i>PLoS ONE</i> , 2018, 13, e0207520.	1.1	24
11	Massive myeloid sarcoma affecting the central nervous system, mediastinum, retroperitoneum, liver, and rectum associated with acute myeloblastic leukaemia: a case report. <i>Journal of Clinical Pathology</i> , 2005, 58, 325-327.	1.0	22
12	Reduced IgG anti-small nuclear ribonucleoprotein autoantibody production in systemic lupus erythematosus patients with positive IgM anti-cytomegalovirus antibodies. <i>Arthritis Research and Therapy</i> , 2009, 11, R27.	1.6	22
13	Tumor necrosis factor receptor 2 M196R polymorphism in rheumatoid arthritis and osteoarthritis: relationship with sTNFR2 levels and clinical features. <i>Rheumatology International</i> , 2006, 27, 53-59.	1.5	21
14	The Montreal Cognitive Assessment Test. <i>Journal of Clinical Rheumatology</i> , 2019, 25, 325-328.	0.5	21
15	The 482Ser of <i>PPARGC1A</i> and 12Pro of <i>PPARG2</i> Alleles Are Associated with Reduction of Metabolic Risk Factors Even Obesity in a Mexican-Mestizo Population. <i>BioMed Research International</i> , 2015, 2015, 1-14.	0.9	19
16	High prevalence of autoantibodies to RNA helicase A in Mexican patients with systemic lupus erythematosus. <i>Arthritis Research and Therapy</i> , 2010, 12, R6.	1.6	18
17	Circulating TNFRI and TNFRII levels correlated with the disease activity score (DAS28) in rheumatoid arthritis. <i>Scandinavian Journal of Rheumatology</i> , 2009, 38, 332-335.	0.6	17
18	Association of ADIPOQ +45T>G polymorphism with body fat mass and blood levels of soluble adiponectin and inflammation markers in a Mexican-Mestizo population. <i>Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy</i> , 2012, 5, 369.	1.1	17

#	ARTICLE	IF	CITATIONS
19	Serum leptin and serum leptin/serum leptin receptor ratio imbalance in obese rheumatoid arthritis patients positive for anti-cyclic citrullinated peptide antibodies. <i>Arthritis Research and Therapy</i> , 2015, 17, 335.	1.6	16
20	Characterization of B7H6, an endogenous ligand for the NK cell activating receptor NKp30, reveals the identity of two different soluble isoforms during normal human pregnancy. <i>Immunobiology</i> , 2018, 223, 57-63.	0.8	16
21	PAI-1 mRNA expression and plasma level in rheumatoid arthritis: relationship with 4G/5G PAI-1 polymorphism. <i>Rheumatology International</i> , 2012, 32, 3951-3956.	1.5	13
22	Ageing influences the relationship of circulating miR-33a and miR-33b levels with insulin resistance and adiposity. <i>Diabetes and Vascular Disease Research</i> , 2019, 16, 244-253.	0.9	13
23	Inverse Relationship of the CMKLR1 Relative Expression and Chemerin Serum Levels in Obesity with Dysmetabolic Phenotype and Insulin Resistance. <i>Mediators of Inflammation</i> , 2016, 2016, 1-9.	1.4	12
24	Downregulation of hippocampal NR2A/2B subunits related to cognitive impairment in a pristane-induced lupus BALB/c mice. <i>PLoS ONE</i> , 2019, 14, e0217190.	1.1	12
25	CCL2 Serum Levels and Adiposity Are Associated with the Polymorphic Phenotypes -2518A on CCL2 and 641LE on CCR2 in a Mexican Population with Insulin Resistance. <i>Journal of Diabetes Research</i> , 2016, 2016, 1-11.	1.0	11
26	Disease Activity Score on 28 Joints and Polypharmacy Are Independent Predictors for Health-Related Quality of Life Evaluated by INCAVISA in Patients With Rheumatoid Arthritis. <i>Journal of Clinical Rheumatology</i> , 2016, 22, 399-404.	0.5	11
27	Low Levels of CD36 in Peripheral Blood Monocytes in Subclinical Atherosclerosis in Rheumatoid Arthritis: A Cross-Sectional Study in a Mexican Population. <i>BioMed Research International</i> , 2014, 2014, 1-6.	0.9	10
28	FAS -670A>G promoter polymorphism is associated with soluble Fas levels in primary Sjögren's syndrome. <i>Genetics and Molecular Research</i> , 2014, 13, 4831-4838.	0.3	9
29	Circulating E-selectin and tumor necrosis factor- α in extraarticular involvement and joint disease activity in rheumatoid arthritis. <i>Rheumatology International</i> , 2009, 29, 281-286.	1.5	8
30	Effect of pregnancy hormone mixtures on cytokine production and surface marker expression in naïve and LPS-activated THP-1 differentiated monocytes/macrophages. <i>Innate Immunity</i> , 2020, 26, 84-96.	1.1	8
31	sE-Selectin expression and A561C polymorphism in relation to rheumatoid arthritis clinical activity. <i>Journal of Rheumatology</i> , 2006, 33, 1968-72.	1.0	5
32	TNFR1-383 A>C polymorphism association with clinical manifestations in primary Sjögren's syndrome patients. <i>Genetics and Molecular Research</i> , 2016, 15, .	0.3	3
33	Extrapulmonary Tuberculosis Infection in Mexican Patients With Idiopathic Inflammatory Myopathies. <i>Journal of Clinical Rheumatology</i> , 2015, 21, 331-332.	0.5	2
34	Low CD36 and LOX-1 Levels and CD36 Gene Subexpression Are Associated with Metabolic Dysregulation in Older Individuals with Abdominal Obesity. <i>Journal of Diabetes Research</i> , 2016, 2016, 1-10.	1.0	2
35	Sensorineural Hearing Impairment and Subclinical Atherosclerosis in Rheumatoid Arthritis Patients Without Traditional Cardiovascular Risk Factors. <i>Archives of Rheumatology</i> , 2016, 31, 208-214.	0.3	1
36	CCR2/CCL2 and CMKLR1/RvE1 chemokines system levels are associated with insulin resistance in rheumatoid arthritis. <i>PLoS ONE</i> , 2021, 16, e0246054.	1.1	1

#	ARTICLE	IF	CITATIONS
37	Evaluación clínica y ultrasonográfica de la glándula tiroides en pacientes con artritis reumatoide. Gaceta Medica De Mexico, 2018, 154, 432-437.	0.5	1