## Muhammad Zaid Iskandar

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/2546621/publications.pdf

Version: 2024-02-01

1684188 1588992 12 70 5 8 citations g-index h-index papers 13 13 13 104 docs citations times ranked citing authors all docs

#	Article	IF	Citations
1	Plasma Desmosine and Abdominal Aortic Aneurysm Disease. Journal of the American Heart Association, 2019, 8, e013743.	3.7	22
2	5-Fluorouracil cardiotoxicity: reversible left ventricular systolic dysfunction with early detection. BMJ Case Reports, 2015, 2015, bcr2015209347-bcr2015209347.	0.5	13
3	Sacubitril and valsartan fixed combination to reduce heart failure events in post-acute myocardial infarction patients. Drugs of Today, 2017, 53, 545.	1.1	12
4	Biomarkers of Aortopathy in Marfan Syndrome. Cardiology in Review, 2020, 28, 92-97.	1.4	9
5	The future of pharmacogenetics in the treatment of heart failure. Pharmacogenomics, 2015, 16, 1817-1827.	1.3	7
6	Inferior Vena Cava Dilatation Predicts Mortality and Worsening Renal Function in Patients With Chronic Heart Failure. Journal of Cardiac Failure, 2015, 21, S70.	1.7	2
7	Acute spinal cord compression: a rare complication of dual antiplatelet therapy: FigureÂ1. BMJ Case Reports, 2015, 2015, bcr2015209952.	0.5	2
8	P6040The novel biomarker plasma desmosine, a marker of elastin breakdown, is an independent predictor of abdominal aortic aneurysm events independent of aneurysm size. European Heart Journal, 2018, 39, .	2.2	1
9	5â€Exaggerated elastin turnover in childhood and adolescence in marfan syndrome - correlation with age - new insights from the aims trial. , 2021, , .		1
10	28â€Heart Rate is not Related to Survival in Patients with Heart Failure and Atrial Fibrillation. Heart, 2015, 101, A15.2-A16.	2.9	0
11	126â€Urinary desmosine, a biomarker of elastin degradation is significantly elevated and associated with maximum aortic root size and aortic Z-scores in patients with bicuspid aortic valve. , 2019, , .		0
12	65â€The novel plasma biomarker desmosine, a marker of elastin breakdown, is an independent predictor of abdominal aortic aneurysm events independent of aneurysm size. , 2018, , .		0