

Ashham Mansur

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2548817/publications.pdf>

Version: 2024-02-01

24
papers

385
citations

759055

12
h-index

794469

19
g-index

24
all docs

24
docs citations

24
times ranked

731
citing authors

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 1 | Impact of statin therapy on mortality in patients with sepsis-associated acute respiratory distress syndrome (ARDS) depends on ARDS severity: a prospective observational cohort study. <i>BMC Medicine</i> , 2015, 13, 128. | 2.3 | 57 |
| 2 | Extracorporeal life support with left ventricular decompressionâ€”improved survival in severe cardiogenic shock: results from a retrospective study. <i>PeerJ</i> , 2017, 5, e3813. | 0.9 | 29 |
| 3 | Chronic kidney disease is associated with a higher 90-day mortality than other chronic medical conditions in patients with sepsis. <i>Scientific Reports</i> , 2015, 5, 10539. | 1.6 | 27 |
| 4 | Results of concomitant groin-free percutaneous temporary RVAD support using a centrifugal pump with a double-lumen jugular venous cannula in LVAD patients. <i>Journal of Thoracic Disease</i> , 2019, 11, S913-S920. | 0.6 | 27 |
| 5 | The regulatory toll-like receptor 4 genetic polymorphism rs11536889 is associated with renal, coagulation and hepatic organ failure in sepsis patients. <i>Journal of Translational Medicine</i> , 2014, 12, 177. | 1.8 | 25 |
| 6 | Ninety-Day Survival Rate of Patients with Sepsis Relates to Programmed Cell Death 1 Genetic Polymorphism rs11568821. <i>Journal of Investigative Medicine</i> , 2014, 62, 638-643. | 0.7 | 21 |
| 7 | Primary bacteraemia is associated with a higher mortality risk compared with pulmonary and intra-abdominal infections in patients with sepsis: a prospective observational cohort study. <i>BMJ Open</i> , 2015, 5, e006616-e006616. | 0.8 | 21 |
| 8 | Prehospital ultrasound-guided nerve blocks improve reduction-feasibility of dislocated extremity injuries compared to systemic analgesia. A randomized controlled trial. <i>PLoS ONE</i> , 2018, 13, e0199776. | 1.1 | 21 |
| 9 | The FER rs4957796 TT genotype is associated with unfavorable 90-day survival in Caucasian patients with severe ARDS due to pneumonia. <i>Scientific Reports</i> , 2017, 7, 9887. | 1.6 | 18 |
| 10 | The CD14 rs2569190 TT Genotype Is Associated with an Improved 30-Day Survival in Patients with Sepsis: A Prospective Observational Cohort Study. <i>PLoS ONE</i> , 2015, 10, e0127761. | 1.1 | 16 |
| 11 | The CTLA-4 rs231775 GG genotype is associated with favorable 90-day survival in Caucasian patients with sepsis. <i>Scientific Reports</i> , 2018, 8, 15140. | 1.6 | 13 |
| 12 | Pre-hospital transthoracic echocardiography for early identification of non-ST-elevation myocardial infarction in patients with acute coronary syndrome. <i>Critical Care</i> , 2018, 22, 29. | 2.5 | 13 |
| 13 | CTLA-4 Genetic Variants Predict Survival in Patients with Sepsis. <i>Journal of Clinical Medicine</i> , 2019, 8, 70. | 1.0 | 13 |
| 14 | Favorable 90-Day Mortality in Obese Caucasian Patients with Septic Shock According to the Sepsis-3 Definition. <i>Journal of Clinical Medicine</i> , 2020, 9, 46. | 1.0 | 13 |
| 15 | Gene Expressionâ€”Based Diagnosis of Infections in Critically Ill Patientsâ€”Prospective Validation of the SepsisMetaScore in a Longitudinal Severe Trauma Cohort. <i>Critical Care Medicine</i> , 2021, 49, e751-e760. | 0.4 | 10 |
| 16 | Perioperative Blood Glucose Levelsâ€” <150 â€”mg/dL are Associated With Improved 5-Year Survival in Patients Undergoing On-Pump Cardiac Surgery. <i>Medicine (United States)</i> , 2015, 94, e2035. | 0.4 | 9 |
| 17 | Effect of the Lymphocyte Activation Gene 3 Polymorphism rs951818 on Mortality and Disease Progression in Patients with Sepsisâ€”A Prospective Genetic Association Study. <i>Journal of Clinical Medicine</i> , 2021, 10, 5302. | 1.0 | 9 |
| 18 | Combination of general anesthesia and peripheral nerve block with low-dose ropivacaine reduces postoperative pain for several days after outpatient arthroscopy. <i>Medicine (United States)</i> , 2017, 96, e6046. | 0.4 | 8 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 19 | Lack of an Association between the Functional Polymorphism TREM-1 rs2234237 and the Clinical Course of Sepsis among Critically Ill Caucasian Patients – A Monocentric Prospective Genetic Association Study. <i>Journal of Clinical Medicine</i> , 2019, 8, 301. | 1.0 | 8 |
| 20 | TIM-3 Genetic Variants Are Associated with Altered Clinical Outcome and Susceptibility to Gram-Positive Infections in Patients with Sepsis. <i>International Journal of Molecular Sciences</i> , 2020, 21, 8318. | 1.8 | 8 |
| 21 | Prognostic relevance of hemodialysis for short-term survival in patients after LVAD implantation. <i>Scientific Reports</i> , 2018, 8, 8546. | 1.6 | 7 |
| 22 | Anaemia requiring red blood cell transfusion is associated with unfavourable 90-day survival in surgical patients with sepsis. <i>BMC Research Notes</i> , 2018, 11, 879. | 0.6 | 6 |
| 23 | Paths of femoral nerve catheters placed using ultrasound-guided in plane vs out of plane techniques. <i>Medicine (United States)</i> , 2018, 97, e12958. | 0.4 | 3 |
| 24 | Genetic Polymorphisms in Endothelin-1 as Predictors for Long-Term Survival and the Cardiac Index in Patients Undergoing On-Pump Cardiac Surgery. <i>PLoS ONE</i> , 2015, 10, e0131155. | 1.1 | 3 |