

Ashham Mansur

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2548817/publications.pdf>

Version: 2024-02-01

24
papers

385
citations

759233
12
h-index

794594
19
g-index

24
all docs

24
docs citations

24
times ranked

731
citing authors

#	ARTICLE	IF	CITATIONS
1	Impact of statin therapy on mortality in patients with sepsis-associated acute respiratory distress syndrome (ARDS) depends on ARDS severity: a prospective observational cohort study. BMC Medicine, 2015, 13, 128.	5.5	57
2	Extracorporeal life support with left ventricular decompressionâ€”improved survival in severe cardiogenic shock: results from a retrospective study. PeerJ, 2017, 5, e3813.	2.0	29
3	Chronic kidney disease is associated with a higher 90-day mortality than other chronic medical conditions in patients with sepsis. Scientific Reports, 2015, 5, 10539.	3.3	27
4	Results of concomitant groin-free percutaneous temporary RVAD support using a centrifugal pump with a double-lumen jugular venous cannula in LVAD patients. Journal of Thoracic Disease, 2019, 11, S913-S920.	1.4	27
5	The regulatory toll-like receptor 4 genetic polymorphism rs11536889 is associated with renal, coagulation and hepatic organ failure in sepsis patients. Journal of Translational Medicine, 2014, 12, 177.	4.4	25
6	Ninety-Day Survival Rate of Patients with Sepsis Relates to Programmed Cell Death 1 Genetic Polymorphism rs11568821. Journal of Investigative Medicine, 2014, 62, 638-643.	1.6	21
7	Primary bacteraemia is associated with a higher mortality risk compared with pulmonary and intra-abdominal infections in patients with sepsis: a prospective observational cohort study. BMJ Open, 2015, 5, e006616-e006616.	1.9	21
8	Prehospital ultrasound-guided nerve blocks improve reduction-feasibility of dislocated extremity injuries compared to systemic analgesia. A randomized controlled trial. PLoS ONE, 2018, 13, e0199776.	2.5	21
9	The FER rs4957796 TT genotype is associated with unfavorable 90-day survival in Caucasian patients with severe ARDS due to pneumonia. Scientific Reports, 2017, 7, 9887.	3.3	18
10	The CD14 rs2569190 TT Genotype Is Associated with an Improved 30-Day Survival in Patients with Sepsis: A Prospective Observational Cohort Study. PLoS ONE, 2015, 10, e0127761.	2.5	16
11	The CTLA-4 rs231775 GG genotype is associated with favorable 90-day survival in Caucasian patients with sepsis. Scientific Reports, 2018, 8, 15140.	3.3	13
12	Pre-hospital transthoracic echocardiography for early identification of non-ST-elevation myocardial infarction in patients with acute coronary syndrome. Critical Care, 2018, 22, 29.	5.8	13
13	CTLA-4 Genetic Variants Predict Survival in Patients with Sepsis. Journal of Clinical Medicine, 2019, 8, 70.	2.4	13
14	Favorable 90-Day Mortality in Obese Caucasian Patients with Septic Shock According to the Sepsis-3 Definition. Journal of Clinical Medicine, 2020, 9, 46.	2.4	13
15	Gene Expressionâ€”Based Diagnosis of Infections in Critically Ill Patientsâ€”Prospective Validation of the SepsisMetaScore in a Longitudinal Severe Trauma Cohort. Critical Care Medicine, 2021, 49, e751-e760.	0.9	10
16	Perioperative Blood Glucose Levelsâ€”<150â€”mg/dL are Associated With Improved 5-Year Survival in Patients Undergoing On-Pump Cardiac Surgery. Medicine (United States), 2015, 94, e2035.	1.0	9
17	Effect of the Lymphocyte Activation Gene 3 Polymorphism rs951818 on Mortality and Disease Progression in Patients with Sepsisâ€”A Prospective Genetic Association Study. Journal of Clinical Medicine, 2021, 10, 5302.	2.4	9
18	Combination of general anesthesia and peripheral nerve block with low-dose ropivacaine reduces postoperative pain for several days after outpatient arthroscopy. Medicine (United States), 2017, 96, e6046.	1.0	8

#	ARTICLE	IF	CITATIONS
19	Lack of an Association between the Functional Polymorphism TREM-1 rs2234237 and the Clinical Course of Sepsis among Critically Ill Caucasian Patientsâ€”A Monocentric Prospective Genetic Association Study. <i>Journal of Clinical Medicine</i> , 2019, 8, 301.	2.4	8
20	TIM-3 Genetic Variants Are Associated with Altered Clinical Outcome and Susceptibility to Gram-Positive Infections in Patients with Sepsis. <i>International Journal of Molecular Sciences</i> , 2020, 21, 8318.	4.1	8
21	Prognostic relevance of hemodialysis for short-term survival in patients after LVAD implantation. <i>Scientific Reports</i> , 2018, 8, 8546.	3.3	7
22	Anaemia requiring red blood cell transfusion is associated with unfavourable 90-day survival in surgical patients with sepsis. <i>BMC Research Notes</i> , 2018, 11, 879.	1.4	6
23	Paths of femoral nerve catheters placed using ultrasound-guided in plane vs out of plane techniques. <i>Medicine (United States)</i> , 2018, 97, e12958.	1.0	3
24	Genetic Polymorphisms in Endothelin-1 as Predictors for Long-Term Survival and the Cardiac Index in Patients Undergoing On-Pump Cardiac Surgery. <i>PLoS ONE</i> , 2015, 10, e0131155.	2.5	3