Ashham Mansur

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/2548817/publications.pdf

Version: 2024-02-01

24 papers

385 citations

759055 12 h-index 19 g-index

24 all docs

24 docs citations

24 times ranked 731 citing authors

#	Article	IF	CITATIONS
1	Gene Expression–Based Diagnosis of Infections in Critically Ill Patients—Prospective Validation of the SepsisMetaScore in a Longitudinal Severe Trauma Cohort. Critical Care Medicine, 2021, 49, e751-e760.	0.4	10
2	Effect of the Lymphocyte Activation Gene 3 Polymorphism rs951818 on Mortality and Disease Progression in Patients with Sepsis—A Prospective Genetic Association Study. Journal of Clinical Medicine, 2021, 10, 5302.	1.0	9
3	Favorable 90-Day Mortality in Obese Caucasian Patients with Septic Shock According to the Sepsis-3 Definition. Journal of Clinical Medicine, 2020, 9, 46.	1.0	13
4	TIM-3 Genetic Variants Are Associated with Altered Clinical Outcome and Susceptibility to Gram-Positive Infections in Patients with Sepsis. International Journal of Molecular Sciences, 2020, 21, 8318.	1.8	8
5	Results of concomitant groin-free percutaneous temporary RVAD support using a centrifugal pump with a double-lumen jugular venous cannula in LVAD patients. Journal of Thoracic Disease, 2019, 11, S913-S920.	0.6	27
6	Lack of an Association between the Functional Polymorphism TREM-1 rs2234237 and the Clinical Course of Sepsis among Critically Ill Caucasian Patients—A Monocentric Prospective Genetic Association Study. Journal of Clinical Medicine, 2019, 8, 301.	1.0	8
7	CTLA-4 Genetic Variants Predict Survival in Patients with Sepsis. Journal of Clinical Medicine, 2019, 8, 70.	1.0	13
8	Paths of femoral nerve catheters placed using ultrasound-guided in plane vs out of plane techniques. Medicine (United States), 2018, 97, e12958.	0.4	3
9	Anaemia requiring red blood cell transfusion is associated with unfavourable 90-day survival in surgical patients with sepsis. BMC Research Notes, 2018, 11, 879.	0.6	6
10	The CTLA-4 rs231775 GG genotype is associated with favorable 90-day survival in Caucasian patients with sepsis. Scientific Reports, 2018, 8, 15140.	1.6	13
11	Prehospital ultrasound-guided nerve blocks improve reduction-feasibility of dislocated extremity injuries compared to systemic analgesia. A randomized controlled trial. PLoS ONE, 2018, 13, e0199776.	1.1	21
12	Pre-hospital transthoracic echocardiography for early identification of non-ST-elevation myocardial infarction in patients with acute coronary syndrome. Critical Care, 2018, 22, 29.	2.5	13
13	Prognostic relevance of hemodialysis for short-term survival in patients after LVAD implantation. Scientific Reports, 2018, 8, 8546.	1.6	7
14	Combination of general anesthesia and peripheral nerve block with low-dose ropivacaine reduces postoperative pain for several days after outpatient arthroscopy. Medicine (United States), 2017, 96, e6046.	0.4	8
15	The FER rs4957796 TT genotype is associated with unfavorable 90-day survival in Caucasian patients with severe ARDS due to pneumonia. Scientific Reports, 2017, 7, 9887.	1.6	18
16	Extracorporeal life support with left ventricular decompression—improved survival in severe cardiogenic shock: results from a retrospective study. PeerJ, 2017, 5, e3813.	0.9	29
17	The CD14 rs2569190 TT Genotype Is Associated with an Improved 30-Day Survival in Patients with Sepsis: A Prospective Observational Cohort Study. PLoS ONE, 2015, 10, e0127761.	1.1	16
18	Chronic kidney disease is associated with a higher 90-day mortality than other chronic medical conditions in patients with sepsis. Scientific Reports, 2015, 5, 10539.	1.6	27

#	Article	IF	CITATION
19	Perioperative Blood Glucose Levels <150 mg/dL are Associated With Improved 5-Year Survival in Patients Undergoing On-Pump Cardiac Surgery. Medicine (United States), 2015, 94, e2035.	0.4	9
20	Primary bacteraemia is associated with a higher mortality risk compared with pulmonary and intra-abdominal infections in patients with sepsis: a prospective observational cohort study. BMJ Open, 2015, 5, e006616-e006616.	0.8	21
21	Impact of statin therapy on mortality in patients with sepsis-associated acute respiratory distress syndrome (ARDS) depends on ARDS severity: a prospective observational cohort study. BMC Medicine, 2015, 13, 128.	2.3	57
22	Genetic Polymorphisms in Endothelin-1 as Predictors for Long-Term Survival and the Cardiac Index in Patients Undergoing On-Pump Cardiac Surgery. PLoS ONE, 2015, 10, e0131155.	1.1	3
23	The regulatory toll-like receptor 4 genetic polymorphism rs11536889 is associated with renal, coagulation and hepatic organ failure in sepsis patients. Journal of Translational Medicine, 2014, 12, 177.	1.8	25
24	Ninety-Day Survival Rate of Patients with Sepsis Relates to Programmed Cell Death 1 Genetic Polymorphism rs11568821. Journal of Investigative Medicine, 2014, 62, 638-643.	0.7	21