## Yoon-Kyung Chang

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/2559866/publications.pdf

Version: 2024-02-01

43 papers

835 citations

686830 13 h-index 26 g-index

43 all docs 43 docs citations

43 times ranked 1109 citing authors

#	Article	IF	CITATIONS
1	Rescue Stenting for Failed Mechanical Thrombectomy in Acute Ischemic Stroke. Stroke, 2018, 49, 958-964.	1.0	135
2	Moderate-to-severe obstructive sleep apnea is associated with cerebral small vessel disease. Sleep Medicine, 2017, 30, 36-42.	0.8	75
3	Improved oral hygiene care is associated with decreased risk of occurrence for atrial fibrillation and heart failure: A nationwide population-based cohort study. European Journal of Preventive Cardiology, 2020, 27, 1835-1845.	0.8	73
4	Improved oral hygiene is associated with decreased risk of new-onset diabetes: a nationwide population-based cohort study. Diabetologia, 2020, 63, 924-933.	2.9	67
5	Better oral hygiene is associated with lower risk of stroke. Journal of Periodontology, 2021, 92, 87-94.	1.7	45
6	Oral health and gastrointestinal cancer: A nationwide cohort study. Journal of Clinical Periodontology, 2020, 47, 796-808.	2.3	42
7	Benefit of Targeting a LDL (Low-Density Lipoprotein) Cholesterol <70 mg/dL During 5 Years After Ischemic Stroke. Stroke, 2020, 51, 1231-1239.	1.0	39
8	Low levels of plasma omega 3-polyunsaturated fatty acids are associated with cerebral small vessel diseases in acute ischemic stroke patients. Nutrition Research, 2015, 35, 368-374.	1.3	31
9	Microbiota dysbiosis and functional outcome in acute ischemic stroke patients. Scientific Reports, 2021, 11, 10977.	1.6	28
10	Insulin resistance is associated with poor functional outcome after acute ischemic stroke in non-diabetic patients. Scientific Reports, 2021, 11, 1229.	1.6	22
11	Low Plasma Proportion of Omega 3-Polyunsaturated Fatty Acids Predicts Poor Outcome in Acute Non-Cardiogenic Ischemic Stroke Patients. Journal of Stroke, 2015, 17, 168.	1.4	19
12	Plasma Klotho concentration is associated with the presence, burden and progression of cerebral small vessel disease in patients with acute ischaemic stroke. PLoS ONE, 2019, 14, e0220796.	1.1	17
13	Interarm Blood Pressure Difference is Associated with Early Neurological Deterioration, Poor Short-Term Functional Outcome, and Mortality in Noncardioembolic Stroke Patients. Journal of		

#	Article	lF	CITATIONS
19	Interarm Systolic and Diastolic Blood Pressure Difference Is Diversely Associated With Cerebral Atherosclerosis in Noncardioembolic Stroke Patients. American Journal of Hypertension, 2018, 31, 35-42.	1.0	13
20	Association of Alzheimer's Disease with COVID-19 Susceptibility and Severe Complications: A Nationwide Cohort Study. Journal of Alzheimer's Disease, 2022, 87, 701-710.	1.2	13
21	Association between the fatty liver index and the risk of severe complications in COVID-19 patients: a nationwide retrospective cohort study. BMC Infectious Diseases, 2022, 22, 384.	1.3	13
22	High Dietary Glycemic Load is Associated with Poor Functional Outcome in Patients with Acute		

#	Article	lF	CITATIONS
37	Association between intakes of minerals (potassium, magnesium, and calcium) and diet quality and risk of cerebral atherosclerosis in ischemic stroke patients. Journal of Nutrition and Health, 2015, 48, 167. Association of Periodontal Disease with the Occurrence of Unruptured Cerebral Aneurysm among	0.2	2
38	Adults in Korea: A Nationwide Population-Based Cohort Study     TRANSLATE with x   English      Arabic  Hebrew  Polish     Bulgarian  Hindi  Portuguese    Catalan  Hmong Daw  Romanian    Chinese Simplified	0.8	2
39	Hungarian&#xÓD; Russian&#xOD; &#xOD; &#xOD; Chinese. Medicina (Lithuania), 2021, 57, 910.  Plasma osteoprotegerin levels are associated with the presence and burden of cerebral small vessel disease in patients with acute ischemic stroke. Clinical Neurology and Neurosurgery, 2021, 210, 107010.</td><td>0.6</td><td>2</td></tr><tr><td>40</td><td>The association of plasma osteoprotegerin levels and functional outcomes post endovascular thrombectomy in acute ischemic stroke patients: a retrospective observational study. PeerJ, 2022, 10, e13327.</td><td>0.9</td><td>1</td></tr><tr><td>41</td><td>Plasma osteoprotegerin level is associated with hemorrhagic transformation in stroke patients who underwent endovascular thrombectomy. Clinical Neurology and Neurosurgery, 2022, 219, 107305.</td><td>0.6</td><td>1</td></tr><tr><td>42</td><td>Characteristics for Ischemic Stroke in 18–30 Years Old Patients, Multicenter Stroke Registry Study. The Ewha Medical Journal, 2017, 40, 128.</td><td>0.1</td><td>0</td></tr><tr><td>43</td><td>Prognostic significance of acute hypertensive response in ischemic stroke or transient ischemic attack patients and its relationship with left ventricular myocardial function. Journal of Hypertension, 2019, 37, 2345-2353.</td><td>0.3</td><td>0</td></tr></tbody></table>		