Yoon-Kyung Chang

List of Publications by Year in Descending Order

Source: https://exaly.com/author-pdf/2559866/yoon-kyung-chang-publications-by-year.pdf

Version: 2024-04-20

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

39	417	11	19
papers	citations	h-index	g-index
43	619 ext. citations	3.7	4.04
ext. papers		avg, IF	L-index

#	Paper	IF	Citations
39	Association of Alzheimerは Disease with COVID-19 Susceptibility and Severe Complications: A Nationwide Cohort Study <i>Journal of Alzheimerps Disease</i> , 2022 ,	4.3	2
38	Clinical Implications of Atrial Fibrillation Detection Using Wearable Devices in Patients With Cryptogenic Stroke (CANDLE-AF) Trial: Design and Rationale <i>Frontiers in Cardiovascular Medicine</i> , 2022 , 9, 837958	5.4	1
37	Association between the fatty liver index and the risk of severe complications in COVID-19 patients: a nationwide retrospective cohort study <i>BMC Infectious Diseases</i> , 2022 , 22, 384	4	1
36	The association of plasma osteoprotegerin levels and functional outcomes post endovascular thrombectomy in acute ischemic stroke patients: a retrospective observational study <i>PeerJ</i> , 2022 , 10, e13327	3.1	
35	Plasma osteoprotegerin level is associated with hemorrhagic transformation in stroke patients who underwent endovascular thrombectomy. <i>Clinical Neurology and Neurosurgery</i> , 2022 , 219, 107305	2	
34	Association between Findings in Oral Health Screening and Body Mass Index: A Nation-Wide Longitudinal Study. <i>International Journal of Environmental Research and Public Health</i> , 2021 , 18,	4.6	2
33	Plasma osteoprotegerin levels are associated with the presence and burden of cerebral small vessel disease in patients with acute ischemic stroke. <i>Clinical Neurology and Neurosurgery</i> , 2021 , 210, 107010	2	1
32	Microbiota dysbiosis and functional outcome in acute ischemic stroke patients. <i>Scientific Reports</i> , 2021 , 11, 10977	4.9	5
31	Tooth loss is associated with an increased risk of hypertension: A nationwide population-based cohort study. <i>PLoS ONE</i> , 2021 , 16, e0253257	3.7	1
30	Oral health and longitudinal changes in fasting glucose levels: A nationwide cohort study. <i>PLoS ONE</i> , 2021 , 16, e0253769	3.7	3
29	Better oral hygiene is associated with lower risk of stroke. <i>Journal of Periodontology</i> , 2021 , 92, 87-94	4.6	19
28	Insulin resistance is associated with poor functional outcome after acute ischemic stroke in non-diabetic patients. <i>Scientific Reports</i> , 2021 , 11, 1229	4.9	8
27	Outcome in Patients Treated with Intra-arterial thrombectomy: The optiMAL Blood Pressure control (OPTIMAL-BP) Trial. <i>International Journal of Stroke</i> , 2021 , 17474930211041213	6.3	O
26	Improved oral hygiene care and chronic kidney disease occurrence: A nationwide population-based retrospective cohort study <i>Medicine (United States)</i> , 2021 , 100, e27845	1.8	1
25	Oral health and gastrointestinal cancer: A nationwide cohort study. <i>Journal of Clinical Periodontology</i> , 2020 , 47, 796-808	7.7	13
24	Improved oral hygiene is associated with decreased risk of new-onset diabetes: a nationwide population-based cohort study. <i>Diabetologia</i> , 2020 , 63, 924-933	10.3	24
23	Benefit of Targeting a LDL (Low-Density Lipoprotein) Cholesterol . <i>Stroke</i> , 2020 , 51, 1231-1239	6.7	19

(2016-2020)

22	Plasma Fibroblast Growth Factor 23 Concentration Is Associated with Intracranial Cerebral Atherosclerosis in Acute Ischemic Stroke Patients. <i>Journal of Clinical Neurology (Korea</i> , 2020 , 16, 29-36	1.7	4
21	Improved oral hygiene care is associated with decreased risk of occurrence for atrial fibrillation and heart failure: A nationwide population-based cohort study. <i>European Journal of Preventive Cardiology</i> , 2020 , 27, 1835-1845	3.9	29
20	Association of Tooth Loss with New-Onset Parkinson'd Disease: A Nationwide Population-Based Cohort Study. <i>Parkinson's Disease</i> , 2020 , 2020, 4760512	2.6	5
19	Inter-arm Blood Pressure Difference is Associated with Recurrent Stroke in Non-cardioembolic Stroke Patients. <i>Scientific Reports</i> , 2019 , 9, 12758	4.9	5
18	Plasma Klotho concentrations predict functional outcome at three months after acute ischemic stroke patients. <i>Annals of Medicine</i> , 2019 , 51, 262-269	1.5	6
17	Interarm Blood Pressure Difference has Various Associations with the Presence and Burden of Cerebral Small-Vessel Diseases in Noncardioembolic Stroke Patients. <i>Journal of Clinical Neurology</i> (Korea, 2019 , 15, 159-167	1.7	6
16	Plasma Klotho concentration is associated with the presence, burden and progression of cerebral small vessel disease in patients with acute ischaemic stroke. <i>PLoS ONE</i> , 2019 , 14, e0220796	3.7	9
15	Prognostic significance of acute hypertensive response in ischemic stroke or transient ischemic attack patients and its relationship with left ventricular myocardial function. <i>Journal of Hypertension</i> , 2019 , 37, 2345-2353	1.9	
14	High dietary glycemic load was associated with the presence and burden of cerebral small vessel diseases in acute ischemic stroke patients. <i>Nutrition Research</i> , 2018 , 51, 93-101	4	9
13	Rescue Stenting for Failed Mechanical Thrombectomy in Acute Ischemic Stroke: A Multicenter Experience. <i>Stroke</i> , 2018 , 49, 958-964	6.7	85
12	Increased Thrombogenicity in Chronic Renal Failure in a Rat Model Induced by 5/6 Ablation/Infarction. <i>Yonsei Medical Journal</i> , 2018 , 59, 754-759	3	2
11	High Dietary Glycemic Load is Associated with Poor Functional Outcome in Patients with Acute Cerebral Infarction. <i>Journal of Clinical Neurology (Korea</i> , 2018 , 14, 165-173	1.7	10
10	Interarm Blood Pressure Difference is Associated with Early Neurological Deterioration, Poor Short-Term Functional Outcome, and Mortality in Noncardioembolic Stroke Patients. <i>Journal of Clinical Neurology (Korea</i> , 2018 , 14, 555-565	1.7	15
9	Moderate-to-severe obstructive sleep apnea is associated with cerebral small vessel disease. <i>Sleep Medicine</i> , 2017 , 30, 36-42	4.6	51
8	Is obstructive sleep apnea associated with the presence of intracranial cerebral atherosclerosis?. <i>Sleep and Breathing</i> , 2017 , 21, 639-646	3.1	11
7	Characteristics for Ischemic Stroke in 18B0 Years Old Patients, Multicenter Stroke Registry Study. <i>The Ewha Medical Journal</i> , 2017 , 40, 128	0.1	
6	Interarm Systolic and Diastolic Blood Pressure Difference Is Diversely Associated With Cerebral Atherosclerosis in Noncardioembolic Stroke Patients. <i>American Journal of Hypertension</i> , 2017 , 31, 35-42	2.3	12
5	Recurrent Episodes of Rhabdomyolysis after Seizures in a Patient with Glycogen Storage Disease Type V. <i>Journal of Clinical Neurology (Korea</i> , 2016 , 12, 373-5	1.7	3

4	Low levels of plasma omega 3-polyunsaturated fatty acids are associated with cerebral small vessel diseases in acute ischemic stroke patients. <i>Nutrition Research</i> , 2015 , 35, 368-74	4	26
3	Low-density-lipoprotein particle size predicts a poor outcome in patients with atherothrombotic stroke. <i>Journal of Clinical Neurology (Korea</i> , 2015 , 11, 80-6	1.7	9
2	Association between intakes of minerals (potassium, magnesium, and calcium) and diet quality and risk of cerebral atherosclerosis in ischemic stroke patients. <i>Journal of Nutrition and Health</i> , 2015 , 48, 167	0.8	2
1	Low Plasma Proportion of Omega 3-Polyunsaturated Fatty Acids Predicts Poor Outcome in Acute Non-Cardiogenic Ischemic Stroke Patients. <i>Journal of Stroke</i> , 2015 , 17, 168-76	5.6	18