

Erik PÅ¥lsson

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2559926/publications.pdf>

Version: 2024-02-01

44
papers

1,270
citations

331538

21
h-index

395590

33
g-index

46
all docs

46
docs citations

46
times ranked

2280
citing authors

#	ARTICLE	IF	CITATIONS
1	A serum proteomic study of two case-control cohorts identifies novel biomarkers for bipolar disorder. <i>Translational Psychiatry</i> , 2022, 12, 55.	2.4	7
2	Influence of genetic variations in IL1B on brain region volumes in bipolar patients and controls. <i>Psychiatry Research</i> , 2021, 296, 113606.	1.7	4
3	Association of CACNA1C polymorphisms with serum BDNF levels in bipolar disorder. <i>British Journal of Psychiatry</i> , 2021, 218, 77-79.	1.7	11
4	Genome-wide association study of patients with a severe major depressive episode treated with electroconvulsive therapy. <i>Molecular Psychiatry</i> , 2021, 26, 2429-2439.	4.1	32
5	Central levels of tryptophan metabolites in subjects with bipolar disorder. <i>European Neuropsychopharmacology</i> , 2021, 43, 52-62.	0.3	24
6	Patient educational level and management of bipolar disorder. <i>BJPsych Open</i> , 2021, 7, e63.	0.3	6
7	Long-term trajectory of cognitive performance in people with bipolar disorder and controls: 6-year longitudinal study. <i>BJPsych Open</i> , 2021, 7, e115.	0.3	8
8	Regional lithium prescription rates and recurrence in bipolar disorder. <i>International Journal of Bipolar Disorders</i> , 2021, 9, 18.	0.8	8
9	Serum profiling of anorexia nervosa: A 1H NMR-based metabolomics study. <i>European Neuropsychopharmacology</i> , 2021, 49, 1-10.	0.3	6
10	Characterisation of age and polarity at onset in bipolar disorder. <i>British Journal of Psychiatry</i> , 2021, 219, 659-669.	1.7	20
11	Cerebrospinal fluid proteomics targeted for central nervous system processes in bipolar disorder. <i>Molecular Psychiatry</i> , 2021, 26, 7446-7453.	4.1	5
12	Psychoeducation for bipolar disorder and risk of recurrence and hospitalization – a within-individual analysis using registry data. <i>Psychological Medicine</i> , 2020, 50, 1043-1049.	2.7	19
13	Characteristics of bipolar I and II disorder: A study of 8766 individuals. <i>Bipolar Disorders</i> , 2020, 22, 392-400.	1.1	52
14	International Consortium on the Genetics of Electroconvulsive Therapy and Severe Depressive Disorders (Gen-ECT-ic). <i>European Archives of Psychiatry and Clinical Neuroscience</i> , 2020, 270, 921-932.	1.8	22
15	<i>CACNA1C</i> polymorphism and brain cortical structure in bipolar disorder. <i>Journal of Psychiatry and Neuroscience</i> , 2020, 45, 182-187.	1.4	10
16	Long-term subjective memory after electroconvulsive therapy. <i>BJPsych Open</i> , 2020, 6, e26.	0.3	23
17	Executive functioning but not IQ or illness severity predicts occupational status in bipolar disorder. <i>International Journal of Bipolar Disorders</i> , 2020, 8, 7.	0.8	16
18	Prospective cohort study of early biosignatures of response to lithium in bipolar-I-disorders: overview of the H2020-funded R-LiNK initiative. <i>International Journal of Bipolar Disorders</i> , 2019, 7, 20.	0.8	41

#	ARTICLE	IF	CITATIONS
19	Classification of cognitive performance in bipolar disorder. <i>Cognitive Neuropsychiatry</i> , 2017, 22, 407-421.	0.7	12
20	Cerebrospinal fluid monoamine metabolite profiles in bipolar disorder, ADHD, and controls. <i>Journal of Neural Transmission</i> , 2017, 124, 1135-1143.	1.4	19
21	Personality traits in bipolar disorder and influence on outcome. <i>BMC Psychiatry</i> , 2017, 17, 159.	1.1	32
22	Blood metabolomics analysis identifies abnormalities in the citric acid cycle, urea cycle, and amino acid metabolism in bipolar disorder. <i>BBA Clinical</i> , 2016, 5, 151-158.	4.1	76
23	Polymorphisms of <i>BDNF</i> and <i>CACNA1C</i> are not associated with cognitive functioning in bipolar disorder or healthy controls. <i>Cognitive Neuropsychiatry</i> , 2016, 21, 271-278.	0.7	14
24	Dietary intake, resting energy expenditure, and eating behavior in women with and without polycystic ovary syndrome. <i>Clinical Nutrition</i> , 2016, 35, 213-218.	2.3	54
25	<i>CACNA1C</i> polymorphism and altered phosphorylation of tau in bipolar disorder. <i>British Journal of Psychiatry</i> , 2016, 208, 195-196.	1.7	19
26	Polymorphisms of dopamine pathway genes <i>NRG1</i> and <i>LMX1A</i> are associated with cognitive performance in bipolar disorder. <i>Bipolar Disorders</i> , 2015, 17, 859-868.	1.1	23
27	Cognitive Performance and Cerebrospinal Fluid Biomarkers of Neurodegeneration: A Study of Patients with Bipolar Disorder and Healthy Controls. <i>PLoS ONE</i> , 2015, 10, e0127100.	1.1	38
28	Monocyte and microglial activation in patients with mood-stabilized bipolar disorder. <i>Journal of Psychiatry and Neuroscience</i> , 2015, 40, 250-258.	1.4	75
29	CSF neuroinflammatory biomarkers in bipolar disorder are associated with cognitive impairment. <i>European Neuropsychopharmacology</i> , 2015, 25, 1091-1098.	0.3	47
30	Increased brain nitric oxide levels following ethanol administration. <i>Nitric Oxide - Biology and Chemistry</i> , 2015, 47, 52-57.	1.2	22
31	Increased cerebrospinal fluid interleukin-8 in bipolar disorder patients associated with lithium and antipsychotic treatment. <i>Brain, Behavior, and Immunity</i> , 2015, 43, 198-204.	2.0	51
32	Markers of glutamate signaling in cerebrospinal fluid and serum from patients with bipolar disorder and healthy controls. <i>European Neuropsychopharmacology</i> , 2015, 25, 133-140.	0.3	38
33	Cognitive Functioning in Clinically Stable Patients with Bipolar Disorder I and II. <i>PLoS ONE</i> , 2015, 10, e0115562.	1.1	36
34	Low neuropeptide Y in cerebrospinal fluid in bipolar patients is associated with previous and prospective suicide attempts. <i>European Neuropsychopharmacology</i> , 2014, 24, 1907-1915.	0.3	24
35	Apolipoprotein E Genotype and the Diagnostic Accuracy of Cerebrospinal Fluid Biomarkers for Alzheimer Disease. <i>JAMA Psychiatry</i> , 2014, 71, 1183.	6.0	85
36	Abnormality in serum levels of mature brain-derived neurotrophic factor (BDNF) and its precursor proBDNF in mood-stabilized patients with bipolar disorder: A study of two independent cohorts. <i>Journal of Affective Disorders</i> , 2014, 160, 1-9.	2.0	78

#	ARTICLE	IF	CITATIONS
37	Bloodâ€“cerebrospinal fluid barrier dysfunction in patients with bipolar disorder in relation to antipsychotic treatment. <i>Psychiatry Research</i> , 2014, 217, 143-146.	1.7	36
38	Neurocognitive function in bipolar disorder: a comparison between bipolar I and II disorder and matched controls. <i>BMC Psychiatry</i> , 2013, 13, 165.	1.1	64
39	Brain nitric oxide: Regional characterisation of a real-time microelectrochemical sensor. <i>Journal of Neuroscience Methods</i> , 2012, 209, 13-21.	1.3	25
40	Noise benefit in prepulse inhibition of the acoustic startle reflex. <i>Psychopharmacology</i> , 2011, 214, 675-685.	1.5	13
41	Information processing deficits and nitric oxide signalling in the phencyclidine model of schizophrenia. <i>Psychopharmacology</i> , 2010, 212, 643-651.	1.5	11
42	Increased cortical nitric oxide release after phencyclidine administration. <i>Synapse</i> , 2009, 63, 1083-1088.	0.6	22
43	Agmatine attenuates the disruptive effects of phencyclidine on prepulse inhibition. <i>European Journal of Pharmacology</i> , 2008, 590, 212-216.	1.7	27
44	The amino acid l-lysine blocks the disruptive effect of phencyclidine on prepulse inhibition in mice. <i>Psychopharmacology</i> , 2007, 192, 9-15.	1.5	13