
## **Byung-Hoon Lee**

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/2561415/publications.pdf Version: 2024-02-01



RVUNC-HOON LEE

#	Article	IF	CITATIONS
1	SNX10-mediated degradation of LAMP2A by NSAIDs inhibits chaperone-mediated autophagy and induces hepatic lipid accumulation. Theranostics, 2022, 12, 2351-2369.	10.0	8
2	The role of SHMT2 in modulating lipid metabolism in hepatocytes via glycine-mediated mTOR activation. Amino Acids, 2022, 54, 823-834.	2.7	5
3	CD44 is involved in liver regeneration through enhanced uptake of extracellular cystine. Clinical and Translational Medicine, 2022, 12, e873.	4.0	1
4	Small-Molecule Inhibitors Targeting Proteasome-Associated Deubiquitinases. International Journal of Molecular Sciences, 2021, 22, 6213.	4.1	12
5	Downregulation of PHGDH expression and hepatic serine level contribute to the development of fatty liver disease. Metabolism: Clinical and Experimental, 2020, 102, 154000.	3.4	31
6	Diclofenac impairs autophagic flux via oxidative stress and lysosomal dysfunction: Implications for hepatotoxicity. Redox Biology, 2020, 37, 101751.	9.0	49
7	Deubiquitination Reactions on the Proteasome for Proteasome Versatility. International Journal of Molecular Sciences, 2020, 21, 5312.	4.1	24
8	Protective effect of EX-527 against high-fat diet-induced diabetic nephropathy in Zucker rats. Toxicology and Applied Pharmacology, 2020, 390, 114899.	2.8	18
9	Z-ligustilide and n-Butylidenephthalide Isolated from the Aerial Parts of Angelica tenuissima Inhibit Lipid Accumulation In Vitro and In Vivo. Planta Medica, 2019, 85, 719-728.	1.3	6
10	Hepatic upregulation of fetuin-A mediates acetaminophen-induced liver injury through activation of TLR4 in mice. Biochemical Pharmacology, 2019, 166, 46-55.	4.4	13
11	In vitro analysis of proteasome-associated USP14 activity for substrate degradation and deubiquitylation. Methods in Enzymology, 2019, 619, 249-268.	1.0	3
12	Activation of SIRT1 by l-serine increases fatty acid oxidation and reverses insulin resistance in C2C12 myotubes (l-serine activates SIRT1 in C2C12 myotubes). Cell Biology and Toxicology, 2019, 35, 457-470.	5.3	19
13	The deubiquitinating enzyme Usp14 controls ciliogenesis and Hedgehog signaling. Human Molecular Genetics, 2019, 28, 764-777.	2.9	25
14	The critical role of autophagy in cadmium-induced immunosuppression regulated by endoplasmic reticulum stress-mediated calpain activation in RAW264.7 mouse monocytes. Toxicology, 2018, 393, 15-25.	4.2	39
15	Chemically Induced Cellular Proteolysis: An Emerging Therapeutic Strategy for Undruggable Targets. Molecules and Cells, 2018, 41, 933-942.	2.6	30
16	Ubiquitylation of p62/sequestosome1 activates its autophagy receptor function and controls selective autophagy upon ubiquitin stress. Cell Research, 2017, 27, 657-674.	12.0	143
17	Activation of AMPK by berberine induces hepatic lipid accumulation by upregulation of fatty acid translocase CD36 in mice. Toxicology and Applied Pharmacology, 2017, 316, 74-82.	2.8	45
18	An inhibitor of the proteasomal deubiquitinating enzyme USP14 induces tau elimination in cultured neurons. Journal of Biological Chemistry, 2017, 292, 19209-19225.	3.4	98

**BYUNG-HOON LEE** 

#	Article	IF	CITATIONS
19	Inhibition of homocysteine-induced endoplasmic reticulum stress and endothelial cell damage by I-serine and glycine. Toxicology in Vitro, 2016, 34, 138-145.	2.4	19
20	Guidelines for the use and interpretation of assays for monitoring autophagy (3rd edition). Autophagy, 2016, 12, 1-222.	9.1	4,701
21	Orotic Acid Induces Hypertension Associated with Impaired Endothelial Nitric Oxide Synthesis. Toxicological Sciences, 2015, 144, 307-317.	3.1	13
22	Increased Hepatic Fatty Acid Uptake and Esterification Contribute to Tetracycline-Induced Steatosis in Mice. Toxicological Sciences, 2015, 145, 273-282.	3.1	27
23	Cinnamamides, Novel Liver X Receptor Antagonists that Inhibit Ligand-Induced Lipogenesis and Fatty Liver. Journal of Pharmacology and Experimental Therapeutics, 2015, 355, 362-369.	2.5	13
24	Differential effects of p38 and JNK activation by GSK3 on cadmium-induced autophagy and apoptosis. Toxicology Research, 2015, 4, 976-985.	2.1	4
25	L-Serine Supplementation Attenuates Alcoholic Fatty Liver by Enhancing Homocysteine Metabolism in Mice and Rats. Journal of Nutrition, 2015, 145, 260-267.	2.9	57
26	Expression of CYP3A in chronic ethanol-fed mice is mediated by endogenous pregnane X receptor ligands formed by enhanced cholesterol metabolism. Archives of Toxicology, 2015, 89, 579-589.	4.2	10
27	Abstract 479: Orotic Acid Induces Hypertension Associated With Impaired Endothelial Nitric Oxide Synthesis. Arteriosclerosis, Thrombosis, and Vascular Biology, 2015, 35, .	2.4	Ο
28	Uric acid induces endothelial dysfunction by vascular insulin resistance associated with the impairment of nitric oxide synthesis. FASEB Journal, 2014, 28, 3197-3204.	0.5	164
29	LXR-α antagonist meso-dihydroguaiaretic acid attenuates high-fat diet-induced nonalcoholic fatty liver. Biochemical Pharmacology, 2014, 90, 414-424.	4.4	32
30	Protective effect of resveratrol derivatives on high-fat diet induced fatty liver by activating AMP-activated protein kinase. Archives of Pharmacal Research, 2014, 37, 1169-1176.	6.3	41
31	Uric acid induces fat accumulation via generation of endoplasmic reticulum stress and SREBP-1c activation in hepatocytes. Laboratory Investigation, 2014, 94, 1114-1125.	3.7	196
32	Activation of Autophagy Rescues Amiodarone-Induced Apoptosis of Lung Epithelial Cells and Pulmonary Toxicity in Rats. Toxicological Sciences, 2013, 136, 193-204.	3.1	20
33	Involvement of E2F1 transcriptional activity in cadmiumâ€induced cellâ€cycle arrest at G1 in human lung fibroblasts. Environmental and Molecular Mutagenesis, 2011, 52, 145-152.	2.2	14
34	Gas chromatography–mass spectrometry-based simultaneous quantitative analytical method for urinary oxysterols and bile acids in rats. Analytical Biochemistry, 2011, 408, 242-252.	2.4	48
35	Role of the AMPK/SREBP-1 pathway in the development of orotic acid-induced fatty liver. Journal of Lipid Research, 2011, 52, 1617-1625.	4.2	82
36	Genomics-based screening of differentially expressed genes in the brains of mice exposed to silver nanoparticles via inhalation. Journal of Nanoparticle Research, 2010, 12, 1567-1578.	1.9	74

**BYUNG-HOON LEE** 

#	Article	IF	CITATIONS
37	Analysis of hepatic gene expression during fatty liver change due to chronic ethanol administration in mice. Toxicology and Applied Pharmacology, 2009, 235, 312-320.	2.8	22
38	Salvia miltiorrhiza Bunge and its active component cryptotanshinone protects primary cultured rat hepatocytes from acute ethanol-induced cytotoxicity and fatty infiltration. Food and Chemical Toxicology, 2009, 47, 98-103.	3.6	32
39	Magnolia officinalis Reverses Alcoholic Fatty Liver by Inhibiting the Maturation of Sterol Regulatory Element–Binding Protein-1c. Journal of Pharmacological Sciences, 2009, 109, 486-495.	2.5	31
40	Effects of tanshinone IIA on the hepatotoxicity and gene expression involved in alcoholic liver disease. Archives of Pharmacal Research, 2008, 31, 659-665.	6.3	34
41	Gene expression profiling in human lung fibroblast following cadmium exposure. Food and Chemical Toxicology, 2008, 46, 1131-1137.	3.6	29
42	Elevated Levels of PDGF Receptor and MDM2 as Potential Biomarkers for Formaldehyde Intoxication. Toxicological Research, 2008, 24, 45-49.	2.1	2
43	Temporal Changes in the Hepatic Fatty Liver in Mice Receiving Standard Lieber-DeCarli Diet. Toxicological Research, 2008, 24, 113-117.	2.1	5
44	Changes in the Expression of Ras-family Genes in Rats Exposed to Formaldehyde by Inhalation. Toxicological Research, 2008, 24, 201-206.	2.1	1
45	Differential gene expression and lipid metabolism in fatty liver induced by acute ethanol treatment in mice. Toxicology and Applied Pharmacology, 2007, 223, 225-233.	2.8	87
46	Hepatic Gene Expression Profiling and Lipid Homeostasis in Mice Exposed to Steatogenic Drug, Tetracycline. Toxicological Sciences, 2006, 94, 206-216.	3.1	57
47	Salvia miltiorrhiza Inhibits Biliary Obstruction-Induced Hepatocyte Apoptosis by Cytoplasmic Sequestration of p53. Toxicology and Applied Pharmacology, 2002, 182, 27-33.	2.8	31
48	Mitochondrial NADH-cytochrome b5 reductase plays a crucial role in the reduction of d-erythroascorbyl free radical in Saccharomyces cerevisiae. Biochimica Et Biophysica Acta - General Subjects, 2001, 1527, 31-38.	2.4	33
49	Cytotoxic Lavandulyl Flavanones from Sophora flavescens. Journal of Natural Products, 2000, 63, 680-681.	3.0	128
50	Manganese-containing superoxide dismutase and its gene from Candida albicans. Biochimica Et Biophysica Acta - General Subjects, 1999, 1426, 409-419.	2.4	35
51	D-Arabinose dehydrogenase and its gene from Saccharomyces cerevisiae. BBA - Proteins and Proteomics, 1998, 1429, 29-39.	2.1	60
52	D-Erythroascorbic acid is an important antioxidant molecule inSaccharomyces cerevisiae. Molecular Microbiology, 1998, 30, 895-903.	2.5	105